

Random Knot Diagrams

Jason Cantarella (UGA)

joint w/ Harrison Chapman (UGA), Matt Mastin (Wake Forest)


Crucial Assist: Eric Rawdon (St. Thomas)

CanadAM Conference, June 2, 2015

Natural questions about knot diagrams

Question

What fraction of 8-crossing diagrams are trefoils?




Natural questions about knot diagrams

Question

What fraction of 8-crossing diagrams are trefoils?


12.48%



Natural questions about knot diagrams

Question


What is the average minimal crossing # of an 8-crossing diagram?



Natural questions about knot diagrams

Question


What is the average minimal crossing # of an 8-crossing diagram?
0.52



Natural questions about knot diagrams

Definition


The **untwisting** operator deletes all 1-crossing connect summands of a diagram. (Equivalently, performs all “available” Reidemeister I moves.)



Natural questions about knot diagrams

Question


What is the average crossing # of a untwisted 8-crossing diagram?



Natural questions about knot diagrams

Question


What is the average crossing # of a untwisted 8-crossing diagram?
2.20



Natural questions about knot diagrams

Question

How many 8-crossing diagrams can be untwisted to the unknot?




Natural questions about knot diagrams


Question

How many 8-crossing diagrams can be untwisted to the unknot?


42.05%




Ansatz




Combinatorial approaches



Polymers




Random curves




Knots




Random
combinatorial




???



Polymers




Random curves




Knots



Random
diagrams




Random
combinatorial



Random diagrams

Definition


In the **random diagram model** of random knotting, a n -crossing diagram is drawn uniformly from the finite set of n -crossing knot diagrams.



Random diagrams

Definition

A **knot diagram** is a equivalence class of generic immersions of the oriented S^1 into the sphere S^2 together with over-under strand information at each double point up to diffeomorphism of S^2 .



How to enumerate knot diagrams (like a topologist)

Definition

A **knot shadow** is a equivalence class of generic immersions of the unoriented S^1 into the sphere S^2 up to diffeomorphism of S^2 .

Plan to Enumerate Diagrams

- 1 *Enumerate shadows (and discard isomorphic shadows)*
- 2 *Assign crossing and orientation information (and discard crossing patterns related by an automorphism of the shadow)*

How to enumerate knot diagrams (like a topologist)

Definition

A **knot shadow** is a equivalence class of generic immersions of the unoriented S^1 into the sphere S^2 up to diffeomorphism of S^2 .

Plan to Enumerate Diagrams

- 1 *Enumerate shadows (and discard isomorphic shadows)*
- 2 *Assign crossing and orientation information (and discard crossing patterns related by an automorphism of the shadow)*


Observation (known to all combinatoricists, but new to me)

Symmetry stinks.

Tabulating knot shadows: plantri, two ways

Proposition

Knot shadows \leftrightarrow 1-component 4-valent embedded planar multigraphs up to embedded isomorphism




Tabulating knot shadows: plantri, two ways

Proposition

Knot shadows \leftrightarrow 1-component 4-valent embedded planar multigraphs up to embedded isomorphism

- 1 Add loops and edges to planar simple graphs (slow)




Tabulating knot shadows: plantri, two ways

Proposition

Knot shadows \leftrightarrow 1-component 4-valent embedded planar multigraphs up to embedded isomorphism

- 1 Add loops and edges to planar simple graphs (slow)
- 2 Generate multiquadrangulations of sphere by careful pattern of connect sums, take dual graphs (fast)




Tabulating knot shadows: plantri, two ways

Proposition

Knot shadows \leftrightarrow 1-component 4-valent embedded planar multigraphs up to embedded isomorphism

- 1 Add loops and edges to planar simple graphs (slow)
- 2 Generate multiquadrangulations of sphere by careful pattern of connect sums, take dual graphs (fast)



Actually generate all **link shadows**, then restrict to knot shadows

Verifying against existing shadow counts

oriented	$n = 0$	1	2	3	4	5
S^2, S^1	1	1	3	9	37	182
S^2	1	1	2	6	21	99
S^1	1	1	2	6	21	97
—	1	1	2	6	19	76

Curves on S^2 . The number of types

V.I. Arnol'd. *Topological Invariants of Plane Curves*

A008989 Number of immersions of unoriented circle into unoriented sphere with n double points.

1, 1, 2, 6, 19, 76, 376, 2194 [list](#); [graph](#); [rcf](#); [listen](#); [history](#); [text](#); [internal format](#)

OFFSET

0,3

REFERENCES

V. I. Arnold, Topological Invariants of Plane Curves..., American Math.

LINKS

[Table of \$n, a\(n\)\$ for \$n=0..7\$.](#)

CROSSREFS

Sequence in context: [A159119](#) [A181770](#) [A138800](#) * [A057240](#) [A079564](#) [A079453](#)

Adjacent sequences: [A008986](#) [A008987](#) [A008988](#) * [A008990](#) [A008991](#) [A008992](#)

KEYWORD

nonn

AUTHOR

[N. J. A. Sloane](#).

EXTENSIONS

Two more terms from Guy H. Valette (guy.valette(AT)skynet.be), Feb 09 20

STATUS

approved

OEIS A008989

n	# knot shadows
0	1
1	1
2	2
3	6
4	19
5	76
6	376
7	2194
8	14614
9	106421
10	823832

We have not found any existing counts of **diagrams**.

Assign crossings, orientation, identify

- 1 Orient each component. (2 choices)
- 2 Assign over-under information to each vertex. (2^n choices)

n	# knot shadows	2^{n+1} (# shadows)	# knot diagrams
3	6	96	36
4	19	608	276
5	76	4,864	2,936
6	376	48,128	35,872
7	2,194	561,664	484,088
8	14,614	7,482,368	6,967,942
9	106,421	108,975,104	105,555,336
10	823,832	1,687,207,936	$\sim 1,687,207,936$

Assign crossings, orientation, identify

- 1 Orient each component. (2 choices)
- 2 Assign over-under information to each vertex. (2^n choices)


n	# knot shadows	2^{n+1} (# shadows)	# knot diagrams
3	6	96	36
4	19	608	276
5	76	4,864	2,936
6	376	48,128	35,872
7	2,194	561,664	484,088
8	14,614	7,482,368	6,967,942
9	106,421	108,975,104	105,555,336
10	823,832	1,687,207,936	$\sim 1,687,207,936$

Observation (ktacbntm)


Symmetry becomes rare, quickly!

Size of the automorphism group of a random diagram


Mean number of automorphisms
versus crossing number of diagram




Knotting in diagrams




Ratios of knots in n -crossing diagrams (log scale)




Ratios of knots in n -crossing diagrams (log scale)



Why so many unknots?




The space of shadows




Link shadows. Pictures generated by Eric Lybrand (UGA undergrad).

The space of shadows




Link shadows. Pictures generated by Eric Lybrand (UGA undergrad).

The space of shadows



Link shadows. Pictures generated by Eric Lybrand (UGA undergrad).


The space of shadows



Link shadows. Pictures generated by Eric Lybrand (UGA undergrad).


Most diagrams are (very) composite

Number of diagrams with n connect summands
in 3 crossing diagrams




Most diagrams are (very) composite

Number of diagrams with n connect summands
in 4 crossing diagrams




Most diagrams are (very) composite

Number of diagrams with n connect summands
in 5 crossing diagrams




Most diagrams are (very) composite

Number of diagrams with n connect summands
in 6 crossing diagrams




Most diagrams are (very) composite

Number of diagrams with n connect summands
in 7 crossing diagrams




Most diagrams are (very) composite

Number of diagrams with n connect summands
in 8 crossing diagrams




Most diagrams are (very) composite

Number of diagrams with n connect summands
in 9 crossing diagrams




Most diagrams are (very) composite

Number of diagrams with n connect summands
in 10 crossing diagrams



Mean number of connect summands \simeq crossing number

Mean number of connect summands
versus crossing number of diagram



Obvious Unknots

Proposition

If $d(n, k)$ is the number of diagrams with n crossings and k connect summands, and $d(n)$ is the number of all n crossing diagrams, then the unknot fraction among all n crossing diagrams is at least

$$\frac{1}{d(n)} \left(d(n, n) + \frac{3}{4}d(n, n-2) + \frac{7}{8}d(n, n-3) \right)$$

Proof.

Any diagram in $d(n, n)$ is a connect sum of all 1-crossing diagrams, and so can be simplified to the unknot via RI moves.



Obvious Unknots

Proposition

If $d(n, k)$ is the number of diagrams with n crossings and k connect summands, and $d(n)$ is the number of all n crossing diagrams, then the unknot fraction among all n crossing diagrams is at least

$$\frac{1}{d(n)} \left(d(n, n) + \frac{3}{4}d(n, n-2) + \frac{7}{8}d(n, n-3) \right)$$

Proof.

Any diagram in $d(n, n-2)$ is a connect sum of 1-crossing diagrams, and a prime 3-crossing diagram (turns out there's only one— the trefoil diagram). This diagram is knotted iff those three crossings have the same sign, which occurs 1/4 of the time.



Obvious Unknots

Proposition

If $d(n, k)$ is the number of diagrams with n crossings and k connect summands, and $d(n)$ is the number of all n crossing diagrams, then the unknot fraction among all n crossing diagrams is at least

$$\frac{1}{d(n)} \left(d(n, n) + \frac{3}{4}d(n, n-2) + \frac{7}{8}d(n, n-3) \right)$$

Proof.

You can make a similar argument for $d(n, n-3)$.



Obvious Unknots

Proposition

If $d(n, k)$ is the number of diagrams with n crossings and k connect summands, and $d(n)$ is the number of all n crossing diagrams, then the unknot fraction among all n crossing diagrams is at least

$$\frac{1}{d(n)} \left(d(n, n) + \frac{3}{4}d(n, n-2) + \frac{7}{8}d(n, n-3) \right)$$


Proof.

You can make a similar argument for $d(n, n-3)$.




Question

Does this (crude) bound explain the unknot fraction?



Future Direction: How to enumerate diagrams (like a BOSS combinatoricist)




Two-leg diagrams counted by generating function (Bouttier, et al. 2003):


$$G_0 = \frac{24g - 1 + \sqrt{1 - 12g}}{9g(1 + \sqrt{1 - 12g})} = 1 + 2g + 9g^2 + 54g^3 + 378g^4 + \dots$$

We can enumerate two-leg diagrams using blossom trees (Schaeffer).


Future Direction: Uniform sampling of large diagrams



Future Direction: Uniform sampling of large diagrams




Future Direction: Uniform sampling of large diagrams



Future direction: Link diagrams



n	# link shadows	# knot shadows
0	1	1
1	1	1
2	3	2
3	7	6
4	30	19
5	124	76
6	733	376
7	4586	2194
8	33373	14614
9	259434	106421
10	2152298	823832

Future direction: Knot distances



Thank you!

Coming soon: Cantarella, Chapman, Mastin. *Knot probabilities in random diagrams.*



This research was supported in part by NSF grant DMS-1344994 (RTG in Algebra, Algebraic Geometry, and Number Theory, at the University of Georgia).