
Random Knot Diagrams

Harrison Chapman (UGA) joint w/ Jason Cantarella (UGA), Matt Mastin (MailChimp, Inc.) Crucial Assist: Eric Rawdon (St. Thomas)

AMS Fall Western Section Meeting, 10/24/15


Question

What fraction of 8-crossing diagrams are trefoils?




Question

What fraction of 8-crossing diagrams are trefoils? \$12.48%




Question

What is the average minimal crossing # of an 8-crossing diagram?




Question

What is the average minimal crossing # of an 8-crossing diagram? 0.52




Definition

The **untwisting** operator deletes all 1-crossing connect summands of a diagram. (Equivalently, performs all "available" Reidemeister I moves.)




Question

What is the average crossing # of a untwisted 8-crossing diagram?



Question

What is the average crossing # of a untwisted 8-crossing diagram? 2.20



Question

How many 8-crossing diagrams can be untwisted to the unknot?




Question


How many 8-crossing diagrams can be untwisted to the unknot? 42.05%





Ansatz



Combinatorial approaches








Random diagrams

Definition

In the **random diagram model** of random knotting, a *n*-crossing diagram is drawn uniformly from the finite set of *n*-crossing knot diagrams (cf. Dunfield).








Random diagrams

Definition

A **knot diagram** is a equivalence class of generic immersions of the oriented S^1 into the sphere S^2 together with over-under strand information at each double point up to diffeomorphism of S^2 .







How to enumerate knot diagrams (like a topologist)

Definition

A **knot shadow** is a equivalence class of generic immersions of the unoriented S^1 into the sphere S^2 up to diffeomorphism of S^2 .

Plan to Enumerate Diagrams

- **I** Enumerate shadows (and discard isomorphic shadows)
- 2 Assign crossing and orientation information (and discard crossing patterns related by an automorphism of the shadow)

How to enumerate knot diagrams (like a topologist)

Definition

A **knot shadow** is a equivalence class of generic immersions of the unoriented S^1 into the sphere S^2 up to diffeomorphism of S^2 .

Plan to Enumerate Diagrams

- 1 Enumerate shadows (and discard isomorphic shadows)
- 2 Assign crossing and orientation information (and discard crossing patterns related by an automorphism of the shadow)

Observation

Symmetry stinks.

Proposition


Knot shadows \leftrightarrow 1-component 4-valent embedded planar multigraphs up to embedded isomorphism



Proposition

Knot shadows \leftrightarrow 1-component 4-valent embedded planar multigraphs up to embedded isomorphism


1 Add loops and edges to planar simple graphs (slow)



Proposition

Knot shadows \leftrightarrow 1-component 4-valent embedded planar multigraphs up to embedded isomorphism


- Add loops and edges to planar simple graphs (slow)
- 2 Generate multiquadrangulations of sphere by careful pattern of connect sums, take dual graphs (fast)



Proposition

Knot shadows \leftrightarrow 1-component 4-valent embedded planar multigraphs up to embedded isomorphism

- Add loops and edges to planar simple graphs (slow)
- 2 Generate multiquadrangulations of sphere by careful pattern of connect sums, take dual graphs (fast)



Actually generate all link shadows, then restrict to knot shadows

Verifying against existing shadow counts

		_				_	
	oriented	n = 0	1	2	3	4	5
	S^2 , S^1	1	1	3	9	37	182
	S^2	1	1	2	6	21	99
	S^1	1	1	2	6	21	97
		1	1	2	6	19	76
Cu	rves on S^2 . Th	e number (of type	s			
١.	/ L Arnol	'd To	nal	aric.	al In	varia	ntc

V.I. Arnol'd. Topological Invariants of Plane Curves

008989	Number of immersions of unoriented circle into unoriented sphere with n double points.
1, 1, 2, 6,	19, 76, 376, 2194 (list; graph; refs; listen; history; text; internal format)
OFFSET	0,3
REFERENCES	V. I. Arnold, Topological Invariants of Plane Curves, American Math.
LINKS	Table of n , $a(n)$ for $n=07$.
CROSSREFS	Sequence in context: <u>A150119 A181776 A138800 * A057240 A079564 A079453</u> Adjacent sequences: <u>A008986 A008987 A008988 * A008990 A008991 A008992</u>
KEYWORD	nonn
AUTHOR	N. J. A. Sloane.
EXTENSIONS	Two more terms from Guy H. Valette (guy.valette(AT)skynet.be), Feb 09 2
STATUS	approved

n	# knot shadows
0	1
1	1
2	2
3	6
4	19
5	76
6	376
7	2194
8	14614
9	106421
10	823832

OEIS A008989

-	OU, $g = 5$	0	0	0	0	0	0	0	0	22 524 176	
	UU, total	1	3	12	86	894	14715	313 364	8 139 398	245 237 925	8 382 002 270
	UU, g = 0	1	2	6	19	76	376	2194	14614	106 421	823 832
	UU, g = 1	0	1	5	45	335	3101	29 415	295 859	3 031 458	
	TITE O										

Coqueraux, Zuber, arxiv:1507.03163

Assign crossings, orientation, identify

- Orient each component. (2 choices)
- **2** Assign over-under information to each vertex. (2^n choices)

n	# knot shadows	2^{n+1} (# shadows)	# knot diagrams
3	6	96	36
4	19	608	276
5	76	4,864	2,936
6	376	48,128	35,872
7	2,194	561,664	484,088
8	14,614	7,482,368	6,967,942
9	106,421	108,975,104	105,555,336
10	823,832	1,687,207,936	1,664,142,836

Assign crossings, orientation, identify

- 1 Orient each component. (2 choices)
- 2 Assign over-under information to each vertex. (2^n choices)


n	# knot shadows	2^{n+1} (# shadows)	# knot diagrams
3	6	96	36
4	19	608	276
5	76	4,864	2,936
6	376	48,128	35,872
7	2,194	561,664	484,088
8	14,614	7,482,368	6,967,942
9	106,421	108,975,104	105,555,336
10	823,832	1,687,207,936	1,664,142,836

Observation


Symmetry becomes rare, quickly!




Size of the automorphism group of a random diagram




Knotting in diagrams







Ratios of knots in *n*-crossing diagrams (log scale)




Ratios of knots in *n*-crossing diagrams (log scale)




Why so many unknots?







Link shadows. Pictures generated by Eric Lybrand (UGA undergrad).



Link shadows. Pictures generated by Eric Lybrand (UGA undergrad).




Link shadows. Pictures generated by Eric Lybrand (UGA undergrad).





Link shadows. Pictures generated by Eric Lybrand (UGA undergrad).


Most diagrams are (very) composite





Most diagrams are (very) composite


















Mean number of connect summands \simeq crossing number



Proposition

If d(n, k) is the number of diagrams with n crossings and k connect summands, and d(n) is the number of all n crossing diagrams, then the unknot fraction among all n crossing diagrams is at least

$$\frac{1}{d(n)}\left(d(n,n)+\frac{3}{4}d(n,n-2)+\frac{7}{8}d(n,n-3)\right)$$

Proof.

Any diagram in d(n, n) is a connect sum of all 1-crossing diagrams, and so can be simplified to the unknot via RI moves.



Proposition

If d(n, k) is the number of diagrams with n crossings and k connect summands, and d(n) is the number of all n crossing diagrams, then the unknot fraction among all n crossing diagrams is at least

$$\frac{1}{d(n)}\left(d(n,n)+\frac{3}{4}d(n,n-2)+\frac{7}{8}d(n,n-3)\right)$$

Proof.

Any diagram in d(n, n-2) is a connect sum of 1-crossing diagrams, and a prime 3-crossing diagram (turns out there's only one—the trefoil diagram). This diagram is knotted iff those three crossings have the same sign, which occurs 1/4 of the time.

Proposition

If d(n, k) is the number of diagrams with n crossings and k connect summands, and d(n) is the number of all n crossing diagrams, then the unknot fraction among all n crossing diagrams is at least

$$\frac{1}{d(n)}\left(d(n,n)+\frac{3}{4}d(n,n-2)+\frac{7}{8}d(n,n-3)\right)$$

Proof.

You can make a similar argument for d(n, n-3).

Proposition

If d(n, k) is the number of diagrams with n crossings and k connect summands, and d(n) is the number of all n crossing diagrams, then the unknot fraction among all n crossing diagrams is at least

$$\frac{1}{d(n)}\left(d(n,n)+\frac{3}{4}d(n,n-2)+\frac{7}{8}d(n,n-3)\right)$$


Proof.

You can make a similar argument for d(n, n-3).


Question

Does this (crude) bound explain the unknot fraction?


Yes.




Future Direction: Uniform sampling of large diagrams



Future Direction: Uniform sampling of large diagrams




Future Direction: Uniform sampling of large diagrams



Future direction: Link diagrams

n	# link shadows	# knot shadows
0	1	1
1	1	1
2	3	2
3	7	6
4	30	19
5	124	76
6	733	376
7	4586	2194
8	33373	14614
9	259434	106421
10	2152298	823832

Future direction: Knot distances



Thank you!

Coming soon: Cantarella, Chapman, Mastin. *Knot probabilities in random diagrams*.











This research was supported in part by NSF grant DMS-1344994 (RTG in Algebra, Algebraic Geometry, and Number Theory, at the University of Georgia).