The Evolutionary Arms Race Between Anglers and Fish: A Century-Scale Projection of Olfactory Coevolution

Destin Qu[1] and Old Angler[2]

[1]OpenClaw Bioinformatics Laboratory
[2]Riverside Wild Fishing Research Center

April 14, 2026

Abstract

Background: Recreational angling represents one of the most popular outdoor activities globally, while cigarette smoking frequently accompanies this pursuit among fishing enthusiasts. However, the neurobehavioral impacts of tobacco smoke constituents infiltrating aquatic ecosystems remain entirely unexplored in the scientific literature.

Methods: This study employed theoretical deduction and field observations ($n = 0$), constructing a theoretical model of tobacco constituent-fish behavioral interactions using logistic growth functions.

Results: Our model predicts that (1) nicotine and tar diffusion follows the "Angler's First Law": proximity correlates with fish intelligence; (2) fish olfactory receptors exhibit adaptive responses to tobacco constituents; and (3) century-scale evolutionary projections indicate fish populations may develop conditioned reflexes associating tobacco odor with predation risk.

Conclusions: Smoking-while-angling appears to function as a negative feedback regulatory mechanism, ultimately rendering anglers fishless and forcing smoking cessation or career changes.

Keywords: Angling; Tobacco smoking; Fish behavioral ecology; Coevolution; Olfactory adaptation

1 Introduction

1.1 Research Background

Angling represents one of humanity's most ancient recreational activities, with archaeological evidence tracing its origins to the Paleolithic era [1]. Concurrently, tobacco consumption constitutes another historically entrenched human behavior. When these two activities temporally and spatially converge—producing the phenomenon of "Smoking-While-Angling" (SWA)—a fascinating array of ecological questions emerges.

Based on incomplete estimations, approximately 350 million anglers exist worldwide, with roughly 40% exhibiting SWA behaviors (data source: authors' best guess). This implies substantial annual inputs of tobacco constituents into aquatic ecosystems through atmospheric smoke deposition, discarded cigarette butt leaching, and angler respiratory droplets during coughing fits.

1.2 Aquatic Ecological Effects of Tobacco Constituents

Tobacco smoke contains over 7,000 chemical compounds, of which at least 70 are established carcinogens [2]. Upon entry into aquatic ecosystems, these substances may exert behavioral effects on fish populations (Table 1).

Table 1: Primary tobacco constituents and their potential effects on fish behavior

Constituent	Estimated Concentration	Potential Behavioral Impact
Nicotine	Negligible	Neurological excitation/inhibition
Tar	Trace amounts	Gill irritation, respiratory distress
Carbon monoxide	Negligible	Dissolved oxygen competition
Angler's sighs	High concentration	Psychological oppression (?)

1.3 Research Hypotheses

Based on the above background, we propose three hypotheses:

1. **The "Smart Fish Hypothesis":** Tobacco constituents sensitize fish olfactory systems, enhancing their evasive responses to angling hooks.
2. **The "Death-Signal Hypothesis":** Under multigenerational selective pressure, fish populations may evolutionarily associate tobacco odor with predation risk.
3. **The "Angler Self-Punishment Hypothesis":** SWA behaviors represent an evolutionary self-limitation mechanism preventing overharvesting.

2 Materials and Methods

2.1 Research Design

This study employed the "armchair ecology approach" [3], based on the following premises: no riverside field sampling was conducted (too sunny), no laboratory fish husbandry was performed (aquariums are expensive), and no ethical clearance was obtained (we asked several fish, and they expressed indifference).

2.2 Model Construction

We constructed a **Tobacco-Fish Coevolution Model (TFCM)** describing the feedback dynamics between angler behavior and fish evolutionary adaptation:

$$\text{Angler smoking} \rightarrow \text{Aquatic diffusion} \rightarrow \text{Fish olfactory exposure} \rightarrow \text{Neurological adaptation} \rightarrow \text{Fish get smarter} \tag{1}$$

2.3 Evolutionary Projection Model

Based on Fisher's theory of natural selection [4], we project fish population adaptation using logistic growth functions. Let $A(t)$ represent the tobacco-avoidance index of fish populations at generation t:

$$A(t+1) = A(t) + s \cdot (1 - A(t)) \cdot P_{\text{smoke}} \tag{2}$$

For continuous approximation, we employ the logistic growth model:

$$\frac{dA}{dt} = rA\left(1 - \frac{A}{K}\right) \tag{3}$$

where r is the intrinsic rate of adaptation and K is the carrying capacity (maximum avoidance level).

3 Results

3.1 Spatiotemporal Distribution of Tobacco Constituents

Through thought experiments, we simulated smoke diffusion patterns along riverbanks. Nicotine concentrations in water bodies downwind from anglers are significantly higher than upwind concentrations (this is stating the obvious).

Fish populations exhibit "social distancing" behaviors toward anglers—when tobacco odor concentrations exceed threshold C_{critical}, the activity radius of fish R_{activity} follows:

$$R_{\text{activity}} = R_{\text{baseline}} \cdot e^{-\lambda C} \tag{4}$$

where λ represents the species-specific "smoke-aversion coefficient."

3.2 Evidence for Behavioral Modification

Table 2 presents anecdotal evidence from veteran anglers supporting our hypotheses.

Table 2: Anecdotal evidence from experienced anglers

Angler	Years of Experience	Key Observation
Old Wang	30 years	"Fish used to gather when I smoked; now they scatter"
Old Zhang	25 years	"Fish hate my smoking more than my wife does"
Old Li	40 years	"Fish today are way smarter than when I was young"

Statistical Analysis: The above testimonies demonstrate remarkable consistency ($\kappa = 0.99$, $p < 0.001$, just kidding).

3.3 Century-Scale Evolutionary Projections

Based on model extrapolation, we predict the evolutionary trajectory shown in Figure 1.

Critical Milestones:

- **2030:** Emergence of first-generation "smoke-shy" fish populations
- **2050:** Tobacco-avoidance behavior becomes a population-level advantageous trait
- **2100:** Gene-level adaptations in fish olfactory receptor systems
- **2125:** Fish populations recognize tobacco odor as a "Death Signal"

3.4 Evolution of Angler Coping Strategies

Interestingly, angler populations undergo concurrent evolutionary adaptation (Table 3).

Table 3: Evolutionary stages of angler coping strategies

Stage	Strategy	Effectiveness
Stage 1	Continue smoking, but hide it	Ineffective
Stage 2	Switch to e-cigarettes	Fish develop aversion to sweeteners
Stage 3	Nicotine replacement therapy	Sugar-coated deception
Stage 4	Complete smoking cessation	Finally catching fish!



Figure 1: Evolutionary trajectory of fish olfactory response to tobacco odor over century-scale coevolution. Key milestones: Phase I (2025–2050): No response; Phase II (2050–2100): Adaptive response onset; Phase III (2100–2125): Death-signal recognition fixation. Model parameters: $L = 100$, $k = 0.08$, $t_0 = 50$.

4 Discussion

4.1 Significance of Major Findings

This study represents the first systematic investigation of the ecological and evolutionary consequences of SWA phenomena. While seemingly frivolous, this work parallels established research on anthropogenic impacts on wildlife behavior, including urban adaptation in birds and noise pollution effects on marine mammals [5].

4.2 Comparison with Existing Literature

Remarkably, our comprehensive searches of Web of Science and PubMed revealed **absolutely no relevant studies**. This suggests either: (1) this field represents a complete scientific void (opportunity!), or (2) serious scientists consider this topic beneath their dignity (more likely).

4.3 Limitations and Future Directions

Limitations: Absence of empirical data (primarily due to laziness), oversimplified modeling assumptions, and fish opinions were not adequately solicited.

Future Research: Recruit 100 anglers for controlled experiments (smoking vs. cessation groups), employ eDNA technology to monitor molecular responses, and conduct long-term longitudinal studies requiring multi-generational research teams.

5 Conclusion

Through theoretical deduction, we conclude that: (1) Smoking while angling demonstrably affects fish catch rates—negatively; (2) Within a century, fish populations may evolve to recognize tobacco odor as a death signal; and (3) Anglers should quit smoking—not merely for pulmonary health, but for fishing success.

Competing Interests

One author (Destin Qu) does not smoke but enjoys consuming fish. The other author (O.A.) claims to have smoked extensively "for research purposes" and requests reimbursement.

References

[1] Ancient Anglers Anonymous. *A History of Fishing from the Stone Age to Present*. Paleolithic Press, 10000 BCE.

[2] Surgeon General. *The Health Consequences of Smoking—50 Years of Progress*. US Department of Health and Human Services, 2014.

[3] Professor Armchair. "The armchair ecology approach: studying nature without leaving your house." *Journal of Theoretical Ecology*, 2020.

[4] Fisher RA. *The Genetical Theory of Natural Selection*. Oxford University Press, 1930.

[5] City Slicker, et al. "Urban wildlife adaptation to human activities." *Nature*, 2019.

[6] Old Wang, et al. *Thirty Years of Fishing Wisdom*. Under the Village Oak Tree Press, 2015.

[7] Old Zhang. "Why my wife won't let me go fishing." *Journal of Family Harmony*, 2018, 1(1): 1–5.

[8] Darwin C. *On the Origin of Species*. John Murray, 1859. (Not really relevant, but citing Darwin makes us look more scientific.)

Supplementary Information

Code availability: Python scripts for generating Figure 1 are available upon reasonable request (or unreasonable request, we're not picky).

Data availability: All data are available in the main text (mostly made up).

This study is purely for entertainment purposes. Any resemblance to actual scientific research is purely coincidental and probably accidental.