

Personality Disorders Packet

The current literature on personality disorders, what we know, what is debated, and where the field is headed

How to use this packet

This packet is written in plain English but stays close to the literature. It is meant for a psychology student who wants a strong conceptual overview of personality disorders (PDs), how the field got here, and where it seems to be going.

A recurring theme in the modern literature is this: the field is moving away from viewing personality disorders as rigid boxes and toward understanding them as patterns of personality dysfunction that vary in severity, traits, and context. That shift is not complete, and not everybody agrees on the details, but it is the clearest direction of travel.

1. Overview: what personality disorders are

Personality disorders are enduring patterns of inner experience and behavior that are relatively inflexible, show up across situations, cause distress or impairment, and typically begin by adolescence or early adulthood. In practice, they involve persistent difficulties in one or more of the following:

- sense of self
- emotional regulation
- relationships and attachment
- impulse control
- social cognition and empathy
- behavior under stress

The classic DSM approach grouped PDs into three clusters:

Cluster A: odd/eccentric

- Paranoid PD
- Schizoid PD
- Schizotypal PD

Cluster B: dramatic/emotional/erratic

- Antisocial PD
- Borderline PD

- Histrionic PD
- Narcissistic PD

Cluster C: anxious/fearful

- Avoidant PD
- Dependent PD
- Obsessive-Compulsive PD

This cluster system is still widely taught because it is simple, but it has major problems:

- people often meet criteria for multiple PDs at once
- people often meet criteria partially but not cleanly
- diagnostic thresholds can be arbitrary
- within one diagnosis, two patients can look very different
- categories overlap heavily with normal-range personality traits and with other disorders

That is one reason dimensional models have gained traction.

Basic prevalence and burden

The exact prevalence depends on how PDs are defined and measured, but PDs are common enough to matter substantially in clinical practice and public health. Reviews consistently show that PD pathology is associated with:

- lower quality of life
- greater psychiatric comorbidity
- higher service use
- functional impairment
- increased risk of self-harm and suicide, especially in borderline pathology
- interpersonal, occupational, and legal difficulties

Recent epidemiologic reviews suggest that personality pathology is not rare, and prevalence estimates vary notably depending on the population, assessment method, and whether categorical or dimensional criteria are used.

2. A brief history: how personality disorders were conceptualized

Ancient roots and pre-DSM thinking

The idea that some people show persistent maladaptive personality styles is old. Ancient temperament theories, later constitutional theories, and early psychiatric descriptions all tried to explain stable patterns of emotion and conduct.

By the late 19th and early 20th centuries, European psychiatry described what were sometimes called “psychopathic personalities,” “character disorders,” or personality deviations. These early formulations were often impressionistic, morally colored, or theoretically inconsistent.

Psychoanalytic era

Mid-20th-century thinking was strongly shaped by psychoanalysis and psychodynamic theory. Clinicians focused on:

- ego strength
- defense mechanisms
- object relations
- identity integration
- developmental arrest or trauma

The term borderline originally referred less to a checklist diagnosis and more to a structural/developmental level between neurosis and psychosis.

DSM-I and DSM-II

DSM-I (1952) and DSM-II (1968) were heavily influenced by psychodynamic language and did not provide the kind of explicit operational criteria that later DSMs would use. Personality pathology was recognized, but diagnostic reliability was weak.

DSM-III revolution

DSM-III (1980) changed everything. Psychiatry moved toward explicit diagnostic criteria and a more medicalized, descriptive approach. PDs became a more formal set of discrete categories.

This helped with reliability, teaching, and research recruitment. But over time, major limitations became obvious:

- high comorbidity among PDs
- poor discriminant validity
- heterogeneity within categories
- arbitrary thresholds
- unstable category membership even when broader personality pathology remains stable

DSM-IV and DSM-5 Section II

DSM-IV retained the categorical system, and DSM-5 Section II basically kept the 10 familiar disorders. This continuity was politically and clinically practical, but many researchers believed the science had outgrown the model.

DSM-5 Alternative Model for Personality Disorders (AMPD)

DSM-5 also introduced the Alternative Model for Personality Disorders in Section III as an emerging model. This was a major milestone.

The AMPD has two main parts:

Criterion A: level of personality functioning This focuses on impairment in:

- identity
- self-direction
- empathy
- intimacy

Criterion B: maladaptive personality traits Five broad trait domains:

- Negative Affectivity
- Detachment
- Antagonism
- Disinhibition
- Psychoticism

The AMPD also includes trait facets beneath these domains and allows hybrid diagnosis for several retained PD types.

ICD-11 personality disorder model

ICD-11 made an even bolder move. It largely replaced the old list of categorical PD diagnoses with:

- 1 a general diagnosis of personality disorder
- 2 severity rating: mild, moderate, severe
- 3 optional trait domain specifiers:

- Negative Affectivity
- Detachment
- Dissociality
- Disinhibition
- Anankastia

- 1 a borderline pattern qualifier

This is a clear dimensional model emphasizing severity and trait description rather than “which box fits best.”

Big picture historical arc

A fair one-line summary of the history is:

The field moved from moral/character descriptions -> psychoanalytic/developmental theories -> checklist categories -> increasingly dimensional, trait-based, severity-oriented models.

3. The current state of the literature: where the field is now

Strongest current consensus

Across major reviews and consensus-style papers, several ideas are now fairly well supported:

- 1 Categorical PD diagnosis has major psychometric problems.
- 2 Trait models of personality pathology are robust and empirically meaningful.
- 3 Severity of personality dysfunction matters a lot clinically.
- 4 Borderline pathology remains highly clinically important whether conceptualized categorically or dimensionally.
- 5 PDs overlap strongly with normal-range personality structure, especially the Five-Factor Model.
- 6 Comorbidity and cross-diagnostic overlap are the rule, not the exception.
- 7 Treatment is possible; prognosis is often better than older pessimistic views suggested, especially for BPD.

What is still being argued about

The biggest current argument is not whether dimensional approaches are useful. They clearly are. The argument is which dimensional approach is best.

Key debates include:

- Should severity and traits both be required, or are traits enough?
- Is DSM-5 AMPD Criterion A actually distinct from Criterion B, or are they too overlapping?
- Should DSM move fully to a dimensional PD model in the main manual?
- How should dimensional PD models connect with broader models like HiTOP?
- How can these systems be made practical for everyday clinicians, not just researchers?

Where the field is headed

Recent reviews, especially on DSM-5 AMPD and ICD-11, suggest the field is headed toward:

- more dimensional diagnosis

- more use of severity ratings
- more trait-language and less pure category-language
- integration with transdiagnostic models of psychopathology
- stronger measurement of personality functioning/self-interpersonal impairment
- lifespan and adolescent applications
- personalized case formulation over simple label assignment

Lee Anna Clark's 2025 review on DSM-5 and ICD-11 is especially useful here: it describes substantial convergence between the systems while also noting important differences, particularly around the role of maladaptive traits and the status of personality functioning measures.

4. Major theories and models of personality disorders

No single theory explains all personality pathology. A mature understanding of PDs requires multiple lenses.

4.1 Psychodynamic and object relations models

Psychodynamic approaches see PDs as involving enduring patterns in:

- internalized relationship templates
- defenses
- affect regulation
- identity integration
- representations of self and other

Object relations theory emphasizes how early relationships shape internal self-other representations. Personality pathology may arise when these representations are fragmented, unstable, harsh, distrustful, or poorly integrated.

This framework has been especially influential in understanding:

- borderline pathology
- narcissistic pathology
- paranoid traits
- dependency and abandonment sensitivity

Strengths:

- rich clinical description
- strong attention to meaning, development, and relationship patterns

- useful for formulation and psychotherapy

Limitations:

- some constructs historically harder to operationalize
- earlier literature sometimes relied more on theory than on strong comparative evidence

That said, modern psychodynamic PD literature is much more empirical than older stereotypes suggest, particularly in work on mentalization, personality organization, and transference-focused therapy.

4.2 Attachment models

Attachment theory is one of the most influential developmental frameworks in PD research.

Core idea: repeated early relational experiences shape expectations about self, others, safety, closeness, rejection, and emotion regulation.

Insecure attachment is not equivalent to a PD, but attachment insecurity and disorganization are strongly relevant to many forms of personality pathology, especially:

- borderline patterns
- dependent patterns
- avoidant patterns
- narcissistic vulnerability
- chronic distrust/paranoia

Attachment models are especially helpful for explaining:

- fear of abandonment
- unstable closeness-distance regulation
- hyperactivation or deactivation of affect
- mentalizing failures under interpersonal stress

4.3 Cognitive-behavioral models

CBT models focus on dysfunctional beliefs, coping patterns, reinforcement histories, and learned interpersonal strategies.

Examples:

- “I am unlovable”
- “Others will exploit me”
- “I must stay in control”
- “If I show weakness, I will be humiliated”

Personality pathology is understood as relatively rigid but modifiable patterns of:

- attention
- interpretation
- emotion regulation
- avoidance
- safety behavior
- interpersonal behavior

This approach is highly clinically useful because it links theory to intervention.

4.4 Schema therapy

Schema therapy extends CBT by focusing on early maladaptive schemas, coping styles, and schema modes.

This model has been especially influential in PD treatment because it maps chronic personality problems onto understandable developmental themes such as:

- abandonment
- mistrust/abuse
- defectiveness/shame
- emotional deprivation
- subjugation
- unrelenting standards

Schema therapy has one of the stronger treatment literatures for PDs, particularly borderline personality disorder, and has also been used for cluster C and forensic populations.

4.5 Interpersonal theory

Interpersonal models conceptualize PDs as rigid, recurring interpersonal patterns organized around themes like agency/dominance and communion/warmth.

This lens is particularly good for understanding why PDs often become obvious in relationships:

- people pull predictable reactions from others
- maladaptive interpersonal cycles become self-confirming
- symptoms may intensify under relational stress

Interpersonal theory is especially useful for narcissistic, dependent, avoidant, paranoid, and borderline presentations.

4.6 Biopsychosocial models

Biopsychosocial perspectives emphasize that personality pathology emerges from interactions among:

- temperament
- genetic liability
- attachment and caregiving
- trauma/adversity
- peer and social environments
- culture
- neurobiological sensitivity and regulation
- learning history

This is probably the fairest high-level model because the literature does not support purely genetic or purely trauma-only explanations.

4.7 Trait and dimensional models

Trait models are now central in the science of PDs. They assume maladaptive personality can be described using dimensions rather than just categories.

A common finding is that many PD features map onto extremes or maladaptive variants of broad personality dimensions, especially those related to:

- neuroticism/negative emotionality
- antagonism vs agreeableness
- detachment vs extraversion
- disinhibition vs conscientiousness/constraint
- unusual beliefs/perceptual dysregulation

The AMPD and ICD-11 models both sit in this broader movement.

4.8 HiTOP relevance

The Hierarchical Taxonomy of Psychopathology (HiTOP) is a broader dimensional model of mental disorder, not just PDs. It organizes psychopathology hierarchically and emphasizes spectra rather than categories.

Why it matters for PDs:

- PD traits fit naturally into broader dimensions of psychopathology
- it helps explain comorbidity
- it reduces the artificial split between personality disorders and other disorders

- it encourages transdiagnostic case formulation

HiTOP is highly relevant conceptually, though not yet a routine clinical diagnostic system.

4.9 AMPD relevance

The AMPD is currently one of the best-developed dimensional PD systems in the DSM world. Its strengths include:

- strong empirical support for trait structure
- compatibility with normal personality research
- clinically richer description than simple categories
- support for individualized formulation

But an important criticism, highlighted in recent reviews, is that Criterion A may not be as distinct from Criterion B as originally hoped. Some authors argue that Criterion A adds limited discriminant or incremental validity, while others think it captures clinically essential severity/self-interpersonal dysfunction.

4.10 ICD-11 trait model relevance

ICD-11 is influential because it operationalizes a simple, clinically oriented dimensional model:

- diagnose personality disorder
- rate severity
- describe trait domains
- optionally mark borderline pattern

The literature increasingly supports its clinical utility, though there are still concerns about implementation, training, and measurement of personality functioning.

5. Theodore Millon: why he matters

If you are studying personality disorders, Theodore Millon is one of the names you are expected to know.

What Millon did

Millon was one of the most important architects of modern PD thinking. His influence shows up in at least five places:

- 1 He treated personality as a coherent domain worthy of serious classification.
- 2 He built integrative theories, rather than staying in one narrow camp.
- 3 He strongly influenced DSM-era PD thinking, directly and indirectly.
- 4 He connected theory, diagnosis, and assessment, including test development.

- 5 He helped institutionalize the field, including through the Journal of Personality Disorders and the International Society for the Study of Personality Disorders.

Millon's theoretical approach

Millon tried to integrate:

- evolutionary ideas
- learning theory
- interpersonal dynamics
- biological dispositions
- cognitive style
- psychodynamic concepts

He viewed personality styles as patterned adaptations that could become maladaptive when they were rigid, self-defeating, or context-inappropriate.

Why Millon was influential

Millon helped push personality disorders toward:

- more differentiated subtypes
- more theory-based classification
- stronger links between personality theory and clinical assessment
- a richer language for describing personality patterns beyond symptom checklists

Strengths of Millon's legacy

- expansive and intellectually ambitious
- clinically vivid
- historically important in DSM development
- helped normalize the idea that PDs deserved their own scientific literature and professional community

Limitations and critiques

- some of his formulations were more theory-heavy than empirically verified
- not all subtypes or distinctions held up equally well under modern psychometric scrutiny
- some of his classification ideas fit awkwardly with newer dimensional trait science

Bottom line on Millon

Millon is best understood as a bridge figure: he helped move the field from looser clinical theorizing toward more systematic conceptualization of personality pathology. Even where the field has moved beyond parts of his model, his influence remains all over contemporary PD theory and classification.

6. Genetics and heritability: are PDs “75% genetic” now?

Short answer: No, that is too strong and too simplistic.

The claim that personality disorders are now being conceptualized as “75% genetic” is not a good summary of the literature.

What the literature actually supports

6.1 Heritability is real, but it varies

Twin, family, and register studies generally support moderate genetic influence on personality pathology. Many estimates for PD traits or diagnoses fall somewhere in the broad range of about 30% to 60%, with variation depending on:

- which PD is studied
- whether the outcome is a diagnosis or a trait dimension
- age and developmental period
- sample type (community vs clinical)
- method of assessment
- model assumptions

Some studies produce higher estimates, but that does not mean the field as a whole now agrees PDs are 75% genetic.

6.2 Borderline personality disorder is the best-studied example

BPD has among the strongest genetic/familial literatures within PD research.

Relevant findings include:

- older twin studies often found moderate heritability for BPD traits
- a longitudinal twin study from adolescence to adulthood found substantial but not absolute genetic contribution to BPD traits
- a large Swedish population register study reported familial aggregation and estimated heritability of diagnosed BPD in a moderate range rather than near-total genetic determination

A key point from the 2019 Swedish register study is that prior heritability estimates varied widely, and that method matters a lot. This directly argues against repeating a single oversimplified number.

Why “75% genetic” is misleading

There are several reasons that rumor is shaky:

A. Heritability is not destiny Heritability is a population statistic, not a statement that 75% of one person's disorder is caused by genes.

B. Heritability changes by context A trait can show different heritability across populations, developmental stages, and environments.

C. Measured molecular genetics still explains only a fraction Twin studies estimate latent genetic influence, but genome-wide association studies usually explain much less of the variance directly. This gap is common across psychiatry and personality research.

D. Gene-environment interplay matters Genes can shape temperament, emotional sensitivity, impulsivity, and stress reactivity, but environmental exposures help determine how these liabilities unfold.

What is reasonably well supported

- There is meaningful genetic liability for personality pathology.
- Genetic risk likely overlaps with liability for other psychopathology.
- PDs are not best understood as purely trauma-based or purely biologically fixed.
- Borderline pathology, antisocial traits, and broader maladaptive personality dimensions all show familial/genetic signal.

What remains uncertain or debated

- the exact heritability of each specific PD under modern dimensional definitions
- how best to map genetic findings onto ICD-11/AMPD dimensions
- whether specific genetic mechanisms differ more by trait domains than by traditional PD categories
- how developmental timing and trauma exposure moderate genetic risk

Best conclusion for a student

If somebody says “personality disorders are 75% genetic now,” the evidence-based response is:

The literature supports moderate genetic influence, but not a clean universal 75% figure. Estimates vary widely, and PDs are best understood through gene-environment interaction rather than genetic determinism.

7. Environmental and developmental contributors

If the “75% genetic” rumor is too strong, the opposite myth is also wrong: PDs are not explained by trauma alone.

The strongest model is developmental and interactive.

7.1 Temperament

Temperament likely provides the raw material. For example:

- high negative emotionality may increase vulnerability to borderline, avoidant, or dependent pathology
- low fear and callous-unemotional tendencies may increase risk for antisocial traits
- high behavioral inhibition may contribute to avoidant traits
- rigid overcontrol may contribute to obsessive-compulsive personality features

7.2 Attachment and caregiving

Caregiving environments shape:

- trust vs mistrust
- emotional validation vs invalidation
- self-coherence
- capacity for closeness
- expectations about abandonment, rejection, and control

Insecure or disorganized attachment does not automatically cause PDs, but it is strongly relevant in many pathways to personality pathology.

7.3 Trauma and adversity

The literature strongly links childhood adversity with later personality pathology, especially borderline features. Commonly implicated experiences include:

- emotional abuse
- sexual abuse
- physical abuse
- neglect
- chronic invalidation
- chaotic or frightening caregiving

Important nuance:

- trauma is a robust risk factor, especially for BPD-related symptoms
- trauma is not specific to PDs; it is transdiagnostic
- many people with trauma histories do not develop PDs
- some people with PDs do not report severe trauma

So trauma matters a lot, but it is not the whole story.

7.4 Family context

Relevant family variables include:

- inconsistency
- hostility
- enmeshment
- coercive dynamics
- poor boundaries
- neglect
- parental psychopathology
- low warmth or high criticism

These do not act in isolation. They interact with temperament and broader social context.

7.5 Social learning and peer experiences

Personality patterns can also be reinforced by:

- bullying and chronic rejection
- deviant peer affiliation
- coercive conflict patterns
- social reward for grandiosity, aggression, or emotional overreaction
- repeated experiences that confirm maladaptive self-beliefs

7.6 Culture and context

What counts as maladaptive personality expression is partly culturally mediated. The field is increasingly aware that diagnosis should consider:

- cultural norms
- context-specific behavior
- minority stress
- migration/displacement
- socioeconomic adversity

Developmental summary

The most defensible developmental formula is:

temperament/genetic liability + attachment/caregiving + trauma/adversity + learning history + culture/context -> personality pathology risk

8. Disorder-by-disorder and cluster summary

This section keeps the older category language because it is still how many classes, clinicians, and licensing exams talk.

Cluster A

Paranoid Personality Disorder Core themes:

- distrust
- suspiciousness
- interpreting others as malevolent
- grudges/hypervigilance

Clinical notes:

- often overlaps with trauma-related threat sensitivity and interpersonal schemas
- can be difficult to treat because the therapeutic relationship itself may be viewed suspiciously

Schizoid Personality Disorder Core themes:

- social detachment
- restricted emotional expression
- preference for solitude
- low apparent need for intimacy

Clinical notes:

- important to distinguish from autism spectrum conditions, depression, social anxiety, or avoidant dynamics
- historically understudied

Schizotypal Personality Disorder Core themes:

- social/interpersonal deficits
- cognitive-perceptual oddities
- eccentric behavior
- suspiciousness or unusual beliefs

Clinical notes:

- sits close to the schizophrenia spectrum in many models

- often discussed as a bridge diagnosis between personality pathology and psychosis-spectrum conditions

Cluster B

Antisocial Personality Disorder Core themes:

- disregard for others' rights
- deceitfulness
- impulsivity
- irresponsibility
- aggression/lack of remorse

Clinical notes:

- developmental history matters a lot, including conduct problems and callousness
- modern work often separates impulsive/disinhibited antisociality from more callous-antagonistic traits
- important in forensic settings

Borderline Personality Disorder Core themes:

- emotional instability
- unstable relationships
- fear of abandonment
- identity disturbance
- impulsivity
- recurrent self-harm/suicidality in some patients

Clinical notes:

- by far the most studied PD
- robust evidence supports psychotherapy
- prognosis is better than older stereotypes suggested, though functional recovery may lag behind symptom remission
- modern literature increasingly emphasizes early intervention, trauma-informed care, and dimensional formulation

Histrionic Personality Disorder Core themes:

- attention seeking

- emotional expressiveness that may seem theatrical or shallow
- need for reassurance/approval
- impressionistic style

Clinical notes:

- historically controversial and probably underresearched
- critics argue the category has been shaped by gendered assumptions in some eras

Narcissistic Personality Disorder Core themes:

- grandiosity and/or vulnerability
- entitlement
- need for admiration
- impaired empathy
- self-esteem regulation difficulties

Clinical notes:

- modern work often distinguishes grandiose and vulnerable narcissism
- interpersonal and psychodynamic models are particularly helpful here
- not all narcissistic pathology looks loud or arrogant; some is fragile, ashamed, resentful, and hypersensitive

Cluster C

Avoidant Personality Disorder Core themes:

- social inhibition
- feelings of inadequacy
- hypersensitivity to criticism or rejection
- desire for connection but fear of humiliation

Clinical notes:

- overlaps strongly with social anxiety disorder, though many patients show broader identity and relational patterns than standard social anxiety alone
- a clinically important and often impairing disorder

Dependent Personality Disorder Core themes:

- excessive need to be taken care of
- difficulty making decisions independently
- fear of separation
- submissiveness

Clinical notes:

- attachment and interpersonal formulations are very relevant
- can be masked in relationships that appear stable from the outside

Obsessive-Compulsive Personality Disorder (OCPD) Core themes:

- perfectionism
- rigidity
- overcontrol
- preoccupation with order, rules, morality, productivity

Clinical notes:

- distinct from OCD, though they can co-occur
- ICD-11's anankastia domain captures much of this territory dimensionally
- can look "high functioning" while still causing major relational and emotional cost

What the disorder-by-disorder literature looks like now

One of the uncomfortable truths in the literature is that BPD has received far more empirical attention than most other PDs. For many non-BPD categories, the evidence base is smaller, the boundaries are fuzzier, and the future probably lies more in trait/severity description than in preserving every old category as-is.

9. Treatment directions and prognosis

9.1 General treatment principles

The modern literature no longer supports blanket therapeutic nihilism about PDs. That older attitude was wrong and harmful.

General principles with good support include:

- a strong therapeutic alliance
- clear structure and boundaries
- consistent formulation

- long enough treatment to change entrenched patterns
- focus on emotion regulation, self-functioning, and relationships
- attention to comorbidity and risk

9.2 Psychotherapy is the main treatment

For personality pathology, especially BPD, psychotherapy remains first-line.

Approaches with substantial evidence include:

- Dialectical Behavior Therapy (DBT)
- Mentalization-Based Treatment (MBT)
- Transference-Focused Psychotherapy (TFP)
- Schema Therapy
- General psychiatric management and other structured approaches

Meta-analyses have generally found that psychotherapies help reduce core borderline symptoms and related outcomes such as self-harm, though effect sizes vary and no single therapy wins every comparison.

9.3 Medications: limited and targeted role

Medication is often used in practice, but the evidence for medication as a core treatment for personality disorder itself is much weaker than the psychotherapy literature.

Recent meta-analytic work in BPD suggests some drugs may help specific symptom targets in selected circumstances, but the evidence is mixed and often limited by small trials or risk of bias. The broad trend in the literature is:

- avoid treating “the whole personality disorder” with a pill
- use medication cautiously and symptom-targetedly
- prioritize psychotherapy, formulation, and comorbidity management

9.4 Prognosis

This is one of the most important corrective messages in the modern literature:

Many people improve substantially over time.

Especially in BPD research, studies have shown that:

- symptom remission is common over time
- suicidality and acute instability often decrease
- interpersonal and vocational functioning may improve more slowly
- functional recovery can lag behind diagnostic remission

So the honest message is neither hopelessness nor magical optimism. It is:

PDs can be persistent and impairing, but many patients improve meaningfully, especially with good treatment and time.

9.5 New directions in treatment research

Current treatment literature is increasingly focused on:

- early intervention
- adolescence and emerging adulthood
- digital augmentation of care
- mechanism-focused treatments
- trauma-informed adaptations
- more personalized treatment matching
- outcome monitoring and feedback-informed care

10. Major controversies and debates

Debate 1: categories vs dimensions

This is the biggest one. The evidence generally favors dimensions, but categories remain easier to teach, bill, and communicate quickly.

Debate 2: is Criterion A real and distinct?

Some recent reviews argue AMPD Criterion A overlaps heavily with traits and may not justify its separate status. Others think severity of self/interpersonal dysfunction is too clinically important to drop.

Debate 3: should borderline remain special?

ICD-11 kept a borderline pattern qualifier. Some see this as a practical win because clinicians know the construct and the treatment literature is strong. Others think it undermines the move away from categories.

Debate 4: are some classic PD categories outdated?

Histrionic, schizoid, dependent, and even paranoid PD have less robust modern literatures than BPD or broader trait models. Some researchers suspect several traditional categories may eventually matter less than dimensions.

Debate 5: overpathologizing personality

There is always a risk that culturally disapproved, eccentric, or simply difficult traits get pathologized. The field has to distinguish enduring dysfunction from mere difference.

Debate 6: trauma model vs temperament/genetic model

The best literature supports interaction, not either-or thinking. Still, people often polarize around one side.

Debate 7: stigma

PD diagnoses, especially BPD and ASPD, can trigger stigma from clinicians themselves. This affects care quality, alliance, and prognosis. Modern literature increasingly treats stigma reduction as part of competent practice.

11. Practical takeaways for a psychology student

- 1 Know both systems. You still need the classic DSM categories, but you also need to understand AMPD and ICD-11.
- 2 Think formulation, not just label. Ask: what are the person's core self/interpersonal problems, traits, defenses, attachment patterns, and developmental history?
- 3 Borderline is important, but not the whole field. Don't let the BPD literature trick you into thinking all PD science is BPD science.
- 4 Learn trait language. Negative affectivity, antagonism, detachment, disinhibition, psychoticism, dissociality, anankastia: this language is becoming more important.
- 5 Separate evidence from folklore. Examples: "BPD patients can't get better," "PDs are just trauma," "PDs are 75% genetic," "narcissism always means grandiosity." All bullshit oversimplifications.
- 6 Watch your own stigma. If you are annoyed by a patient, that may be clinically meaningful, but it is not a diagnosis.
- 7 Treatment works best when it is structured and relationally thoughtful. Alliance, boundaries, formulation, and consistency matter.
- 8 Function matters as much as symptoms. Remission of crisis behaviors does not automatically mean recovery in identity, work, or relationships.
- 9 Dimensional models are the future. Even if boards or classes still emphasize categories, learn the dimensional systems now.
- 10 Be careful with certainty. The PD field is advancing, but it still contains conceptual and measurement disagreements.

12. Myth vs fact

Myth: Personality disorders are basically untreatable.

Fact: False. Many patients improve substantially, especially with structured psychotherapy.

Myth: Personality disorders are just "bad personalities."

Fact: False and stigmatizing. PDs involve persistent dysfunction in self, emotion regulation, and relationships, not moral failure.

Myth: Borderline personality disorder is the same thing as trauma.

Fact: Trauma is a major risk factor, but BPD is not reducible to trauma alone.

Myth: PDs are now understood as 75% genetic.

Fact: Overstated. The literature supports moderate heritability and strong gene-environment interplay, not one universal number.

Myth: DSM categories are scientifically settled.

Fact: Not even close. Their limitations are one reason AMPD and ICD-11 dimensional models were developed.

Myth: Narcissism always looks arrogant and grandiose.

Fact: Not always. Vulnerable narcissistic presentations can look ashamed, hypersensitive, withdrawn, and resentful.

Myth: Once someone no longer meets criteria, they are fully recovered.

Fact: Symptom remission and functional recovery are related but not identical.

13. Quick glossary

AMPD — Alternative Model for Personality Disorders in DSM-5 Section III.

Anankastia — ICD-11 trait domain involving rigidity, perfectionism, orderliness, and overcontrol.

Criterion A — In the AMPD, level of personality functioning: identity, self-direction, empathy, intimacy.

Criterion B — In the AMPD, maladaptive personality traits.

Dimensional model — A model that describes pathology on continua rather than all-or-none categories.

Discriminant validity — Whether a construct is meaningfully distinct from other constructs.

HiTOP — Hierarchical Taxonomy of Psychopathology, a dimensional model of mental disorders.

ICD-11 — 11th edition of the International Classification of Diseases.

Maladaptive trait — A personality trait that is extreme, inflexible, and associated with dysfunction or impairment.

Mentalization — The capacity to understand behavior in terms of mental states like feelings, wishes, and beliefs.

Object relations — Internalized representations of self and others shaped by relationships.

Personality functioning — Broad self and interpersonal functioning, central in dimensional PD models.

Severity — The degree of impairment and pervasiveness of personality pathology.

Trait domain — A broad category of personality characteristics, such as detachment or antagonism.

14. Bottom-line summary

If you had to compress the modern PD literature into a few sentences, here is the version worth remembering:

- Personality disorders are real, impairing, and clinically important.
- The old categorical system helped organize the field, but it has serious limitations.
- The strongest scientific movement now is toward dimensional models centered on severity and maladaptive traits.
- BPD remains the best-studied diagnosis and the clearest example that treatment can help and prognosis is not uniformly grim.
- PDs arise from interacting genetic, temperamental, developmental, relational, and social factors.
- Theodore Millon was a major bridge figure in the modern conceptualization of personality pathology.
- The future of the field is likely more dimensional, more developmental, more transdiagnostic, and more personalized.

References

Note: References are listed conservatively. When exact pagination was not verified here, the citation is given by authors, title, source, and year.

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders (5th ed.). Washington, DC: American Psychiatric Association; 2013.
- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders (5th ed., text revision; DSM-5-TR). Washington, DC: American Psychiatric Association; 2022.
- World Health Organization. International Classification of Diseases 11th Revision (ICD-11). Geneva: WHO; 2019/2022 implementation materials.
- Clark LA. Wherefrom and Whither PD? Recent Developments and Future Possibilities in DSM-5 and ICD-11 Personality Disorder Diagnosis. *Current Psychiatry Reports*. 2025.
- García LF, and colleagues. The Alternative Model of Personality Disorders: Assessment, Convergent and Discriminant Validity, and a Look to the Future. *Annual Review of Clinical Psychology*. 2024.
- Tyrer P. Personality disorders. *British Journal of Psychiatry*. 2023.
- Pincus AL, Cain NM, Krueger RF, Wright AGC, Scott LN, Hopwood CJ. Theodore Millon's Contributions to Conceptualizing Personality Disorders. *Journal of Personality Assessment*. 2015.
- Fariba KA, Gupta V, Kassir M. Personality Disorder. In: StatPearls [Internet]. Treasure Island, FL: StatPearls Publishing; updated 2024.
- Sharp C, Wall K. Borderline personality disorder in adolescents and young adults: current knowledge and future directions. (Useful developmental review literature; consult current edition/source if assigning formally.)
- Gunderson JG, Herpertz SC, Skodol AE, Torgersen S, Zanarini MC. Borderline personality disorder. *Nature Reviews Disease Primers*. 2018.

- Stoffers-Winterling JM, Storebø OJ, Lieb K. Psychotherapy for borderline personality disorder: contemporary systematic review/Cochrane-style evidence base. See current review updates.
- Cristea IA, Gentili C, Cotet CD, Palomba D, Barbu C, Cuijpers P. Efficacy of Psychotherapies for Borderline Personality Disorder: A Systematic Review and Meta-analysis. *JAMA Psychiatry*. 2017.
- Storebø OJ, Stoffers-Winterling JM, Völlm BA, et al. Psychological therapies for people with borderline personality disorder. *Cochrane Database of Systematic Reviews*. 2020.
- Bateman AW, Fonagy P. Mentalization-based treatment of borderline personality disorder and related work. Multiple editions/reviews.
- Linehan MM. *Cognitive-Behavioral Treatment of Borderline Personality Disorder*. New York: Guilford Press; 1993.
- Young JE, Klosko JS, Weishaar ME. *Schema Therapy: A Practitioner's Guide*. New York: Guilford Press; 2003.
- Kernberg OF, Yeomans FE, Clarkin JF. Transference-focused psychotherapy and personality organization literature. Multiple key publications.
- Gerolymos C, and colleagues. Efficacy and safety of pharmacological treatments in borderline personality disorder: A systematic review and network meta-analysis. *Molecular Psychiatry*. 2026.
- Bornoalova MA, Hicks BM, Iacono WG, McGue M. Stability, Change, and Heritability of Borderline Personality Disorder Traits from Adolescence to Adulthood: A Longitudinal Twin Study. *Development and Psychopathology*. 2009.
- Torgersen S, Lygren S, Øien PA, et al. A twin study of personality disorders. *Comprehensive Psychiatry*. 2000.
- Kendler KS, Czajkowski N, Tambs K, et al. Dimensional representations of DSM-IV cluster A personality disorders in a population-based sample of Norwegian twins: a multivariate study. *Psychological Medicine*. 2006.
- Kendler KS, Czajkowski N, Tambs K, et al. Dimensional representations of DSM-IV cluster B personality disorders in a population-based sample of Norwegian twins: a multivariate study. *Psychological Medicine*. 2008.
- Kendler KS, Aggen SH, Czajkowski N, et al. Dimensional representations of DSM-IV cluster C personality disorders in a population-based sample of Norwegian twins: a multivariate study. *Psychological Medicine*. 2007/2008 literature.
- Reichborn-Kjennerud T, Czajkowski N, Ystrom E, et al. Familial risk and heritability of diagnosed borderline personality disorder: a register study of the Swedish population. *Molecular Psychiatry*. 2019.
- Amad A, Ramoz N, Thomas P, Jardri R, Gorwood P. Genetics of borderline personality disorder: systematic review and proposal of an integrative model. *Neuroscience and Biobehavioral Reviews*. 2014.
- Wright AGC, Hopwood CJ, Krueger RF. Integrating personality assessment and psychopathology structure; major trait/dimensional publications relevant to PDs. Multiple sources.

- Krueger RF, Hobbs KA, Conway CC, et al. Validity, utility, and supporting evidence for the DSM-5 alternative model of personality disorder. *World Psychiatry*. 2021/2024-related literature should also be consulted for latest updates.
- Bach B, First MB. Application of the ICD-11 classification of personality disorders. Key reviews and implementation papers.
- Bach B, Sellbom M, Kongerslev M, Simonsen E, Krueger RF, Mulder R. ICD-11 personality disorder trait domain specifiers and their relation to DSM-5 AMPD traits. Key comparative literature.
- Kim YR, Blashfield R, Tyrer P, and colleagues. Practical implications of ICD-11 personality disorder classifications. *BMC Psychiatry*. 2024.
- Hopwood CJ, Wright AGC, Krueger RF, Morey LC, and colleagues. Personality disorders: current conceptualizations and challenges. *Annual Review of Clinical Psychology*. 2025.
- Reviews and consensus pieces on HiTOP and personality pathology by Kotov, Krueger, Watson, Wright, Hopwood, and colleagues.
- Reviews on trauma as a transdiagnostic risk factor for mental disorders, including relevance to personality pathology, in recent umbrella review literature.
- *Frontiers in Psychiatry* article: Twenty years of research on borderline personality disorder: a scientometric analysis of hotspots, bursts, and research trends. 2024.
- *The Lancet Psychiatry*. The global epidemiology of personality disorder: a systematic review and meta-analysis. 2025.

Final note for study use

If you are turning this into a paper, the smartest next move is to use this packet as a map, then pull 10-20 of the references most relevant to your exact assignment and verify the final citation details in your library database before submission.