

Physics 2: Advanced PHYC10002

Edward Wang

Semester 2, 2025

Contents

1 Electricity

1

1 Electricity

Definition 1. *Electrostatics* concerns forces between charges at rest.

Theorem 1 (Coulomb's law). *The electrostatic force experienced by a charge q_1 in the vicinity of another charge q_2 is equal to*

$$\mathbf{F}_1 = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{|\mathbf{r}_{12}|^2} \hat{\mathbf{r}}_{12},$$

where $\epsilon_0 \approx 8.85 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$.

Definition 2 (Coulomb's constant). Coulomb's law is sometimes written as

$$\mathbf{F}_1 = k_e \frac{q_1 q_2}{|\mathbf{r}_{12}|^2} \hat{\mathbf{r}}_{12},$$

where $k_e \approx 8.99 \times 10^9 \text{ N m}^2/\text{C}^2$.

Remark 1. The electromagnetic force at a nuclear scale is far stronger than the gravitational force. Consider an electron and a proton about 10^{-10} m apart. Given that $8.99 \times 10^9 \text{ N m}^2/\text{C}^2$, $G \approx 6.67 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$, $m_e \approx 9.1 \times 10^{-31} \text{ kg}$, $m_p \approx 1.6 \times 10^{-27} \text{ kg}$, and $e \approx 1.6 \times 10^{-19} \text{ C}$, we would have

$$|\mathbf{F}_E| = k_e \frac{q_1 q_2}{r^2} \approx 2.3 \times 10^8 \gg 9.7 \times 10^{-48} \approx G \frac{m_1 m_2}{r^2} = |\mathbf{F}_g|.$$

It should also be noted that the strong nuclear force, the force of the gluons binding the quarks together within nucleons, is far stronger than the electromagnetic force.