

Space Studies of the Earth-Moon System, Planets, and Small Bodies of the Solar System (B)  
Lunar Science and Exploration (B3.1)  
Consider as poster only.

## **POLARIMETRIC EXPERIMENTS OF LUNAR SOIL SAMPLES BY PARTICLE SIZE**

Serin Kim, fullmoon929@naver.com

[Korea Astronomy and Space Science Institute], [University of Science and Technology], Yongin-si, Korea

Shuai Li

University of Hawaii at Manoa, Honolulu, United States, shuaili@hawaii.edu

Minsup Jeong

Korea Astronomy and Space Science Institute, Daejeon, Korea, msjeong@kasi.re.kr

Kilho Baek

Kyung Hee University, Yongin, Korea, kilho.baek@gmail.com

Sungsoo Kim

Kyung Hee University, Yongin, Korea, sungsoo.kim@khu.ac.kr

Eunjin Cho

Chungnam National University, Daejeon, Korea, jonjin4368@gmail.com

Young-Jun Choi

Korea Astronomy and Space Science Institute, Daejeon, Korea, yjchoi@kasi.re.kr

Polarization properties of the moon provide valuable insights into lunar soil information such as grain size and composition. Polarization phase curves provide these properties but there is a lack of related research and laboratory measurements using lunar samples. For this reason, we have performed polarimetric experiments at four phase angles ( $15^\circ$ ,  $20^\circ$ ,  $25^\circ$ , and  $100^\circ$ ) using Apollo soil samples in multi-band (B, V, and R). A total of five Apollo samples (14163, 14260, 61141, 61221, and 65701) were used. The samples have been divided into different size groups ( $<25$ , 25–45, 45–90, and 90–150  $\mu\text{m}$ ) including the bulk group ( $<150$   $\mu\text{m}$ ). Our investigation focuses on the effect of grain size on the degree of polarization (DoP) and its wavelength dependence. We show that DoP increases as the grain size increases up to the small size group (25–45  $\mu\text{m}$ ), but the larger size groups show various trends. This is different from the results of experiments using terrestrial samples in which particle size and DoP are proportional. Also, for all samples, DoP decreases with increasing the wavelength. In addition, we analyze the negative branch at phase angles between  $15^\circ$  and  $25^\circ$ . In the negative branch, minimum DoP and inversion angle are measured to study their relationship to grain size.