

CHAPTER – 6

ANATOMY OF FLOWERING PLANTS

EXERCISES

2 Mark Questions

Q1: Where is the product of photosynthesis stored?

Answer: Phloem

Q2: What cells curl the leaves in plants during water stress?

Answer: Bulliform cells present in the upper epidermis of monocot leaves make the leaves curl during water stress.

Q3: What is a cambial ring comprised of?

Answer: Cambium is a layer of actively dividing cells between the xylem and phloem tissues which is responsible for the secondary growth of roots and stem.

Q4: If a tree is debarked, what part of the plant is being removed?

Answer: All the tissues external to the vascular cambium, including the secondary phloem, are removed.

Q.5. What are the three types of meristematic tissues?

Answer: Meristematic tissues are a group of young cells, which consists of actively dividing cells. The three main types of **meristematic tissues** are:

1. Apical Meristem.
2. Lateral Meristem.
3. Intercalary Meristem.

4 Mark Questions

Q1: The transverse section of a plant material shows the following anatomical features – (a) the vascular bundles are conjoint, scattered and surrounded by sclerenchymatous bundle sheaths. (b) phloem parenchyma is absent. What will you identify it as?

Answer: The traverse section is of Monocot stem. It is because the vascular bundles are dispersed in monocot stems. The phloem parenchyma is not found

Q2: Why are the xylem and phloem called complex tissues?

Answer: Xylem and Phloem are called complex tissues because they are made of more than one type of cells which work together as a unit to perform the function. Xylem transports water, while phloem transports food.

Q3: How is the study of plant anatomy useful to us?

Answer:

The study of plant anatomy is useful to us in the following ways:

- To understand structural adaptations in plants to different climatic conditions.
- Helpful in identifying monocots, dicots and gymnosperms.
- Physiological conditions can be studied, which helps in crop improvement.
- The study of plant fibres, such as flax, jute, etc., helps in their commercial exploitation, as it enables the prediction of the strength of wood which can be utilised to its potential.

Q4: What is periderm? How does periderm formation take place in the dicot stems?

Answer:

Phellogen, phellem, and phelloderm are collectively known as periderm. While plants undergo secondary growth, the outer epidermal layer and the cortical layer are ripped due to cambium. In order to replace them, the cortex cells turn

meristematic, which produces the cork cambium or the phellogen, which comprises thin-walled, narrow and rectangular cells.

The phellogen sheds cells on either side. The cells which shed from the exterior give rise to the cork or phellem. The suberin accumulates in its cell wall, making it impermeable to water, while the inner cells emerge to become the secondary cortex or phelloderm, which is parenchymatous.

Q5: Cork cambium forms tissues that form the cork. Do you agree with this statement? Explain.

Answer:

Yes, cork cambium forms tissues that, in turn, form the cork. When the stem increases in girth, another meristematic tissue known as phellogen or cork cambium grows in the cortex region of the stem. This phellogen cuts off cells on both sides. The outer cells differentiate into the phellem or the cork, while the inner cells differentiate into the phelloderm or secondary cortex. The cork is impermeable to water because of suberin, rendering protection to the tissues underneath.

Q6:How do open vascular bundles differ from closed vascular bundles?

Answer:

Open Vascular bundles: In open vascular bundles cambium is present between the xylem and the phloem. It has the capability to form secondary xylem and phloem tissues.

Closed Vascular bundles: These in no presence of cambium in closed vascular bundles. Also, these are not capable of forming secondary xylem and phloem tissues.


7 Mark Questions

Q1: Explain the process of secondary growth in the stems of woody angiosperms with the help of schematic diagrams. What is its significance?

Answer:

The primary xylem and phloem exhibit the presence of a strip of cambium in woody dicot roots known as the interfascicular cambium, which is formed from the cells of the medullary rays connecting the interfascicular cambium. Hence, the continuous cambium ring is formed. The cambium separates from the newly formed cells on either side, while the cells found towards the exterior differentiate into the secondary phloem. The cells detach towards the pith giving rise to the secondary xylem. The secondary xylem is synthesised in excess compared to the secondary phloem.

When there is secondary growth in plants, the girth of the plants increases, along with an increase in the water content and nutrients in order to assist the ever-growing leaves, rendering support to the plants.



Q2: Draw illustrations to bring out the anatomical difference between


(a) Monocot root and Dicot root (b) Monocot stem and Dicot stem.

Answer:

(a) Monocot root and Dicot root


Monocot root

MONOCOT ROOT




Dicot root

DICOT ROOT




(b) Monocot stem and Dicot stem

Monocot stem



Dicot stem




Q3: Cut a transverse section of the young stem of a plant from your school garden and observe it under a microscope. How would you ascertain whether it is a monocot stem or a dicot stem? Give reasons.

Answer: Vascular bundles in dicot stem are arranged in a ring whereas in monocot stem vascular bundles are scattered throughout the ground tissue. On the basis of arrangement of vascular bundles it can be ascertained whether the young stem is dicot or monocot. Besides undifferentiated ground tissue, sclerenchymatous hypodermis, oval or circular vascular bundles with Y shaped xylem are other differentiating features of monocot stem.

Q4: What is the stomatal apparatus? Explain the structure of stomata with a labelled diagram.

Answer: Stomata are structures present in the epidermis of leaves. Stomata regulate the process of transpiration and gaseous exchange. Each stoma is composed of two bean-shaped cells known as guard cells which enclose the stomatal pores.

Guard cells are dumbbell-shaped, where their outer wall is thin, and the inner wall is highly thickened. These structures possess chloroplasts and regulate the closing and opening of the stomata. The epidermal cells near the guard cells, in some cases, become specialised in their structure, shape and size; they are referred to as subsidiary cells. The guard cells, the stomatal aperture and girdling subsidiary cells are collectively referred to as the stomatal apparatus.



Q5: Name the three basic tissue systems in flowering plants. Give the tissue names under each system.

Answer:Following are the three basic tissue systems in flowering plants:

Epidermal tissue system

The epidermal tissue system includes the epidermis and epidermal appendages. The epidermis comprises epidermal cells and guard cells, while the epidermal appendages include root hair, stem hair, stinging hair and glandular hair.

The ground tissue system

The ground tissue system is made up of simple tissues such as parenchyma, collenchyma and sclerenchyma.


Vascular tissue system

The vascular tissue system consists of complex tissues such as Xylem, phloem and vascular cambium.

Q6:Describe the internal structure of a dorsiventral leaf with the help of a labelled diagram.

Answer:Dicots exhibit dorsiventral leaves. When examined, the vertical section of a dorsiventral leaf consists of three different parts, and they are

- **Epidermis** – It is found on the adaxial epidermis (upper surface) and the abaxial epidermis (lower surface). On the outside, the epidermis is covered with a thick cuticle. Compared to the upper surface, the abaxial epidermis comprises more stomata.
- **Mesophyll** – It is a tissue found in between the abaxial and adaxial epidermises. This tissue is differentiated into the palisade parenchyma and the spongy parenchyma. The palisade parenchyma is composed of tall, compactly-arranged cells, while the spongy parenchyma comprises round or oval, loosely-arranged cells possessing intercellular spaces. Mesophyll comprises of chloroplasts that carry out photosynthesis.
- **Vascular system** – The vascular bundles that are found in leaves are closed and conjoint, which are engirdled by thick layers of bundle-sheath cells.



Multiple Choice Questions

1. Root hair develop from the region of

- a) Maturation
- b) Meristematic activity
- c) Root cap
- d) Elongation

Answer: a) Maturation

2. Phloem in gymnosperms lacks

- a) Sieve tubes only
- b) Companion cells only
- c) Albuminous cells and sieve cells
- d) Both sieve tubes and companion cells

Answer: d) Both sieve tubes and companion cells

3. The main water-conducting elements of xylem in gymnosperms are

- a) Tracheids
- b) Fibers
- c) Transfusion tissue
- d) Vessels

Answer: a) Tracheids

4. Which of the following yields fibres?

- a) Sisso
- b) Teak
- c) Oak
- d) Coconut

Answer: d) Coconut

5. In a ring girdled plant

- a) The shoot dies first
- b) The root dies first
- c) Neither shoot nor root dies
- d) The root and shoot die together

Answer: b) The root dies first

6. Casparian strips occur in

- a) Cortex
- b) Endodermis
- c) Epidermis
- d) Pericycle

Answer: b) Endodermis

7. Gymnosperms are also called soft wood spermatophytes because they lack

- a) Xylem vessels
- b) Phloem fibres
- c) Cambium
- d) Thick-walled tracheids

Answer: a) Xylem vessels

8. Palisade parenchyma is absent in leaves of

- a) Sorghum
- b) Soybean
- c) Gram
- d) Mustard

Answer: a) Sorghum

9. Xylem translocates

- a) Water, mineral salts, some organic nitrogen and hormones
- b) Water, mineral salts and some organic nitrogen only
- c) Water and mineral salts only
- d) Water only

Answer: a) Water, mineral salts, some organic nitrogen and hormones

10. The common bottle cork is a product of

- a) Xylem
- b) Phellogen