


```
In [2]: import pandas as pd
import seaborn as sns
import matplotlib.pyplot as plt
```


```
In [3]: df=pd.read_csv("Titanic-Dataset.csv")
```

```
In [5]: sns.countplot(x="Sex", data=df)
plt.title("Genderwise Distribution")
plt.show()
```



Observation: Male passengers are more than the female passengers.

```
In [7]: sns.countplot(x="Sex", hue="Survived", data=df)
plt.title("Genderwise Survival")
plt.show()
```




Observation 1: More female passengers survived as compared to male passengers.

Observation 2: More female passengers survived and the the death rate was less.

Observation 3: The death rate of male was more than the survival rate of male.

```
In [8]: sns.countplot(x="Sex", hue="Pclass", data=df)
plt.title("Passenger classwise distribution")
plt.show()
```




Observation 1: Class 3 passengers were in the maximum amount.

Observation 2: Class 2 passengers were the least in number.

Observation 3: The ratio of male and female passengers in all three classes were similar.


```
In [11]: sns.countplot(x="Sex", hue="Age", data=df)
plt.title("Passenger agewise distribution")
plt.show()
```



Observation 1: Passengers in the age range 20-50 were in the most amount.

Observation 2: Passengers in the age range 60-75 were in the least amount.

```
In [15]: sns.histplot(df['Fare'], bins=30, kde=True)
plt.title('Fare Distribution')
plt.xlabel('Fare')
plt.ylabel('Count')
plt.show()
```



Observation 1: Histogram shows a positive skew.

Observation 2: High fares are the outliers.

Observation 3: Majority of fares are concentrated in the lower range.

Observation 4: Higher fares likely correspond to Class 1 passengers.