PNEUMONIA



< Back To Pneumonia

PNEUMONIA


Causes and Risk Factors





Most of the time your body filters germs out of the air that you breathe. Sometimes germs, such as bacteria, viruses, or fungi, get into your lungs and cause infections.

When these germs get into your lungs, your immune system, which is your body's natural defense against germs, goes into action. Immune cells attack the germs and may cause <u>inflammation</u> of your air sacs, or alveoli. Inflammation can cause your air sacs to fill up with fluid and pus and cause pneumonia <u>symptoms</u>. Learn about <u>how the lungs work</u>.



01:36

This video shows how the immune system works to get rid of germs and how pneumonia develops. <u>Medical Animation Copyright © 2022 Nucleus Medical Media Inc. All rights reserved</u> ☑ .

Causes of pneumonia

Bacteria

Bacteria are a common cause of pneumonia in adults. Many types of bacteria can cause pneumonia, but *Streptococcus pneumoniae* (also called pneumococcus bacteria) is the most common cause in the United States.

Viruses

Viruses that infect your lungs and airways can cause pneumonia. The flu (influenza virus) and the common cold (rhinovirus) are the most common causes of viral pneumonia in adults. Respiratory syncytial virus (RSV) is the most common cause of viral pneumonia in young children.

Many other viruses can cause pneumonia, including SARS-CoV-2, the virus that causes COVID-19. Our <u>video about how SARS-CoV-2 affects the lungs</u> ☑ talks about this.

Fungi

Fungi such as *Pneumocystis jirovecii* may cause pneumonia, especially for people who have weakened immune systems. Some fungi found in the soil in the southwestern United States and in the Ohio and Mississippi River valleys can also cause pneumonia.

Risk factors

Your risk for pneumonia may be higher because of your age, environment, lifestyle habits, and other medical conditions.

Age

Pneumonia can affect people of all ages. However, two age groups are at higher risk of developing pneumonia and having more serious pneumonia.

- **Babies and children, 2 years old or younger,** are at higher risk because their immune systems are still developing. This risk is higher for premature babies.
- Older adults, age 65 or older, are also at higher risk because their immune systems generally weaken as people age. Older adults are also more likely to have other chronic (long-term) health conditions that raise the risk of pneumonia.

Babies, children, and older adults who do not get the recommended vaccines to prevent pneumonia have an even higher risk.

Environment or occupation

Most people get pneumonia when they catch an infection from someone else in their community. Your chance of getting pneumonia is higher if you live or spend a lot of time in a crowded place such as military barracks, prison, homeless shelters, or nursing homes.

Your risk is also higher if you regularly breathe in air pollution or toxic fumes.

Some germs that cause pneumonia can infect birds and other animals. You are most likely to encounter these germs if you work in a chicken or turkey processing center, pet shop, or veterinary clinic.

Lifestyle habits

- **Smoking cigarettes** can make you less able to clear mucus from your airways.
- Using drugs or alcohol can weaken your immune system. You are also more likely to accidentally breathe in saliva or vomit into your windpipe if you are sedated or unconscious from an overdose.

Other medical conditions

You may have an increased risk of pneumonia if you have any of the following medical conditions.

• **Brain disorders,** such as a <u>stroke</u>, a head injury, dementia, or Parkinson's disease can affect your ability to cough or swallow. This can lead to food, drink, vomit, or saliva going down your windpipe instead of your esophagus and getting into your lungs.

- Conditions that weaken your immune system may also increase your risk.
 These include pregnancy, HIV/AIDS, or an organ or bone marrow transplant.
 Chemotherapy, which is used to treat cancer, and long-term use of steroid medicines can also weaken your immune system.
- **Critical diseases that require hospitalization,** including receiving treatment in a hospital intensive care unit, can raise your risk of hospital-acquired pneumonia. Your risk is higher if you cannot move around much or are sedated or unconscious. Using a <u>ventilator</u> raises the risk of a type called ventilator-associated pneumonia.
- **Lung diseases**, such as <u>asthma</u>, <u>bronchiectasis</u>, <u>cystic fibrosis</u>, or <u>COPD</u>, also increase your pneumonia risk.
- Other serious conditions, such as malnutrition, <u>diabetes</u>, <u>heart failure</u>, <u>sickle</u> <u>cell disease</u>, or liver or kidney disease, are additional risk factors.

Last updated on March 24, 2022

Back to top



Web Policies & Notices

Accessibility >
No FEAR Act Data >
Privacy Policy >
Freedom of Information Act (FOIA) >

rreedom of information Act (FOIA)

HHS Vulnerability Disclosure ぱ

Related Government Websites

Stay Connected

型 Live Chat 🛮 🖒

Live Chat with us, Monday through Friday, 8:30 a.m. to 5:00 p.m. EST.



Receive automatic alerts about NHLBI related news and highlights from across the Institute.



Contact Us

Site Index

<u>Jobs</u>





