

**MP456**

**Remote Sensing Fundamentals and Applications**

**ANALYSIS OF DISSOLVED OXYGEN**



**Devansh Shukla**  
**I18PH021**  
Integrated Masters Program - Physics

**Department of Physics**

Sardar Vallabhbhai National Institute of Technology

## Acknowledgement

I take an immense pleasure in acknowledging the kind help and guidance provided by Dr. Kamlesh N. Pathak, Professor, Department of Physics, Sardar Vallabhbhai National Institute of Technology, Surat as my instructor for this Project.

I would also like to acknowledge NASA Goddard Space Flight Center, Ocean Ecology Laboratory, Ocean Biology Processing Group; for their ocean color web dataset.

*DEVANSH SHUKLA*

*I18PH021*

# Contents

I	Introduction . . . . .	3
II	Data & Methodology . . . . .	3
II.I	In-situ data: USGS WaterQualityWatch . . . . .	4
II.II	Remote data: Aqua-MODIS . . . . .	5
II.III	Methodology . . . . .	9
III	Analysis . . . . .	9
III.I	Field data . . . . .	9
III.II	AQUA-MODIS data . . . . .	11
III.III	Verification . . . . .	12
IV	Conclusion . . . . .	13
V	References . . . . .	13

## I Introduction

### Motivation

Remote sensing has been an essential means of monitoring dynamic changes such as rainfall, wind, temperature, et al. for weather monitoring and disaster forecast.

Although water quality remains an essential element for the survival of flora and fauna, it is still monitored through in-situ monitoring stations. Even though these stations provide accurate data, their temporal and spatial resolution is inadequate. Thus, there is an opening for satellite remote sensing to develop a method to monitor water quality.

One such parameter for monitoring water quality is dissolved oxygen, which is defined as the amount of free and non-compound oxygen dissolved in the water body(Chi et al., 2020). Monitoring DO can help us predict changes in that marine ecosystem and implement remedies, considering that even a minor change can cause a significant negative impact.

Literature such as Keeling et al., 2010 [?] suggests that dissolved oxygen in the open ocean has been declining in the past centuries due to natural causes or increased temperature or changes in salinity.

DO decreases exponentially with the increase in salinity(Chi et al., 2020).

In summer and winter, the temperature differential is small between day and night, affecting the convection process and hampering the mixture of upper and lower layers of the ocean; due to this and continuous consumption of DO in the lower layers can lead to hypoxic situations. The temperature differential is high in spring and autumn between day and night, allowing the layers to mix and keep the DO concentration consistent.

The traditional method of in-situ measurements is very time and labor-intensive, so this project aims to develop and verify a correlation between dissolved oxygen, temperature, salinity, and chlorophyll content.

## II Data & Methodology

This project aims to develop and verify a correlation between the water parameters such as chlorophyll concentration, temperature, salinity, pH, and specific concentration with Dissolved Oxygen(DO). Therefore, we require both in-situ and remote observation to make quantitative predictions.

Here we have utilized the USGS real-time water parameters for in-situ observations and AQUA-MODIS from NASA ocean color(NASA OBPG) as the remote dataset, discussed in section II.I and II.II respectively.

For analyzing and predicting the dissolved oxygen concentration, a multiple regression model is developed using the data from 2018-01-01 to 2021-12-31, discussed in section II.III




Figure 1. Requirement of DO for marine life

## II.I In-situ data: USGS WaterQualityWatch

### Introduction

USGS provides an almost real-time environment to gather multiple parameters from water in/nearby the United States, i.e., water temperature, pH, salinity, and specific conduction.

It provides the data in tab-separated *.rdb* files with more information about the format available at [waterdata.usgs.gov](http://waterdata.usgs.gov).




Figure 2. USGS WaterQualityWatch stations

### Data processing

The location chosen for analysis is *ORIENT HARBOR AT ORIENT NY*, located at  $41.13663889^{\circ}N$  and  $-72.30675^{\circ}E$ , accurate to  $\pm 1s$

The rationale behind choosing this location was the easy availability of data for 2018-01-01 to 2021-12-31 and its closeness to the Atlantic ocean.

Its closeness to the Atlantic ocean made it easy to obtain the subset of data from the ocean color dataset of Sea Surface Temperature(*SST*) and Chlorophyll(*Chlor-a*)




Figure 3. Satellite blown-up view




Figure 4. Satellite close-up view

#### *NOTE:*

For better results and analysis choosing multiple locations with adequate separation would be ideal, but the turbidity and water contamination may bias our analysis, so we have restricted to a single location as a proof-of-concept.

### II.II Remote data: Aqua-MODIS




Figure 5. MODIS (credits: wikipedia.org)

#### *Introduction*

MODIS (or Moderate Resolution Imaging Spectroradiometers) is an instrument onboard the Terra(EOS AM) and Aqua(EOS PM) satellites. Terra's orbit around the Earth is timed so that it passes from north to south across the equator in the morning, while Aqua passes south to north over the equator in the afternoon. MODIS is designed to provide measurements in large-scale global dynamics, including oceans, land, and cloud cover. The Scan Mirror Assembly uses a continuously rotating double-sided scan mirror to scan  $\pm 55$ -degrees and is driven by a motor encoder built to operate at 100 percent duty cycle throughout the 6-year instrument design life

#### *Specification*

It captures 36 spectral bands ranging in wavelength from  $0.4\mu\text{m}$  to  $14.4\mu\text{m}$  and at varying spatial resolutions. Together the instruments image the entire Earth every 1 to 2 days. It has a sun-synchronous, near-polar orbit of 705km with a swath of 2300km cross-track by 10km along




Figure 6. AQUA-MODIS orbit, credits:NASA GSFC

the track at nadir. It has a scan rate of 20.3rpm cross track. MODIS's temporal resolution is 1-2 days and was launched and designed to work for at least 6-years.

### *Data processing*

#### **Sea Surface Temperature (SST)**

The Sea surface temperature is observed at  $11\mu m$  and  $4\mu m$  for day and night respectively with  $4km$  and  $9km$  spatial resolution. In this analysis, we have utilized the  $11\mu m$  with  $4km$  resolution daily data.




Figure 7. Sea Surface Temperature( $^{\circ}C$ ;  $11\mu m$ ) 2018/01/01

Processed and plotted in python with using 'nctoolkit'; the white portions represent land mass, swaths and cloud cover.




Figure 8. Sea Surface Temperature( $^{\circ}\text{C}$ ;  $11\mu$ ) 2018/01/02  
Processed and plotted using NASA Panoply with interpolations enabled

#### *Monthly averaged*

The daily fluctuations vanishes when we average for a month; the remaining fluctuations are called monthly temperature anomalies.




Figure 9. Montly averaged temperatures for January 2018

## Chlorophyll

The chlorophyll content cannot be directly measured using remote methods, so in this analysis the chlorophyll-A is computed using the NASA OCI algorithm.

This algorithm returns the near-surface concentration of Chlorophyll-A(chlor\_a) in  $mg\ m^{-3}$ , calculated using an empirical relationship derived from in-situ measurements of chlor\_a and remote sensing reflectances ( $R_{rs}$ ) in the blue-to-green region of the visible spectrum.

In this analysis, we have utilized the 4km spatial resolution daily dataset.




Figure 10. Chlorophyll-A 2018/01/04  
Processed and plotted using NASA Panoply with interpolations enabled




Figure 11. Montly averaged chlorophyll-a (OCI) for January 2018

### II.III Methodology

The in-situ data source is *ORIENT HARBOR AT ORIENT NY*, but the remote data is worldwide with 4km resolution. Therefore, we limit our remote data to a polygon around the monitoring station. To reduce complexity, we choose this polygon to be a box with a latitudes range of  $[40, 41.250]^\circ N$  and a longitudes range of  $[-73, -71]^\circ E$ ; this box encompasses an area of  $25000 \text{ km}^2$  (refer to Fig 8).

The remote data subset is spatial averaged to account for fluctuations inside the box.




Figure 12. Subset for remote data

After gathering the data, we use the supervised machine learning algorithm called the **Multiple Linear Regression** to develop a relationship.

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_n x_n \quad (1)$$

We have utilized pH, salinity, field temperature, remote temperature and specific conductance as our input parameters, so our equation will become

$$DO = \beta_0 + \beta_1 pH + \beta_2(\text{salinity}) + \beta_3(\text{field temperature}) + \beta_4(\text{MODIS SST}) + \beta_5(\text{specific conductance}) \quad (2)$$

We split our dataset into two sets, labeled for supervised learning and unlabeled data for testing with 0.8 and 0.2 ratio respectively.

## III Analysis

### III.I Field data

Time-series data for field data from *ORIENT NY* monitoring station; the data is collected at intervals of 6mins, with no data available represented by *NaN*.

As represented in Fig.13a, the temperature shows a seasonal pattern at a frequency of 6-8 months, representing the gap between summer and winter seasons. Other parameters pH, dissolved oxygen, specific conductance and salinity also follows a similar pattern with frequency around 6-months.




Figure 13. Orient Harbour New York, US

### Correlation with dissolved oxygen

By using our field data we can compute correlations matrices to solidify our understanding of correlations between dissolved oxygen and pH, dissolved oxygen and specific conductance et al. Through these correlation matrices we can verify that our parameters have a positive correlation with the concentration of dissolved oxygen.




Figure 14. Correlation matrices with dissolved oxygen

### III.II AQUA-MODIS data

Fig.15 represents the time-series data from the spatially averaged AQUA-MODIS dataset for SST and Chlorophyll-A for our geofenced location of *ORIENT NY*




Figure 15. AQUA-MODIS

### III.III Verification

By plotting a correlation matrix between the field sea temperature and MODIS sea surface temperature we can make sure they are consistent with each other thus reducing the possibility of an error. we verify our correlation of in-situ measured temperature and remotely observed SST.




Figure 16. Verification of in-situ temperature and AQUA-MODIS SST( $11\mu\text{m}$ )




Figure 17. Model test

Fig.17 suggests a good correlation with predictions and test labels of the data. We find that there is 24% error associated with our predictions which could be reduced by using more longer and vivid datasets.

#### IV Conclusion

In this study, DO measurements of Orient NY monitoring station and MODIS data was used to develop and validate the multiple linear regression model.

As discussed in section III, the MLR model predicts with adequate accuracy which could be improved by adding data from multiple sources with multiple water characteristics.

All codes for extraction and processing of data are available at [devanshshukla99/Analysis-of-DO](https://github.com/devanshshukla99/Analysis-of-DO).

#### V References

- Y. H. Kim, S. Son, H.-C. Kim, B. Kim, Y.-G. Park, J. Nam, and J. Ryu, “Application of satellite remote sensing in monitoring dissolved oxygen variabilities: A case study for coastal waters in korea,” Environment International, vol. 134, p. 105301, 2020. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S0160412019327291>

2. NASA Ocean Biology Processing Group. (2018). MODIS-TERRA Level 3 Mapped Chlorophyll Data Version R2018.0 [Data set]. NASA Ocean Biology DAAC. <https://doi.org/10.5067/TEP>
3. R. F. Keeling, A. Körtzinger, and N. Gruber, “Ocean deoxygenation in a warming world,” Annual Review of Marine Science, vol. 2, no. 1, pp. 199-229, 2010, pMID: 21141663. [Online]. Available: <https://doi.org/10.1146/annurev.marine.010908.163855>
4. K. Triana and A. J. Wahyudi, “Dissolved oxygen variability of indonesian seas over decades as detected by satellite remote sensing,” IOP Conference Series: Earth and Environmental Science, vol. 925, no. 1, p. 012003, nov 2021. [Online]. Available: <https://doi.org/10.1088/1755-1315/925/1/012003>
5. M. H. Gholizadeh, A. M. Melesse, and L. Reddi, “A comprehensive review on water quality parameters estimation using remote sensing techniques,” Sensors (Switzerland), vol. 16, 8 2016
6. Chi, L., Song, X., Yuan, Y., Wang, W., Cao, X., Wu, Z., Yu, Z., 2020. Main factors dominating the development, formation and dissipation of hypoxia off the Changjiang Estuary (CE) and its adjacent waters, China. Environ. Pollut. <https://doi.org/10.1016/j.envpol.2020.117551>
7. Schulzweida, Uwe. (2021, October 31). CDO User Guide (Version 2.0.0). Zenodo. <http://doi.org/10.5281/zenodo.5614769>