

Physics 89 - Introduction to Mathematical Physics

deval deliwala

January 27, 2023

Contents

1	Difference between Mathematics and Physics	2
2	Taylor Series	3
2.1	Testing for Convergence	5
3	Complex Numbers	8
3.1	Taylor Series	8
3.2	Complex Numbers	8
3.2.0.1	Rules of Complex Numbers	9
3.3	Applications	10
3.3.0.1	Hydrodynamics	10
3.3.0.2	The Complex Plane	12
3.3.0.3	Euler's Identity	13
3.3.0.4	Trigonometric Functions	13
3.4	Hyperbolic Functions	14
3.4.0.1	Identities	14
3.4.0.2	Applications to Special Relativity	14
3.4.0.3	Taylor Series	18
3.4.0.4	Path Integrals	20
3.4.0.5	Contour Integrals	21

1 Difference between Mathematics and Physics

Example 1 - Electrostatics

Math Question

$$x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots = ?$$

Math Solution

$$x + \frac{x^2}{2} + \frac{x^3}{3} + \dots = -\log(1-x), \quad \text{for } -1 \leq x \leq 1$$

So,

$$-1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} - \frac{1}{5} + \dots = -\log(2)$$

Example 2 - Diffusion

$f(x, y, z, t)$ = density of diffusing material at time t

Let there exist a cube containing moles

$$\frac{\partial f}{\partial t} = D \left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \right)$$

where D is the *diffusion coefficient*, and the diffusion equation describes how f evolves with time

Math Question

Solve

$$\frac{\partial f}{\partial t} = D \left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \right)$$

given initial condition

$f(x, y, z, 0)$ = concentrated lump at the origin

Math Solution

$$f(x, y, z, t) = \frac{N}{(4\pi Dt)^{3/2}} e^{-\frac{x^2+y^2+z^2}{4Dt}}$$

where N is the number of moles released

2 Taylor Series

- Techniques for obtaining series
- Estimate error, converge?




Figure 1: Taylor Series Visualization

$$f(x) \approx f(0) + f'(0)x + \cdots + \frac{1}{n!}f^n(0)x^n$$

$$f(x) \approx f(a) + f'(a)(x-a) + \frac{1}{2}f''(a)(x-a)^2 + \dots = \sum_{k=0}^{\infty} \frac{1}{k!}f^k(a)(x-a)^k$$

Question

How good is this approximation?

Big O notation

$$\sum_{k=0}^n \frac{1}{k!}f^k(0)x^k + O(x^{n+1})$$

Formally,

$$F(x) = o(x^{n+1}) \quad \text{as } x \rightarrow 0$$

$$|F| \leq C|x|^{n+1} \quad \text{for some unexpected constant } c$$

$$\lim_{x \rightarrow 0} \frac{F}{|x|^{n+1}} = 0$$

Example

$$e \approx 1.9 \text{ GeV} \approx 3700 mc^2$$

Special Relativity

$$\begin{aligned} E_k &= m_0 c^2 - mc^2 = \frac{mc^2}{\sqrt{1 - \frac{v^2}{c^2}}} - mc^2 \\ &\approx 0 + \frac{1}{2}mv^2 + \frac{3}{8}m\frac{v^4}{c^2} + \frac{5}{16}m\frac{v^8}{c^4} \\ f(v) &= \frac{1}{2}mv^2 + \frac{3}{8}m\frac{v^4}{c^2} + \dots \end{aligned}$$

$$\frac{1}{\sqrt{1-x}} \rightarrow \frac{mc^2}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$(1+x)^P, \quad \text{then set } p = \frac{1}{2}$$

$$\begin{aligned} f(x) &= (1+x)^n \\ f'(x) &= p(1+x)^{p-1} \\ f^k(x) &= p(p-1)\dots(p-k+1)(1+x)^{p-k} \rightarrow f^k(0) \\ &= p\dots(p-k+1) \end{aligned}$$

$$(1+x)^n \approx 1 + px + \frac{p(p-1)}{2!}x^2 + \dots + \frac{p!}{k!(p-k)!}x^k = \binom{p}{k}x^k$$

$$\sum_{k=0}^n \binom{p}{k} x^k \quad \text{generalized binomial coefficient}$$

$$(1+x)^P = \sum_{k=0}^n \binom{p}{k} x^k + O(x^{n+1})$$

Question

Given $\frac{1}{\sqrt{1+x}}$ Taylor series, how good is this approximation if $x = 0.1$?

Solution

$$\text{Actual Answer} \rightarrow \frac{1}{\sqrt{1.1}} = 0.9534626$$

$$\text{Taylor Polynomials } x, x^2 \rightarrow 1 - \frac{0.1}{2} = 0.95 \quad / \quad 1 - \frac{0.5}{2} + \frac{3(0.5)^2}{8} = 0.95375 \quad \text{good approx}$$

More Taylor Series

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots$$

$$\cosh x = \frac{e^x + e^{-x}}{2} = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \dots$$

$$\sinh x = \frac{e^x - e^{-x}}{2} = x - \frac{x^3}{3!} + \frac{x^5}{5!} + \dots$$

2.1 Testing for Convergence

If $\sum_0^\infty a_n x^n \leq \infty$ converges,

$$\sum_0^\infty a_n (\lambda X)^n \leq \infty \quad |\lambda| \leq 1$$

Taylor Series have interval of convergence of the form

$$[-L, L] \quad (-L, L) \quad [-L, L) \quad (-L, L]$$

Truncated Taylor Series Approximation

$$R_0(x) = f(x) - f(0) = f'(c)x$$




Figure 2: Remainder Visualized

Remainder Theorem

$$R_n(x) = f^{n+1}(c) \frac{x^{n+1}}{(n+1)!} \quad \text{for some } 0 \leq c \leq x$$

$$\begin{aligned} x &= \frac{\pi}{2} \\ R &= \sin \frac{\pi}{2} - \left(x - \frac{x^3}{6} + \frac{x^5}{120} - \frac{x^7}{5040} + \frac{x^9}{362880} + 0 \right) \\ &= f^{10}(c) \frac{x^{10}}{10!} \quad 0 \leq c \leq \frac{\pi}{2} \end{aligned}$$

$$\begin{aligned} |f^{11}(c)| &= |-\cos c| < 1 \\ |R_{10}| &\leq \frac{1}{11!} \left(\frac{\pi}{2} \right)^{11} \approx 3.6 \times 10^{-6} \end{aligned}$$

Technique for Solving Taylor Series by dividing two polynomials

$$f(x) = a_0 + a_1x + \dots$$

$$g(x) = b_0 + b_1x + \dots$$

$$\frac{f(x)}{g(x)} = (c_0 + c_1x + c_2x^2 + \dots)$$

$$a_0 + a_1x + \dots = (b_0 + b_1x + \dots)(c_0 + c_1x + \dots)$$

$$a_0 = b_0c_0$$

3 Complex Numbers

- Definition
- Functions: $\log z, \sqrt{z}, \sin z, \dots$, etc.
- Applications: AC Circuits, Hydrodynamics
- Math Applications: $\int_{-\infty}^{\infty}$

3.1 Taylor Series

$$f(x) = \frac{1}{1+x^2} = \frac{1}{1-(-x^2)} = 1 - x^2 + x^4 - x^6 + \dots = \sum_{n=0}^{\infty} (-1)^n x^{2n}$$

The interval of convergence for the Taylor series of $\frac{1}{1+x^2}$ is from $(-1, 1)$, which is not readily apparent since

$$\text{@ } x \pm 1, f(x) = \frac{1}{2}$$




Figure 3: Taylor series of e^{1/x^2}

3.2 Complex Numbers

Introduced by *Cardano* in the 1500s with the intent of solving cubic equations.

Quadratic Equations

$$0 = x^2 + bx + c \quad x = -b \pm \frac{\sqrt{b^2 - 4ac}}{2a}$$

Cubic Equations

$$0 = x^3 + ax + b \quad \left(\frac{-b}{2} + \sqrt{\frac{b^2}{4} - \frac{a^3}{27}} \right)^{\frac{1}{3}}$$

$$x^3 - x = 0 \rightarrow x = \frac{1}{\sqrt{3}} \left[\sqrt{-1}^{1/3} + (-\sqrt{-1})^{1/3} \right]$$

- consistency
- final answer is **real**
- simplifies computations

3.2.0.1 Rules of Complex Numbers

$$z = a + bi$$

$$i^2 = -1$$

$$(a + bi)(c + di) = (ac - bd) + (ad + bc)i$$

Example

$$(1 + i)^2 = 2i$$

$$i^4 = 1$$

$$\begin{aligned} \frac{1}{a + bi} &= \frac{(a + bi)}{(a - bi)(a + bi)} = \frac{(a - bi)}{a^2 + b^2} \\ &= \left(\frac{a}{a^2 + b^2} \right) - \left(\frac{b}{a^2 + b^2} \right) i \end{aligned}$$

3.3 Applications




Figure 4: 2D diagram of Sphere from above

3.3.0.1 Hydrodynamics

$$\vec{v}(x, y) = v_x \hat{i} + v_y \hat{j}$$

Problem

$$V_x, V_y = ?$$

Model

1. Incompressible

$$(a). \quad 0 = \nabla \cdot \vec{v} = \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y}$$

2. Irrotational

$$(b.) \quad 0 = (\nabla \times \vec{v})_z = \frac{\partial v_x}{\partial x} - \frac{\partial v_x}{\partial y}$$

Solving (a) and (b)

Set of **coupled** partial differential equations (PDEs)

- What are the Boundary Conditions?
 - an additional set of equations at the edges

$$(1.) \quad r = \sqrt{x^2 + y^2} \rightarrow \infty \quad \vec{v} \rightarrow v_0 \hat{i}$$

$$(2.) \quad \vec{v} \cdot \hat{r} = 0$$

Fact: Complex Numbers

Define $z = x + iy$, z is **not** the third coordinate

Define $U = v_x \hat{i} - iv_y$ and $U = f(z) \rightarrow$ Equations (a.) and (b.) are automatically satisfied.

Solution

$$U = v_0 \left(1 - \frac{R^2}{z^2} \right)$$

$$\frac{1}{z} = \frac{1}{x + iy} = \frac{x - iy}{x^2 + y^2}$$

$$\frac{1}{z^2} = \frac{x^2 - y^2 - 2ixy}{(x^2 + y^2)^2}$$

$$v_x = v_0 - \frac{v_0 R^2 (x^2 - y^2)}{(x^2 + y^2)^2}$$




Figure 5: complex plane




Figure 6: quadrant's of complex plane in polar coordinates

3.3.0.2 The Complex Plane

3.3.0.3 Euler's Identity

$$\cos \theta + i \sin \theta = e^{i\theta}$$

$$e^x = 1 + \frac{x}{1} + \frac{x^2}{2} + \frac{x^3}{3!} + \dots \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$\begin{aligned} e^{iy} &= 1 + \frac{iy}{1} - \frac{y^2}{2!} - \frac{iy^3}{3!} + \frac{y^4}{4!} + \dots \\ &= \left(1 - \frac{y^2}{2!} + \frac{y^4}{4!} + \dots\right) + \left(\frac{y}{1} - \frac{y^3}{3!} + \dots\right)i = \cos y + i \sin y \end{aligned}$$

Euler's Identities

$$\begin{aligned} e^{i\pi} &= -1 \\ 1 = e^{2\pi i} &= e^{2\pi ni} \quad n = 0, \pm 1, \pm 2, \dots \end{aligned}$$

$$\log z = ?$$

$$z = re^{i\theta}$$

$$\log z = \log r + i(\theta + 2\pi n)$$

$$\sqrt{z}$$

$$\begin{aligned} \sqrt{re^{iz}} &= \sqrt{r}e^{i\theta/2} \\ &= \sqrt{r}e^{\frac{i(\theta+2\pi)}{2}} \\ &= -\sqrt{r}e^{i\theta/2} \end{aligned}$$

3.3.0.4 Trigonometric Functions

$$\cos z = \frac{e^{iz} + e^{-iz}}{2}$$

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i}$$

$$\cos(iy) = \frac{e^{-y} + e^y}{2} = \cosh y$$

$$\sin(iy) = i \frac{e^y - e^{-y}}{2} = i \sinh y$$

3.4 Hyperbolic Functions

$$\tanh = \frac{\sinh y}{\cosh y}$$

Everything is **Real** from now on.

3.4.0.1 Identities

$$\sinh(\alpha + \beta) = \sinh \alpha \cosh \beta + \cosh \alpha \sinh \beta$$

$$\cosh(\alpha + \beta) = \cosh \alpha \cosh \beta + \sinh \alpha \sinh \beta$$

$$\tanh(\alpha + \beta) = \frac{\tanh \alpha + \tanh \beta}{1 + \tanh \alpha \tanh \beta}$$

3.4.0.2 Applications to Special Relativity Relativistic Addition to Velocities




Figure 7: a train moving with a car moving inside of it, what would an observer calculate for the speed of the interior car?

$$iW = \frac{u + v}{1 + \frac{uv}{c^2}} = c \frac{\tanh \alpha + \tanh \beta}{1 + \tanh \alpha \tanh \beta} = c \tanh(\alpha + \beta)$$




Figure 8: rapidity - using hyperbolic tangent establishes the bounds of velocity as c and $-c$

Jan 26

Functions of Complex Variables

- Cauchy - Riemann Eqns
- Taylor Series
- $\int_{-\infty}^{\infty}$

- Singularities, Poles, Residue

The Complex Conjugate




Figure 9: graphing complex numbers

$$\bar{z} = a - bi \quad z\bar{z} = a^2 + b^2 = |z|^2$$

Functions of $z = x + iy$

$$\begin{aligned} \operatorname{Re} Z &= x \\ \operatorname{Im} Z &= y \\ |z| &= \sqrt{x^2 + y^2} \\ \bar{z} &= x - iy \end{aligned}$$

Analytic Functions

$$\begin{aligned} & \frac{1}{z} \\ & z^2 \\ & e^z \\ & \sin z \\ & \cos z \end{aligned}$$

$$f(z) = u + iv$$

$$\begin{aligned} u &= u(x, y) \\ v &= v(x, y) \end{aligned}$$

Analytic Functions are the functions where you can write

$$f(z + \Delta z) - f(z) = f'(z)\Delta z + O(\Delta z^2) \quad \Delta z \rightarrow 0$$

Note

$$f(x + \Delta x, y + \Delta y) - f(x, y) = \left(\frac{\partial f}{\partial x} \right) \Delta x + \left(\frac{\partial f}{\partial y} \right) \Delta y + \dots$$

Check

$$\begin{aligned} e^{z+\Delta z} &= e^z e^{\Delta z} = e^z (1 + \Delta z + \dots) \\ e^{z+\Delta z} - e^z &= e^z \delta z + \dots = f''(z) \delta z + \dots \end{aligned}$$

Therefore, $e^{z+\Delta z}$ is *analytic*. However,

$$\bar{z} + \Delta \bar{z} - \bar{z} = \bar{\Delta} z$$

$$\Delta z = \Delta x \rightarrow \bar{\Delta} z = (1) \Delta z \quad \text{Horizontal}$$

$$\Delta z = i \Delta y \rightarrow \bar{\Delta} z = -i \Delta y = (-1) \Delta z \quad \text{Vertical}$$

$$f(z) = u + iv$$

$$\frac{f(x + \Delta x, y) - f(x)}{\Delta x} = f'(z)$$

$$\frac{f(x, y + \Delta y) - f(x, y)}{i\Delta y} = f'(z) = \frac{\partial f}{\partial y} \frac{1}{i}$$

$$\begin{aligned} \frac{\partial f}{\partial x} &= \frac{1}{i} \frac{\partial f}{\partial y} \\ \left(\frac{\partial u}{\partial x} \right) + i \left(\frac{\partial v}{\partial x} \right) &= \frac{1}{i} \left[\left(\frac{\partial u}{\partial y} \right) + i \left(\frac{\partial v}{\partial y} \right) \right] \\ &= \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y} \end{aligned}$$

Cauchy Riemann Equations

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial v}{\partial y} \\ \frac{\partial u}{\partial y} &= -\frac{\partial v}{\partial x} \end{aligned}$$

Example

$$f(z) = x^2 - y^2 + 2ixy$$

$$\begin{aligned} u &= x^2 - y^2 \\ v &= 2xy \end{aligned}$$

$$\frac{\partial u}{\partial x} = 2x = \frac{\partial v}{\partial y}$$

$$\frac{\partial u}{\partial y} = -2y = -\frac{\partial v}{\partial x}$$




Figure 10: a region in the complex plane

3.4.0.3 Taylor Series

$$f(z) = \sum_{n=0}^{\infty} a_n z^n \quad \text{converges if analytic}$$

Given a constant λ

$$0 < \lambda < 1$$

$$f(\lambda z) = \sum_{n=0}^{\infty} a_n (\lambda z)^n \quad \text{definitely converges}$$

$$|a_n z^n \lambda^n| = |a_n z^n| |\lambda|^n$$

Let $\lambda = r e^{i\theta}$, $0 < r < 1$, $|\lambda| < 1$




Figure 11: disk of convergence of power series



Figure 12: disk of convergence is disk with maximum radius inside R

Example

$$f(x) = \frac{1}{x^2 + 1}$$

$$f(z) = \frac{1}{z^2 + 1}$$




Figure 13: using complex number instead of real numbers to calculate interval of convergence

Example

$$if(z) = \log(z)$$




Figure 14: graph of complex log to determine convergence

3.4.0.4 Path Integrals

$$\int_{-\infty}^{\infty} \left(\frac{\sin x}{x} \right) dx = \pi$$

$$\int_{-\infty}^{\infty} \frac{dx}{x^2 + 1} = \arctan x \Big|_{-\infty}^{\infty} = \pi$$




Figure 15: path P

Technique

Parametrize P from $0 < t < \pi$ as $z(t)$




Figure 16: example of parameterizing $z(t)$

$$\int_P f(z) dz = \int_a^b f(z(t)) \left(\frac{dz}{dt} \right) dt$$

$$f(z) = z^3$$

$$\frac{dz}{dt} = iRe^{it}$$

$$f(z(t)) = (Re^{it})^3 = R^3 e^{3it}$$

Collect

$$\begin{aligned}\int_0^\pi f(z(t)) \left(\frac{dz}{dt} \right) dt &= \int_0^\pi R^3 e^{3it} (iR e^{it}) dt \\ &= iR^4 \int_0^\pi e^{4it} dt = \frac{iR^4}{4i} e^{4it} \Big|_0^\pi = 0 \\ e^{4\pi i} &= 1\end{aligned}$$




Figure 17: contour integral path

3.4.0.5 Contour Integrals Contour Integrals = 0 for any analytic function that is analytic for the entire region C

$$\begin{aligned}\oint_C f(z) dz &= 0 \\ &= \int_0^{2\pi} R^n e^{nit} \frac{dz}{dt} dt \\ &= \int_0^{2\pi} R^n e^{nit} iR e^{it} dt \\ &= iR^{n+1} \int_0^{2\pi} e^{i(n+1)t} dt \\ &= \frac{iR^{n+1}}{i(n+1)} e^{i(n+1)t} \Big|_0^{2\pi} = 0\end{aligned}$$

Effect of Singularities

Let $n = -1$

$$\begin{aligned}f(z) &= \frac{1}{z} \\ \oint_C \frac{dz}{z} &= iR^{n+1} \int_0^{2\pi} e^{i(n+1)t} dt \\ &= i \int_0^{2\pi} dt = 2\pi i \neq 0\end{aligned}$$




Figure 18: Fundamental Theorem of Contour Integrals

$$\int_a^b f(z) dz = F(b) - F(a)$$

Example

$$f(z) = \frac{1}{z^2 + 1} \quad \alpha = i$$

$$= \frac{1}{(z - i)(z + i)} = \frac{1}{z - i} \left(\frac{1}{z + i} \right)$$

$$f(z) = \frac{g(z)}{z - \alpha}$$

$$\begin{aligned} f(z) &= \frac{1}{z - \alpha} [g(\alpha) + (z - \alpha)g'(\alpha) + \dots] \\ &= \frac{g(\alpha)}{z - \alpha} + g'(\alpha) + \dots \end{aligned}$$

$$\oint_C f(z) dz = \oint_C' f(z) dz = \oint_D f(z) dz = \oint \frac{g(\alpha)}{z - \alpha} dz$$