

Problem

Use the recursion theorem to give an alternative proof of Rice's theorem in Problem 5.28.

Problem 5.28

Rice's theorem. Let P be any nontrivial property of the language of a Turing machine. Prove that the problem of determining whether a given Turing machine's language has property P is undecidable.

In more formal terms, let P be a language consisting of Turing machine descriptions where P fulfills two conditions. First, P is nontrivial—it contains some, but not all, TM descriptions. Second, P is a property of the TM's language—whenever

$L(M_1) = L(M_2)$, we have $\langle M_1 \rangle \in P$ iff $\langle M_2 \rangle \in P$. Here, M_1 and M_2 are any

TMs. Prove that P is an undecidable language.

Step-by-step solution

Step 1 of 2

Rice's theorem: Let P be any nontrivial property of the language of a Turing machine.

In more formal terms, Let P be a Language consisting of Turing machine descriptions where P fulfills two conditions.

- First, P is nontrivial – It contains some, but not all, TM descriptions.

- Second, P is a property of the TM 's language- whenever $L(M_1) = L(M_2)$, we have $\langle M_1 \rangle \in P$ iff $\langle M_2 \rangle \in P$. Here M_1 and M_2 are any TMs .

Then we have to prove that P is an undecidable Language.

Assume that P is a decidable Language satisfying the Properties of Rice theorem.

[Comment](#)

Step 2 of 2

Let the Turing machine T decides the language P . According to the first condition of Rice theorem, P contains some, but not all, TM descriptions.

From above properties we assume TM_1 & TM_2 be the Turing machines and $\langle TM_1 \rangle \in P$ and $\langle TM_2 \rangle \notin P$.

Now construct the Turing machine TM_3 using TM_1 & TM_2 .

TM_3 on input w :

1. By using recursion theorem, construct our own description for $\langle TM_3 \rangle$

2. Run T on Turing machine $\langle TM_3 \rangle$

3. If T accepts $\langle TM_3 \rangle$, simulate $\langle TM_2 \rangle$ on w .

If T rejects $\langle TM_3 \rangle$, simulate $\langle TM_1 \rangle$ on w .

- Suppose $\langle TM_3 \rangle \in P$

Then T accepts $\langle TM_3 \rangle$ and $L(TM_3) = L(TM_2)$

According to second condition of Rice's theorem $\langle TM_2 \rangle \in P$ but our condition is $\langle TM_2 \rangle \notin P$. It is a contradiction here.

- So, we will get similar contradiction if $\langle TM_3 \rangle \notin P$.

As a result of that our assumption is wrong for P is a decidable language. Hence every property satisfying the conditions of Rice's theorem is undecidable.

[Comment](#)