

Lie Groups

Devesh Rajpal

Contents

1	4th January 23	1
2	9th Jan 2023	1
3	11th Jan 2023	3
4	18 Jan 2023	5
5	23 Jan 2023	7
6	25 Jan 2023	8
7	30 Jan 2023	10

1 4th January 23

One can study Lie Groups from several points of view. The course is aimed to understand the structure of Lie Groups.

Definition 1.1. A smooth manifold M is a Hausdorff space which is locally Euclidean with a smooth atlas i.e. (i) given any $x \in M$, \exists a chart (U, ϕ) , $x \in U \subset M$ with $\phi : U \rightarrow \phi(U)$ open in \mathbb{R}^m .

(ii) We have collection $\{(U, \phi)\}$ of charts such that

$$\phi(U \cap V) \xrightarrow{\psi \circ \phi^{-1}} \psi(U \cap V)$$

is a diffeomorphism.

Suppose $f : M \rightarrow N$ is a continuous map between manifolds. We say that f is smooth if for $(U, \phi) \in \mathcal{H}(M)$, $(V, \psi) \in \mathcal{H}(N)$ such that $f(U) \subset V$ and $\psi \circ f \circ \phi^{-1}$ is smooth.

TO DO : Construction of tangent bundle and vector bundle

2 9th Jan 2023

Definition 2.1. G is a Lie group if

1. G is a smooth manifold
2. G is also a group s.t

$$\begin{aligned}\mu : G \times G &\rightarrow G \\ (g, h) &\mapsto gh\end{aligned}$$

and

$$\begin{aligned}i : G &\rightarrow G \\ g &\mapsto g^{-1}\end{aligned}$$

are smooth maps.

Definition 2.2. A real (or complex) vector space V together with a bilinear map

$$[,] : V \times V \rightarrow V$$

is called a **Lie Algebra** if

1. $[X, Y] = -[Y, X]$ - skew symmetry
2. $[[X, Y], Z] + [[Y, Z], X] + [[Z, X], Y] = 0$ - Jacobi identity

Example. 1. $(\mathbb{R}, +), (\mathbb{C}, +), V$ any f.d vector space over \mathbb{R} or \mathbb{C} .

2. $(\mathbb{R}^\times, \cdot), (\mathbb{C}^\times, \cdot)$

3. $S^1 = \{z \in \mathbb{C}^\times \mid |z| = 1\}$

4. $\mathrm{GL}_n(\mathbb{R}), \mathrm{GL}_n(\mathbb{C})$

5. $\mathbb{R}^n / \mathbb{Z}^n \cong (\mathbb{R}^n / \mathbb{Z}^n) \cong (S^1)^n$

6. Suppose $\Gamma \subset V$ is a discrete subgroup. Then V/Γ is a Lie group.

7. N = unipotent upper triangular matrices, B = upper triangular matrices. As manifolds $N \cong \mathbb{R}^{\binom{n}{2}}$ and $B \cong (\mathbb{R}^\times)^n \times N$.

8. $\mathrm{SL}_n(\mathbb{R}) = \{X \in \mathrm{GL}_n(\mathbb{R}) \mid \det X = 1\}, \mathrm{SL}_n(\mathbb{C})$.

9. $O(n), SO(n)$.

10. $U(n), SU(n)$.

11. \mathbb{H}^\times, S^3 with quaternion multiplication.

12. $Sp(n) = \{X \in \mathrm{GL}_n(\mathbb{R}) \mid X \text{ preserves quaternion structure as a subset of } \mathrm{Aut}_{\mathbb{H}} \mathbb{H}^n\}$

Problem. $V/\Gamma \cong \mathbb{R}^k \times (S^1)^{n-k}$ for n -dimensional vector space V .

Theorem 2.1. Suppose G is a compact, connected, simple Lie group. Then G is locally isomorphic to

1. $SU(n), n \geq 2$ denoted by A_{n-1}
2. $SO(2n+1), n \geq 2$ denoted by B_n
3. $Sp(n), n \geq 1$ denoted by C_n
4. $SO(2n), n \geq 2$ denoted by D_n

or one of the following exceptional Lie group G_2, F_4, E_6, E_7, E_8 .

Problem. Prove that $SL_n(\mathbb{R})$ and $O(n)$ are smooth manifold, hence Lie groups.

Examples of Lie algebra -

- Example.**
1. $(V, [\cdot, \cdot] \equiv 0)$ is called trivial Lie algebra.
 2. $(\mathfrak{gl}_n(\mathbb{R}), [A, B] = AB - BA), \mathfrak{gl}_n(\mathbb{C})$
 3. $\mathfrak{sl}_n(\mathbb{R}) (\mathfrak{sl}_n(\mathbb{C}))$ is the Lie subalgebra of $\mathfrak{gl}_n(\mathbb{R}) (\mathfrak{gl}_n(\mathbb{C}))$ consisting of trace 0.
 4. \mathfrak{so}_n is Lie subalgebra of $\mathfrak{gl}_n(\mathbb{R})$ consisting of skew-symmetric matrices.

Definition 2.3. A vector field X on a Lie group G is called left invariant if $(L_g)_*(X_h) = X_{gh}$

3 11th Jan 2023

Recall $\mathbb{H} = \{a + bi + cj + dk : (a, b, c, d) \in \mathbb{R}^4, i^2 = -1, j^2 = -1, k^2 = -1, ij = k, jk = i, ki = j\}$ is the quaternion division algebra with the norm

$$\|a + bi + cj + dk\|^2 = a^2 + b^2 + c^2 + d^2$$

which satisfies $\|q_1 \cdot q_2\| = \|q_1\| \cdot \|q_2\|$

We can put this multiplication on $S^3 \cong SU(2)$ to get a compact Lie group. To get the isomorphism $SU(2) \cong S^3$, we define a map

$$\begin{aligned} \phi : S^3 &\rightarrow SU(2) \\ (a, b, c, d) &\mapsto \begin{bmatrix} a + bi & c + di \\ -(c - di) & a - bi \end{bmatrix} \end{aligned}$$

which is an algebra isomorphism.

Definition 3.1. The Lie algebra of G is the space of all left-invariant vector fields on G .

We have an isomorphism

$$\begin{aligned} \mathfrak{g} = \text{Lie}(G) &\rightarrow T_e G \\ X &\mapsto X_e \end{aligned}$$

Example. Let $G = \mathbb{R}^n$, with identity element $0 \in \mathbb{R}^n$ and left-invariant vector fields $\{\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\}$. Then the Lie bracket is

$$[\cdot, \cdot] \equiv 0$$

Remark. In general for any abelian Lie group G , the Lie bracket is $[\cdot, \cdot] \equiv 0$.

Theorem 3.1. Let G be a connected Lie group. Then

1. $\text{Lie}(G) = \mathfrak{g}$ is isomorphic as a vector space to $T_e(G)$.
2. Left-invariant vector fields are smooth.
3. $\text{Lie}(G)$ is closed under Lie bracket.

Proof. 1. Let X be a left-invariant vector field on G . We need to show that Xf is smooth for each $f \in C^\infty(G)$.

$$\begin{aligned} (Xf)(g) &= X_g f \\ &= (d\lambda_g X_e) f \\ &= X_e(f \circ \lambda_g) \end{aligned}$$

To show that Xf is smooth, it suffices to show that $X_e(f \circ \lambda_g)$ is smooth. We realize $X_e(f \circ \lambda_g)$ as evaluation of a smooth function on a smooth function.

Let Y be a smooth vector field on G such that $Y_e = X_e$

$$Y_e(f \circ \lambda_g) = X_e(f \circ \lambda_g)$$

We look at λ_g as the composition of

$$\begin{aligned} G &\xrightarrow{i_g^2} G \times G \xrightarrow{\mu} G \\ x &\mapsto (g, x) \mapsto gx \end{aligned}$$

Regard Y as the vector field $(0, Y)$ on $G \times G$. Now

$$\begin{aligned} (0, Y)(f \circ \mu) \circ i_e^1(g) &= (0, Y)_{(g, e)}(f \circ \mu) \\ &= 0_g(f \circ \mu \circ i_g^1) + Y_e(f \circ \mu \circ i_g^2) \\ &= Y_e(f \circ \lambda_g) \end{aligned}$$

which proves the smoothness.

2. Let X, Y left-invariant vector fields on G . We must show that $[X, Y]$ is a left-invariant

vector field.

$$\begin{aligned}
d\lambda_g([X, Y]_e)f &= [X, Y]_g f \\
&= [X, Y]_e(f \circ \lambda_g) \\
&= X_e(Y(f \circ \lambda_g)) - Y_e(X(f \circ \lambda_g)) \\
&= X_e(d\lambda_g(Yf)) - Y_e(d\lambda_g(Xf))
\end{aligned}$$

□

4 18 Jan 2023

Lemma 4.1. Suppose $\psi : M \rightarrow N$ is a smooth map. Let X_1, X_2 be vector fields on M , Y_1, Y_2 be vector fields on N such that X_i is ψ -related to Y_i . Then $[X_1, X_2]$ is ψ -related to $[Y_1, Y_2]$.

Proof. Notice that

$$\begin{aligned}
d\psi[X_1, X_2](f) &= [X_1, X_2](f \circ \psi) \\
&= X_1(X_2 f \circ \psi) - X_2(X_1 f \circ \psi) \\
&= X_1(d\psi X_2 f) - X_2(Y_1 f \circ \psi) \\
&= X_1(Y_2 f \circ \psi) - X_2(Y_1 f \circ \psi) \\
&= d\psi X_1(Y_2 f) - d\psi X_2(Y_1 f) \\
&= Y_1 Y_2 f \circ \psi - Y_2 Y_1 f \circ \psi \\
&= [Y_1, Y_2](f) \circ \psi
\end{aligned}$$

□

This lemma proves that the set of left-invariant vector fields forms a Lie algebra.

Consider the Lie group $\mathrm{GL}_n(\mathbb{R})$. We want to verify the Lie algebra structure on $\mathfrak{gl}_n(\mathbb{R}) = M_n(\mathbb{R})$ with the isomorphism

$$\begin{aligned}
\mathrm{Lie}(\mathrm{GL}_n(\mathbb{R})) &\rightarrow \mathfrak{gl}_n(\mathbb{R}) \\
X &\xrightarrow{\beta} X_e
\end{aligned}$$

Lemma 4.2.

$$\beta([X, Y]) = [\beta(X), \beta(Y)]$$

Proof. Evaluating the bracket on coordinate function x_{ij} .

$$[X, Y]_e(x_{ij}) = X_e(Y(x_{ij})) - Y_e(X(x_{ij})) \quad (1)$$

Now

$$\begin{aligned} Y(x_{ij})(g) &= d\lambda_g Y_e(x_{ij}) \\ &= Y_e(x_{ij} \circ \lambda_g) \\ &= \sum_k x_{ik}(g) Y_e(x_{kj}) \end{aligned}$$

Considering the above as function of g and substituting this in Eq. (1) we get

$$\begin{aligned} [X, Y]_e(x_{ij}) &= X_e Y_e(x_{ij}) - Y_e X_e(x_{ij}) \\ &= \sum_k \{X_e(x_{ik}) Y_e(x_{kj}) - Y_e(x_{ik}) X_e(x_{kj})\} \\ &= [X_e, Y_e](x_{ij}) \end{aligned}$$

□

Definition 4.1. A **Lie subgroup** H of a Lie group G is a submanifold $H \xrightarrow{\alpha} G$ where α is smooth and a group homomorphism.

We say that H is closed Lie subgroup if it is Lie subgroup such that $H \rightarrow \alpha(H)$ is a diffeomorphism.

Example. Consider the map $\mathbb{R} \rightarrow S^1 \times S^1$ given by

$$t \mapsto (e^{2\pi i t}, e^{2\pi i \sqrt{2} t})$$

The image is a Lie subgroup of $S^1 \times S^1$ but it is not a closed Lie subgroup. It is also known as “Skew-line” in the torus.

Definition 4.2. Let $\mathfrak{g}, \mathfrak{h}$ be Lie algebras and $f : \mathfrak{g} \rightarrow \mathfrak{h}$ be a vector space homomorphism. Then we say that f is a Lie algebra homomorphism if

$$f([X, Y]) = [f(X), f(Y)]$$

Theorem 4.3. Suppose that $\psi : G \rightarrow H$ is a Lie group homomorphism. Let X be a left-invariant vector field on G . Extend $d\psi(X_e) = Y_e \in T_e H$ to a left-invariant vector field Y on H . Then X and Y are ψ -related. This implies $d\psi_e : \mathfrak{g} \rightarrow \mathfrak{h}$ is a Lie algebra homomorphism.

Proof. Consider the commutative diagram

$$\begin{array}{ccc}
TG & \xrightarrow{d\psi} & TH \\
\downarrow X & & \downarrow Y \\
G & \xrightarrow{\psi} & H
\end{array}$$

We want to show that $Y \circ \psi = d\psi \circ X$. Now

$$\lambda_{\psi(g)} \circ \psi = \psi \circ \lambda_g$$

so

$$\begin{aligned}
Y_{\psi(g)} &= d\lambda_{\psi(g)} Y_e \\
&= d\lambda_{\psi(g)} d\psi X_e \\
&= d(\lambda_{\psi(g)} \circ \psi)(X_e) \\
&= d(\psi \circ \lambda_g)(X_e) \\
&= d\psi d\lambda_g(X_e) \\
&= d\psi X_g
\end{aligned}$$

□

Theorem 4.4. Let G, H be Lie groups with G connected. Let

$$\phi, \psi : G \rightarrow H$$

be homomorphism of Lie groups such that

$$d\phi = d\psi : T_e G \rightarrow T_e H$$

Then $\phi = \psi$.

5 23 Jan 2023

I missed the class. Regardless here are some definitions from Warner covered on this day.

Definition 5.1. Let M be a smooth d -dimensional manifold. For any integer $1 \leq c \leq d$, a **c -dimensional distribution** \mathcal{D} on manifold is a choice of c -dimensional subspace $\mathcal{D}_p \subset T_p M$. \mathcal{D} is smooth if for each $p \in M$ there is an open neighborhood U of p and there are c smooth vector fields X_1, \dots, X_c on U which span \mathcal{D}_m for each $p \in U$.

We say \mathcal{D} is **involutive** if $[X, Y] \in \mathcal{D}$ whenever $X, Y \in \mathcal{D}$.

Definition 5.2. A submanifold (N, ϕ) of M is an integral manifold of a distribution \mathcal{D} if

$$d\phi(N_p) = \mathcal{D}_{\phi(p)}$$

Suppose there exists an integral manifold N for a distribution \mathcal{D} , then for the points on N the distribution \mathcal{D} is necessarily involutive. Frobenius theorem states that it is sufficient condition for a distribution to be integral.

Theorem 5.1. (Frobenius) Let \mathcal{D} be a c -dimensional involutive smooth distribution on M . Then there exists an integral manifold of \mathcal{D} passing through each point of M .

Differential Ideals

Let $E^*(M) = \bigoplus_{i=0}^{\infty} E^i(M)$ denote the graded algebra of smooth differential forms over manifold M .

Definition 5.3. Let \mathcal{D} be a smooth p -dimensional distribution on M . A q -form ω is said to **annihilate** \mathcal{D} if for each $x \in M$

$$\omega_x(v_1, \dots, v_q) = 0 \quad \text{whenever } v_1, \dots, v_q \in \mathcal{D}_x$$

A form $\omega \in E^*(M)$ is said to annihilate \mathcal{D} if each of the homogenous components of ω annihilate \mathcal{D} . Define

$$\mathcal{I}(\mathcal{D}) \doteq \{\omega \in E^*(M) : \omega \text{ annihilates } \mathcal{D}\}$$

Definition 5.4. An ideal $\mathcal{I} \in E^*(M)$ is called a **differential ideal** if it is closed under exterior differentiation; i.e.

$$d(\mathcal{I}) \subset \mathcal{I}.$$

Theorem 5.2. A smooth distribution \mathcal{D} on M is involutive if and only if the ideal $\mathcal{I}(\mathcal{D})$ is a differential ideal.

6 25 Jan 2023

Theorem 6.1. If $\phi : H \rightarrow G$ is a homomorphism of Lie groups and if ω is a left-invariant differential form on G , then $\phi^*(\omega)$ is again a left-invariant form on H .

Suppose that $\phi : H \rightarrow G$ is a homomorphism of Lie groups. Let $\omega_1, \dots, \omega_d$ be a basis for $E_{\text{inv}}^1(G)$. Then

$$\mathcal{I}_\phi = \langle \{\pi_1^* \phi^*(\omega_j) - \pi_2^*(\omega_j)\} : 1 \leq j \leq d \rangle$$

is a left-invariant differential ideal of $H \times G$.

Lemma 6.2. Suppose X_1, \dots, X_d is a basis of \mathfrak{g} dual to $\omega_1, \dots, \omega_d$. Suppose the Lie bracket is given by

$$[X_i, X_j] = \sum c_{ij}^k X_k$$

Then the C^∞ functions c_{ij}^k are constant. Further,

$$d\omega_i = -c_{kj}^i \omega_k \wedge \omega_j$$

Proof. Notice that

$$\begin{aligned} d\omega_k(X_i, X_j) &= -\omega_k([X_i, X_j]) \\ &= -c_{ij}^k \end{aligned}$$

which is a constant because a left-invariant 1-form evaluated on a left-invariant vector field is a constant. \square

Remark. The c_{ij}^k are called the structural constants of G with respect to the basis $\{X_i\}$ of \mathfrak{g} .

Proof. Theorem 4.4. Notice that $\mathcal{I}_\psi = \mathcal{I}_\phi$ since $d\phi = d\psi$ and these are invariant differential ideals; hence integral manifolds of \mathcal{I}_ϕ and \mathcal{I}_ψ passing through (e, e) are the same. Thus, $\phi = \psi$. \square

Lemma 6.3. Suppose G is any Hausdorff topological group which is connected. Suppose $e \in U \subset G$ is any open set. Then

$$G = \bigcup_{n \geq 1} U^n$$

where $U^n = \{x_1 \cdots x_n | x_i \in U\}$

Proof. Since $e \in U$ is open, $U^{-1} = \{x^{-1} | x \in U\}$ is also an open neighborhood of e . Let $V = U \cap U^{-1}$. Note that

$$H \doteq \bigcup_{n \geq 1} V^n$$

is a subgroup of G , and it is open. Since the cosets gH are also open it follows that $G = \cup_g gH$ being connected must be H . \square

Theorem 6.4. Let G be a Lie group and $\mathfrak{h} \subset \mathfrak{g}$ be a Lie subalgebra of \mathfrak{g} . Then there exists connected Lie subgroup H of G such that $T_e H = \mathfrak{h}$.

Proof. Consider the distribution \mathcal{D} defined as

$$\mathcal{D}_g = \{X_g | X \in \mathfrak{h}\}$$

on G . Suppose X_1, \dots, X_c is a basis of \mathfrak{h} . Then \mathcal{D} is generated by X_1, \dots, X_c and \mathcal{D} is involutive. \square

Corollary. (a) There is a one-to-one correspondence between connected Lie subgroups of G and Lie subalgebras of \mathfrak{g} .

(b) Suppose $(H, i) \leftrightarrow \mathfrak{h} \subset \mathfrak{g}$. Then (H, i) is an embedded manifold if and only if H is closed.

Theorem 6.5. Suppose that $A \subset G$ is an abstract subgroup of G and if A has a manifold structure such that $(A, i) \rightarrow G$ is a submanifold. Then the manifold structure is unique, A is a Lie group and hence (A, i) is a Lie subgroup of G .

Theorem 6.6. (Adó) Suppose that \mathfrak{g} is a finite dimensional Lie algebra. Then \mathfrak{g} can be realized as a subalgebra of $\mathfrak{gl}(n, \mathbb{R})$.

Given any connected Lie group G , it has a universal cover $\tilde{G} \xrightarrow{\pi} G$. Choose $\tilde{e} \in \pi^{-1}(e) \in \tilde{G}$ such that the following diagram

$$\begin{array}{ccc} \tilde{G} \times \tilde{G} & \longrightarrow & \tilde{G} \\ \pi \times \pi \downarrow & & \downarrow \pi \\ G \times G & \longrightarrow & G \end{array}$$

commutes.

7 30 Jan 2023

Lemma 7.1. Suppose that G is a connected Lie group. Then $\pi_1(G)$ is abelian.

Proof. Suppose $\sigma, \tau : I \rightarrow G$ be two loops. Define $\sigma \cdot \tau$ by

$$(\sigma \cdot \tau)(s) = \sigma(s) \cdot \tau(s)$$

Then we have

$$\sigma * \tau \cong \sigma \cdot \tau$$

where $*$ denote the product in the fundamental group $\pi_1(G)$ (given by concatenation) and \cong denotes equivalent in homotopy. Also

$$\sigma \cdot \tau \cdot \sigma^{-1} \cong \tau$$

which implies $\sigma\tau \cong \tau \cdot \sigma$ \square

Theorem 7.2. Suppose that G and H are Lie groups with Lie algebras \mathfrak{g} and \mathfrak{h} with G simply connected. Let $\tilde{\phi} : \mathfrak{g} \rightarrow \mathfrak{h}$ be a Lie algebra homomorphism. Then there exists a Lie group homomorphism

$$\phi : G \rightarrow H$$

such that $d\phi_e : T_e(G) = \mathfrak{g} \rightarrow \mathfrak{h} = T_eH$ is equal to $\tilde{\phi}$.

Proof. Let $\{\omega_i\}$ be a basis for invariant differential forms in $E^1(H)$. Let \mathcal{I} be the ideal generated by $\{\pi_1^*\tilde{\phi}^*(\omega_j) - \pi_2^*(\omega_j) | 1 \leq j \leq d\}$. Then \mathcal{I} is an invariant differential ideal of $G \times H$ so it comes from vanishing of an integrable submanifold of $G \times H$ passing through (e, e) .

Then M is a Lie subgroup of $G \times H$ and $M \xrightarrow{p} G$ obtained by restriction of π_1 is a group homomorphism and also a local diffeomorphism. So $p : M \rightarrow G$ is a covering projection but G is simply connected so p is a diffeomorphism

$$G \xrightarrow{p^{-1}} M \hookrightarrow G \times H \rightarrow H.$$

□

Corollary. 1. Suppose $\mathfrak{g} \cong \mathfrak{h}$ as Lie algebras and G and H are simply connected. Then $G \cong H$ as Lie groups.

2. There exists a one-to-one correspondence between (finite dimensional) Lie algebras and simply connected Lie groups.
3. The differential structure of a Lie group is determined by its Lie algebra.

If G is a topological group which is locally Euclidean, does it support a Lie group structure? The answer is yes but the proof is quite difficult.

Exponential map

Let X be a left-invariant vector field on G . We have a Lie algebra homomorphism

$$\begin{aligned} \text{Lie}(\mathbb{R}) &\cong \mathbb{R} \rightarrow \mathfrak{g} \\ c \frac{d}{dt} &\rightarrow cX \end{aligned}$$

This yields a Lie group homomorphism

$$\begin{aligned} \mathbb{R} &\xrightarrow{\exp_X} G \\ x &\mapsto \exp_X(x) \end{aligned}$$

then $d\exp_X(c \frac{d}{dt}) = cX$. The map

$$\begin{aligned} \mathfrak{g} &\xrightarrow{\exp} G \\ X &\mapsto \exp_X(1) \end{aligned}$$

is called the **exponential map**.

Theorem 7.3. Let $X \in \text{Lie}(G)$. Then

1. $\exp(tX) = \exp_X(t)$
2. $\exp(t_1X_1 + t_2X) = \exp(t_1X) \cdot \exp(t_2X)$
3. $\exp(-tX) = (\exp(tX))^{-1}$
4. $\exp : \mathfrak{g} \rightarrow G$ is smooth and $d\exp : T_0\mathfrak{g} \rightarrow T_eG = \mathfrak{g}$ is the identity map
5. $\lambda_g \circ \exp_X : \mathbb{R} \rightarrow G$ is the unique integral curve of X which is based at g .
6. The left-invariant vector fields are complete, i.e. their integral curves exist for all time.
7. The one-parameter group of diffeomorphism $\psi_{X,t}$ for $t \in \mathbb{R}$ is given by

$$\psi_{X,t} = \rho_{\exp_X(t)}$$

where ρ_g denote right-multiplication by g .

Theorem 7.4. Suppose $\psi : H \rightarrow G$ is a Lie group homomorphism. Then

$$\begin{array}{ccc} \mathfrak{h} & \xrightarrow{d\psi} & \mathfrak{g} \\ \exp \downarrow & & \downarrow \exp \\ H & \xrightarrow{\psi} & G \end{array}$$

commutes. Also

$$\mathbb{R} \xrightarrow{\exp} \mathfrak{h} \quad \mathfrak{g}$$

$$H \quad G$$

[DO THIS COMMUTATIVE DIAGRAM.]