

# MA 205 Complex Analysis: Some More Theorems

September 2, 2017

# Examples

Recall the following example from last time :

Show that  $\int_0^\infty \frac{x^{-c}}{1+x} dx = \frac{\pi}{\sin \pi c}$  if  $0 < c < 1$ .

# Examples

Recall the following example from last time :

Show that  $\int_0^\infty \frac{x^{-c}}{1+x} dx = \frac{\pi}{\sin \pi c}$  if  $0 < c < 1$ .

We'll integrate

$$f(z) = \frac{z^{-c}}{1+z},$$

where  $z^{-c}$  is the branch corresponding to branch cut being the positive real axis, and integrate along the contour:

# Examples


Recall the following example from last time :

Show that  $\int_0^\infty \frac{x^{-c}}{1+x} dx = \frac{\pi}{\sin \pi c}$  if  $0 < c < 1$ .

We'll integrate

$$f(z) = \frac{z^{-c}}{1+z},$$

where  $z^{-c}$  is the branch corresponding to branch cut being the positive real axis, and integrate along the contour:



By residue theorem,

$$\int_{\gamma} \frac{z^{-c}}{1+z} dz = 2\pi i e^{-i\pi c}.$$

# Real Integrals

By residue theorem,

$$\int_{\gamma} \frac{z^{-c}}{1+z} dz = 2\pi i e^{-i\pi c}.$$

The integral is the sum of four integrals; one on  $L_1$ , one on  $\gamma_R$ , one on  $L_2$ , one on  $\gamma_r$ .

# Real Integrals

By residue theorem,

$$\int_{\gamma} \frac{z^{-c}}{1+z} dz = 2\pi i e^{-i\pi c}.$$

The integral is the sum of four integrals; one on  $L_1$ , one on  $\gamma_R$ , one on  $L_2$ , one on  $\gamma_r$ . Note that

$$\int_r^R \frac{t^{-c}}{1+t} dt = \lim_{\delta \rightarrow 0} \int_{L_1} \frac{z^{-c}}{1+z} dz.$$

Similarly,

$$\lim_{\delta \rightarrow 0} \int_{L_2} \frac{z^{-c}}{1+z} dz = -e^{-2\pi i c} \int_r^R \frac{t^{-c}}{1+t} dt.$$

Also,

$$\left| \int_{\gamma_\rho} \frac{z^{-c}}{1+z} dz \right| \leq \frac{\rho^{-c}}{|1-\rho|} 2\pi\rho.$$



Also,

$$\left| \int_{\gamma_\rho} \frac{z^{-c}}{1+z} dz \right| \leq \frac{\rho^{-c}}{|1-\rho|} 2\pi\rho.$$

This is zero in the limit as  $\rho \rightarrow 0$  or  $\rho \rightarrow \infty$ . Thus we get:

$$2\pi i e^{-i\pi c} = (1 - e^{-2i\pi c}) \int_0^\infty \frac{t^{-c}}{1+t} dt.$$

Also,

$$\left| \int_{\gamma_\rho} \frac{z^{-c}}{1+z} dz \right| \leq \frac{\rho^{-c}}{|1-\rho|} 2\pi\rho.$$

This is zero in the limit as  $\rho \rightarrow 0$  or  $\rho \rightarrow \infty$ . Thus we get:

$$2\pi i e^{-i\pi c} = (1 - e^{-2i\pi c}) \int_0^\infty \frac{t^{-c}}{1+t} dt.$$

Thus,

$$\int_0^\infty \frac{t^{-c}}{1+t} dt = \frac{2\pi i e^{-i\pi c}}{1 - e^{-2i\pi c}} = \frac{\pi}{\sin \pi c}.$$

# Real Integral

Integrate  $I = \int_{-\infty}^{\infty} \frac{e^{x/2} dx}{\cosh x}$

In this case  $\cosh x$  has infinitely many poles along the imaginary axis, namely at  $z = i(\pi/2 + n\pi)$ ,  $n \in \mathbb{Z}$  and so we do not choose the previous kind of contours. Instead we choose a rectangular contour  $\gamma$  consisting of vertices  $L, -L, L + i\pi$  and  $-L + i\pi$ .

# Real Integral

Integrate  $I = \int_{-\infty}^{\infty} \frac{e^{x/2} dx}{\cosh x}$

In this case  $\cosh x$  has infinitely many poles along the imaginary axis, namely at  $z = i(\pi/2 + n\pi)$ ,  $n \in \mathbb{Z}$  and so we do not choose the previous kind of contours. Instead we choose a rectangular contour  $\gamma$  consisting of vertices  $L, -L, L + i\pi$  and  $-L + i\pi$ .

By residue theorem,  $\int_{\gamma} \frac{e^{z/2} dz}{\cosh z} = 2\pi i \operatorname{Res}(f, i\frac{\pi}{2}) = 2\pi e^{i\frac{\pi}{4}}$ .

# Real Integral

$$\text{Integrate } I = \int_{-\infty}^{\infty} \frac{e^{x/2} dx}{\cosh x}$$

In this case  $\cosh x$  has infinitely many poles along the imaginary axis, namely at  $z = i(\pi/2 + n\pi)$ ,  $n \in \mathbb{Z}$  and so we do not choose the previous kind of contours. Instead we choose a rectangular contour  $\gamma$  consisting of vertices  $L, -L, L + i\pi$  and  $-L + i\pi$ .

By residue theorem,  $\int_{\gamma} \frac{e^{z/2} dz}{\cosh z} = 2\pi i \text{Res}(f, i\frac{\pi}{2}) = 2\pi e^{i\frac{\pi}{4}}$ .

$$\text{Now } |\cosh(L + iy)| = |e^{L+iy} + e^{-L-iy}|/2 \geq \frac{1}{2}(|e^{L+iy}| - |e^{-L-iy}|) = (e^L - e^{-L})/2 \geq e^L/4$$

# Real Integral

$$\text{Integrate } I = \int_{-\infty}^{\infty} \frac{e^{x/2} dx}{\cosh x}$$

In this case  $\cosh x$  has infinitely many poles along the imaginary axis, namely at  $z = i(\pi/2 + n\pi)$ ,  $n \in \mathbb{Z}$  and so we do not choose the previous kind of contours. Instead we choose a rectangular contour  $\gamma$  consisting of vertices  $L, -L, L + i\pi$  and  $-L + i\pi$ .

By residue theorem,  $\int_{\gamma} \frac{e^{z/2} dz}{\cosh z} = 2\pi i \text{Res}(f, i\frac{\pi}{2}) = 2\pi e^{i\frac{\pi}{4}}$ .

$$\text{Now } |\cosh(L + iy)| = |e^{L+iy} + e^{-L-iy}|/2 \geq \frac{1}{2}(|e^{L+iy}| - |e^{-L-iy}|) = (e^L - e^{-L})/2 \geq e^L/4$$

From this it follows from the ML-inequality that as  $L$  tends to  $\infty$ , the integral along the right vertical side tends to zero. Similarly one checks that the integral along the left vertical side also tend to zero.

## Example cont ..

Now since  $\cosh(x + i\pi) = -\cosh x$ , the integrals along the horizontal sides are related by

$$\int_L^{-L} \frac{e^{(x+i\pi)/2} dx}{\cosh(x + i\pi)} = e^{i\pi/2} \int_{-L}^L \frac{e^{x/2} dx}{\cosh x}$$

## Example cont ..

Now since  $\cosh(x + i\pi) = -\cosh x$ , the integrals along the horizontal sides are related by

$$\int_L^{-L} \frac{e^{(x+i\pi)/2} dx}{\cosh(x + i\pi)} = e^{i\pi/2} \int_{-L}^L \frac{e^{x/2} dx}{\cosh x}$$

Taking  $L$  tending to  $\infty$ , we see that

$$I = \frac{2\pi e^{i\pi/4}}{(1+e^{i\pi/2})} = \frac{\pi}{\cos(\pi/4)} = \pi\sqrt{2}.$$



# Choice of contour for integration

It might be a bit of a mystery as to which contour one should consider for a given contour integral. Here is a general recipe.

# Choice of contour for integration

It might be a bit of a mystery as to which contour one should consider for a given contour integral. Here is a general recipe.

Suppose the improper integral is of the form  $\int_{-\infty}^{\infty} f(x)dx$ . The general idea of course is to find a contour which contains the real line as part of the contour in the “limit”. The choice should be made so that by residue theory one knows the integral over the full contour and such that the integral over the extra added part goes to zero in the limit.

# Choice of contour for integration

It might be a bit of a mystery as to which contour one should consider for a given contour integral. Here is a general recipe.

Suppose the improper integral is of the form  $\int_{-\infty}^{\infty} f(x)dx$ . The general idea of course is to find a contour which contains the real line as part of the contour in the “limit”. The choice should be made so that by residue theory one knows the integral over the full contour and such that the integral over the extra added part goes to zero in the limit.

I. For instance suppose there exists a constant  $C$  such that  $|f(z)| \leq \frac{C}{|z|^r}$  for sufficiently large  $|z|$  and for some  $r > 1$  (here  $f(z)$  is an extension of  $f(x)$  to a function of the complex variable).

Note that this happens for instance in the case when  $f(x) = P(x)/Q(x)$  where  $\deg Q(x) \geq \deg P(x) + 2$ . Then close up the interval with a semicircle into the upper half plane and integrate along the contour and take limit as the radius of semicircle goes to infinity. Use ML inequality to show that the integral along the semicircle goes to zero as radius goes to  $\infty$ .

# Choice of contour for integration

In case the integral is from 0 to  $\infty$ , try and relate it to some integral from  $-\infty$  to  $\infty$ . For instance the function may have a natural continuation to the negative reals. In case this is not possible, often because  $f(x)$  has a singularity at origin; usually a pole, then try using a half annular region  $A(0; r, R)$  like we have done in earlier examples.

# Choice of contour for integration

In case the integral is from 0 to  $\infty$ , try and relate it to some integral from  $-\infty$  to  $\infty$ . For instance the function may have a natural continuation to the negative reals. In case this is not possible, often because  $f(x)$  has a singularity at origin; usually a pole, then try using a half annular region  $A(0; r, R)$  like we have done in earlier examples. This will avoid the pole and then show that in the limit as  $R$  tends to  $\infty$  and  $r$  tends to zero, the integrals along the larger and smaller circle tend to zero. Hence in the limit we will get integral over the real line. Then try and relate the integral over the positive real to that over the negative reals.

# Choice of contour for integration

In case the integral is from 0 to  $\infty$ , try and relate it to some integral from  $-\infty$  to  $\infty$ . For instance the function may have a natural continuation to the negative reals. In case this is not possible, often because  $f(x)$  has a singularity at origin; usually a pole, then try using a half annular region  $A(0; r, R)$  like we have done in earlier examples. This will avoid the pole and then show that in the limit as  $R$  tends to  $\infty$  and  $r$  tends to zero, the integrals along the larger and smaller circle tend to zero. Hence in the limit we will get integral over the real line. Then try and relate the integral over the positive real to that over the negative reals. See the example of  $\int_0^\infty \frac{\log(x)}{1+x^2}$  as an illustration of this. Otherwise, try a rectangular contour and show that the integral over the extra sides goes to zero in the limit.

# Choice of contour for integration

II. If the integrand is of the form  $\int_{-\infty}^{\infty} \frac{P(x)}{Q(x)} \sin(x) dx$ , where  $P(x)$  and  $Q(x)$  are polynomials with  $\deg(Q(x))$  at least one more than that of  $P(x)$ , close up the interval by the semicircular region in the upper half plane and use Jordan's lemma to show that the integral over the semicircle goes to zero using Jordan's lemma. (We have seen this when we integrated  $\sin(x)/x$ ).

# Choice of contour for integration

II. If the integrand is of the form  $\int_{-\infty}^{\infty} \frac{P(x)}{Q(x)} \sin(x) dx$ , where  $P(x)$  and  $Q(x)$  are polynomials with  $\deg(Q(x))$  at least one more than that of  $P(x)$ , close up the interval by the semicircular region in the upper half plane and use Jordan's lemma to show that the integral over the semicircle goes to zero using Jordan's lemma. (We have seen this when we integrated  $\sin(x)/x$ ).

III. If the integrand is of the type  $\int_0^{2\pi} P(\cos(t), \sin(t)) dt$ , set  $z = e^{it}$  and use  $\cos(t) = \frac{z+z^{-1}}{2}$  and  $\sin(t) = \frac{z-z^{-1}}{2i}$ .  $dt$  becomes  $\frac{dz}{iz}$  and then the integral assumes the form  $\int_{|z|=1} P\left(\frac{z+z^{-1}}{2}, \frac{z-z^{-1}}{2i}\right) \frac{dz}{iz}$  which can then be computed by using residue theorem.



# Choice of contour for integration

II. If the integrand is of the form  $\int_{-\infty}^{\infty} \frac{P(x)}{Q(x)} \sin(x) dx$ , where  $P(x)$  and  $Q(x)$  are polynomials with  $\deg(Q(x))$  at least one more than that of  $P(x)$ , close up the interval by the semicircular region in the upper half plane and use Jordan's lemma to show that the integral over the semicircle goes to zero using Jordan's lemma. (We have seen this when we integrated  $\sin(x)/x$ ).

III. If the integrand is of the type  $\int_0^{2\pi} P(\cos(t), \sin(t)) dt$ , set  $z = e^{it}$  and use  $\cos(t) = \frac{z+z^{-1}}{2}$  and  $\sin(t) = \frac{z-z^{-1}}{2i}$ .  $dt$  becomes  $\frac{dz}{iz}$  and then the integral assumes the form  $\int_{|z|=1} P\left(\frac{z+z^{-1}}{2}, \frac{z-z^{-1}}{2i}\right) \frac{dz}{iz}$  which can then be computed by using residue theorem.

IV. If the integrand has infinitely many poles going to infinity, you are usually better off using a rectangular contour which encompasses only finitely many poles.

# Choice of contour for integration

As before one tries to show that in the limit, the integral over the extra added vertical sides goes to zero in the limit and the integrals over the two horizontal sides are related ; usually proportional to each other. Thus taking limit as the length of the rectangular sides goes to infinity, one gets the desired answer.

# Choice of contour for integration

As before one tries to show that in the limit, the integral over the extra added vertical sides goes to zero in the limit and the integrals over the two horizontal sides are related ; usually proportional to each other. Thus taking limit as the length of the rectangular sides goes to infinity, one gets the desired answer.

V. In case the function involves a branch cut, choose a contour which avoids (goes around) the branch cut like in the earlier example.

# Choice of contour for integration

As before one tries to show that in the limit, the integral over the extra added vertical sides goes to zero in the limit and the integrals over the two horizontal sides are related ; usually proportional to each other. Thus taking limit as the length of the rectangular sides goes to infinity, one gets the desired answer.

V. In case the function involves a branch cut, choose a contour which avoids (goes around) the branch cut like in the earlier example.

The following rather obvious lemma is often useful for contour integration:

# Choice of contour for integration

As before one tries to show that in the limit, the integral over the extra added vertical sides goes to zero in the limit and the integrals over the two horizontal sides are related ; usually proportional to each other. Thus taking limit as the length of the rectangular sides goes to infinity, one gets the desired answer.

V. In case the function involves a branch cut, choose a contour which avoids (goes around) the branch cut like in the earlier example.

The following rather obvious lemma is often useful for contour integration:

**Fractional Residue Theorem :** Suppose  $z_0$  is a simple pole of  $f(z)$  and  $C_\delta$  is an arc of the circle  $|z - z_0| = \delta$  of angle  $\alpha$ , then

$$\lim_{\delta \rightarrow 0} \int_{C_\delta} f(z) dz = \alpha i \operatorname{Res}(f(z), z_0)$$

# Argument principle

The following is a useful variant of the residue theorem and is known as the argument principle. It is used to count zero's and poles of a meromorphic function on a domain.

# Argument principle

The following is a useful variant of the residue theorem and is known as the argument principle. It is used to count zero's and poles of a meromorphic function on a domain.

**Argument Principle** : Let  $\gamma$  be a closed contour contained in  $\mathbb{C}$  and let  $f(z)$  be a meromorphic function on an open set containing  $\gamma$  and its interior such that  $\gamma$  does not pass through any of the zeros and poles of  $f(z)$ . Then

$$\int_{\gamma} \frac{f'(z)}{f(z)} dz = N - P$$

where  $N$  and  $P$  denote the number of zero's and poles enclosed by  $\gamma$  with each zero and pole counted as many times as its order.

# Argument principle

The following is a useful variant of the residue theorem and is known as the argument principle. It is used to count zero's and poles of a meromorphic function on a domain.

**Argument Principle** : Let  $\gamma$  be a closed contour contained in  $\mathbb{C}$  and let  $f(z)$  be a meromorphic function on an open set containing  $\gamma$  and its interior such that  $\gamma$  does not pass through any of the zeros and poles of  $f(z)$ . Then

$$\int_{\gamma} \frac{f'(z)}{f(z)} dz = N - P$$

where  $N$  and  $P$  denote the number of zero's and poles enclosed by  $\gamma$  with each zero and pole counted as many times as its order.

**(Proof sketched on the board)**



# Rouche's Theorem

A nice and useful corollary of the argument principle is the following theorem:

# Rouche's Theorem

A nice and useful corollary of the argument principle is the following theorem:

**Rouche's Theorem:** Let  $\gamma$  be a simple closed contour and let  $f(z)$  and  $g(z)$  be two functions holomorphic on an open set containing  $\gamma$  and its interior. Suppose  $|f(z) - g(z)| < |f(z)|$  at all points on  $\gamma$ . Then  $\gamma$  encloses the same number of zero's of  $f(z)$  and  $g(z)$ .

# Rouche's Theorem

A nice and useful corollary of the argument principle is the following theorem:

**Rouche's Theorem:** Let  $\gamma$  be a simple closed contour and let  $f(z)$  and  $g(z)$  be two functions holomorphic on an open set containing  $\gamma$  and its interior. Suppose  $|f(z) - g(z)| < |f(z)|$  at all points on  $\gamma$ . Then  $\gamma$  encloses the same number of zero's of  $f(z)$  and  $g(z)$ . The proof of Rouché's theorem follows easily from the argument principle applied to the function  $f/g$  and noting that after cancelling common factors, the zeros of  $f$  (resp.  $g$ ) are the zero's (resp. poles) of  $f/g$ . I'll leave the details of the proof as an exercise.

# Example

Let us compute the number of zero's of  
 $f(z) = z^6 + 11z^4 + z^3 + 2z + 4$  inside the unit disc.

# Example

Let us compute the number of zero's of

$f(z) = z^6 + 11z^4 + z^3 + 2z + 4$  inside the unit disc.

Take  $g(z) = 11z^4$ . Then  $|g(z) - f(z)| < |g(z)|$  on the unit circle.

Hence  $g(z)$  has the same number of roots as  $f(z)$  inside the unit circle. But the number of roots of  $g(z)$  inside unit circle is 4 (counting multiplicity) which therefore equals number of roots of  $f(z)$ .

# Example

Lets count number of roots of  $f(z) = e^z - 2z - 1$  inside the unit circle.

# Example

Lets count number of roots of  $f(z) = e^z - 2z - 1$  inside the unit circle.

Let us consider  $g(z) = -2z$ . Then

$$|g(z) - f(z)| = |e^z - 1| = \left| \sum_{n=1}^{\infty} \frac{z^n}{n!} \right| \leq \sum_{n=1}^{\infty} \frac{|z^n|}{n!} = e - 1 < |g(z)|$$

on the unit circle. Hence by Rouché's theorem  $f(z)$  and  $g(z)$  have equal number of roots in the unit circle, namely 1.

Here's another quick and pretty proof of FTA using Rouché's theorem.

Let  $f(z) = a_0 + a_1z + \cdots + z^n$  be a non-constant polynomial. Take  $g(z) = z^n$ . Then on a sufficiently large circle around 0 of radius  $R$ ,  $|f(z) - g(z)| < |f(z)|$ . Hence  $f(z)$  and  $g(z)$  have same number of zero's in the disc of radius  $R$ . Since  $g(z)$  has  $n$  zero's, so does  $f(z)$  !



# Picard's theorem

I now state another absolutely spectacular theorem in complex analysis called Picard's theorem on the values taken by a holomorphic function.

# Picard's theorem

I now state another absolutely spectacular theorem in complex analysis called Picard's theorem on the values taken by a holomorphic function.

**Picard's Theorem:** Let  $z_0$  be an essential singularity of  $f(z)$ . Then in any punctured neighborhood of  $z_0$ , the image of  $f(z)$  can miss at most one point.  
This theorem is also called the Big Picard Theorem in view of what comes next.

# Picard's theorem

I now state another absolutely spectacular theorem in complex analysis called Picard's theorem on the values taken by a holomorphic function.

**Picard's Theorem:** Let  $z_0$  be an essential singularity of  $f(z)$ . Then in any punctured neighborhood of  $z_0$ , the image of  $f(z)$  can miss at most one point.  
This theorem is also called the Big Picard Theorem in view of what comes next.

A consequence of this theorem is the **Little Picard theorem** which states that any non-constant entire function can miss at most one point.

# Picard's theorem

I now state another absolutely spectacular theorem in complex analysis called Picard's theorem on the values taken by a holomorphic function.

**Picard's Theorem:** Let  $z_0$  be an essential singularity of  $f(z)$ . Then in any punctured neighborhood of  $z_0$ , the image of  $f(z)$  can miss at most one point.  
This theorem is also called the Big Picard Theorem in view of what comes next.

A consequence of this theorem is the **Little Picard theorem** which states that any non-constant entire function can miss at most one point.

The little Picard Theorem can be seen to be a corollary of the Big Picard Theorem as follows :

# Picard's Theorem

Recall the following fact mentioned earlier: An entire function has a pole at infinity if and only if it is a non-constant polynomial.

Let  $f(z)$  be a non-constant entire function. We wish to show it misses at most one point. If  $f(z)$  is a polynomial, then it is surjective by FTA. If  $f(z)$  is not a polynomial, then it has an essential singularity at infinity (WHY?). That is  $f(\frac{1}{z})$  has an essential singularity at 0. Thus by Big Picard theorem, in any punctured neighborhood of 0, say of radius  $r$ ,  $f(\frac{1}{z})$  misses at most one point. But this implies that in the complement of the circle of radius  $1/r$ ,  $f(z)$  misses at most one point. This is what we wanted.

In combination with an earlier theorem discussed in the course, we now have the following theorem: A non-constant entire function is either surjective or misses one point in which case it is of the form  $e^{g(z)} + c$  for some holomorphic function  $g(z)$  and some constant  $c$ .

# Liouville (1809-1882) & Picard (1856-1941); Wiki

**Joseph Liouville**



**Émile Picard**



# Liouville (1809-1882) & Picard (1856-1941); Wiki

**Joseph Liouville**



**Émile Picard**



Picard was a top rate mathematician who did fundamental work in many disciplines; analysis, function theory, differential equations, and analytic geometry to name a few. In physics he worked on elasticity, heat and electricity. Hadamard wrote about his teacher Picard:- A striking feature of Picard's scientific personality was the perfection of his teaching, one of the most marvellous, if not the most marvellous, that I have ever known.

It is a remarkable fact that between 1894 and 1937 he trained over 10000 engineers who were studying at the cole Centrale des Arts et Manufactures.