

MA 205 Complex Analysis: Examples of Contour Integration

August 24, 2017

Last time we began by understanding the notion of an isolated singularity at ∞ . Much the same way as isolated singularity at a point in \mathbb{C} , we can classify isolated singularities at infinity into removable, pole and essential singularity.

Last time we began by understanding the notion of an isolated singularity at ∞ . Much the same way as isolated singularity at a point in \mathbb{C} , we can classify isolated singularities at infinity into removable, pole and essential singularity.

We then looked at various examples of computing residues and contour integrals. Let us begin by looking at some more today.

Improper Integral

Let $f : [0, \infty] \rightarrow \mathbb{R}$ be a function such that $\int_0^R f(x)dx$ exists for each $R \geq 0$. One then defines the Improper integral $\int_0^\infty f(x)$ to be

$$\lim_{R \rightarrow \infty} \int_0^R f(x)dx.$$

Improper Integral

Let $f : [0, \infty] \rightarrow \mathbb{R}$ be a function such that $\int_0^R f(x)dx$ exists for each $R \geq 0$. One then defines the Improper integral $\int_0^\infty f(x)$ to be

$$\lim_{R \rightarrow \infty} \int_0^R f(x)dx.$$

Similarly if $f : [-\infty, \infty] \rightarrow \mathbb{R}$ is a function such that $\int_{-a}^b f(x)dx$ exists for each $a, b \geq 0$, then the improper integral $\int_{-\infty}^\infty f(x)$ is

defined as $\lim_{a, b \rightarrow \infty} \int_{-a}^b f(x)dx$.

Improper Integral

Let $f : [0, \infty] \rightarrow \mathbb{R}$ be a function such that $\int_0^R f(x)dx$ exists for each $R \geq 0$. One then defines the Improper integral $\int_0^\infty f(x)$ to be

$$\lim_{R \rightarrow \infty} \int_0^R f(x)dx.$$

Similarly if $f : [-\infty, \infty] \rightarrow \mathbb{R}$ is a function such that $\int_{-a}^b f(x)dx$ exists for each $a, b \geq 0$, then the improper integral $\int_{-\infty}^\infty f(x)$ is

defined as $\lim_{a, b \rightarrow \infty} \int_{-a}^b f(x)dx$. f is said to be integrable if the limit exists. If f is integrable, then its integral can be computed as

$$\lim_{R \rightarrow \infty} \int_{-R}^R f(x)dx.$$

Improper Integral

Let $f : [0, \infty] \rightarrow \mathbb{R}$ be a function such that $\int_0^R f(x)dx$ exists for each $R \geq 0$. One then defines the Improper integral $\int_0^\infty f(x)$ to be

$$\lim_{R \rightarrow \infty} \int_0^R f(x)dx.$$

Similarly if $f : [-\infty, \infty] \rightarrow \mathbb{R}$ is a function such that $\int_{-a}^b f(x)dx$ exists for each $a, b \geq 0$, then the improper integral $\int_{-\infty}^\infty f(x)$ is

defined as $\lim_{a, b \rightarrow \infty} \int_{-a}^b f(x)dx$. f is said to be integrable if the limit exists. If f is integrable, then its integral can be computed as

$$\lim_{R \rightarrow \infty} \int_{-R}^R f(x)dx.$$

For instance the function $\frac{1}{1+x^2}dx$ is integrable on \mathbb{R} while the integral $\int_{-\infty}^\infty \sin(x)dx$ does not exist. Intuitively, for such an improper integral to exist, the function has to decay to zero sufficiently rapidly outside “small set”.

Improper Integral

Recall the following bound we implicitly discussed in the last lecture:

Improper Integral

Recall the following bound we implicitly discussed in the last lecture:

If $P(z)/Q(z)$ is a rational function such that with $\deg Q(z) \geq \deg P(z) + 2$. Then there exists a constant C such that for $|P(z)/Q(z)| \leq C/|z|^2$ for $|z|$ sufficiently large. Thus for a large real number R , $|P(z)/Q(z)| \leq \frac{C}{R^2}$ on the circle of radius R .

Example

Compute $\int_{-\infty}^{\infty} \frac{dx}{(1+x^2)^n}$.

Example

Compute $\int_{-\infty}^{\infty} \frac{dx}{(1+x^2)^n}$.

Let $f(x) = \frac{1}{(1+x^2)^n}$. Proceed as in previous lecture by taking a contour γ to be the union of the segment from $-R$ to R along with the upper half semicircle of radius R , oriented positively.

Example

Compute $\int_{-\infty}^{\infty} \frac{dx}{(1+x^2)^n}$.

Let $f(x) = \frac{1}{(1+x^2)^n}$. Proceed as in previous lecture by taking a contour γ to be the union of the segment from $-R$ to R along with the upper half semicircle of radius R , oriented positively. There is just one pole inside γ which is i . Compute $\text{Res}(f; i)$, where $f(z) = \frac{1}{(1+z^2)^n}$.

Example

Compute $\int_{-\infty}^{\infty} \frac{dx}{(1+x^2)^n}$.

Let $f(x) = \frac{1}{(1+x^2)^n}$. Proceed as in previous lecture by taking a contour γ to be the union of the segment from $-R$ to R along with the upper half semicircle of radius R , oriented positively. There is just one pole inside γ which is i . Compute $\text{Res}(f; i)$, where $f(z) = \frac{1}{(1+z^2)^n}$. This is given by $\frac{g^{(n-1)}(i)}{(n-1)!}$, where $g(z) = \frac{1}{(z+i)^n}$. Check:

Example

Compute $\int_{-\infty}^{\infty} \frac{dx}{(1+x^2)^n}$.

Let $f(x) = \frac{1}{(1+x^2)^n}$. Proceed as in previous lecture by taking a contour γ to be the union of the segment from $-R$ to R along with the upper half semicircle of radius R , oriented positively. There is just one pole inside γ which is i . Compute $\text{Res}(f; i)$, where $f(z) = \frac{1}{(1+z^2)^n}$. This is given by $\frac{g^{(n-1)}(i)}{(n-1)!}$, where $g(z) = \frac{1}{(z+i)^n}$. Check:

$$\text{Res}(f; i) = \frac{-i}{2^{2n-1}} \binom{2n-2}{n-1}.$$

Example

Compute $\int_{-\infty}^{\infty} \frac{dx}{(1+x^2)^n}$.

Let $f(x) = \frac{1}{(1+x^2)^n}$. Proceed as in previous lecture by taking a contour γ to be the union of the segment from $-R$ to R along with the upper half semicircle of radius R , oriented positively. There is just one pole inside γ which is i . Compute $\text{Res}(f; i)$, where $f(z) = \frac{1}{(1+z^2)^n}$. This is given by $\frac{g^{(n-1)}(i)}{(n-1)!}$, where $g(z) = \frac{1}{(z+i)^n}$. Check:

$$\text{Res}(f; i) = \frac{-i}{2^{2n-1}} \binom{2n-2}{n-1}.$$

$\int_{\gamma} \frac{dz}{(1+z^2)^n} = \int_{-R}^R \frac{dx}{(1+x^2)^n} + \int_{C_R} \frac{dz}{(1+z^2)^n}$
where C_R is the upper circle of radius R .

if function is defined at 0 take semi circle

Example

Compute $\int_{-\infty}^{\infty} \frac{dx}{(1+x^2)^n}$.

Let $f(x) = \frac{1}{(1+x^2)^n}$. Proceed as in previous lecture by taking a contour γ to be the union of the segment from $-R$ to R along with the upper half semicircle of radius R , oriented positively. There is just one pole inside γ which is i . Compute $\text{Res}(f; i)$, where $f(z) = \frac{1}{(1+z^2)^n}$. This is given by $\frac{g^{(n-1)}(i)}{(n-1)!}$, where $g(z) = \frac{1}{(z+i)^n}$. Check:

$$\text{Res}(f; i) = \frac{-i}{2^{2n-1}} \binom{2n-2}{n-1}.$$

$\int_{\gamma} \frac{dz}{(1+z^2)^n} = \int_{-R}^R \frac{dx}{(1+x^2)^n} + \int_{C_R} \frac{dz}{(1+z^2)^n}$
where C_R is the upper circle of radius R .

By the earlier remark, there exists a constant C , such that $|\frac{1}{(1+z^2)^n}| \leq \frac{C}{R^2}$ on C_R .

Example

Compute $\int_{-\infty}^{\infty} \frac{dx}{(1+x^2)^n}$.

Let $f(x) = \frac{1}{(1+x^2)^n}$. Proceed as in previous lecture by taking a contour γ to be the union of the segment from $-R$ to R along with the upper half semicircle of radius R , oriented positively. There is just one pole inside γ which is i . Compute $\text{Res}(f; i)$, where $f(z) = \frac{1}{(1+z^2)^n}$. This is given by $\frac{g^{(n-1)}(i)}{(n-1)!}$, where $g(z) = \frac{1}{(z+i)^n}$. Check:

$$\text{Res}(f; i) = \frac{-i}{2^{2n-1}} \binom{2n-2}{n-1}.$$

$\int_{\gamma} \frac{dz}{(1+z^2)^n} = \int_{-R}^R \frac{dx}{(1+x^2)^n} + \int_{C_R} \frac{dz}{(1+z^2)^n}$
where C_R is the upper circle of radius R .

By the earlier remark, there exists a constant C , such that $|\frac{1}{(1+z^2)^n}| \leq \frac{C}{R^2}$ on C_R . Thus by ML inequality the second integral tends to zero as $R \rightarrow \infty$. Thus, the answer is $\frac{\pi}{4^{n-1}} \binom{2n-2}{n-1}$.

Jordan's lemma

Theorem (Jordan's Lemma)

Let f be a continuous function defined on the semicircular contour $C_R = \{Re^{i\theta} \mid \theta \in [0, \pi]\}$ of the form

$$f(z) = e^{iaz} g(z),$$

where $g(z)$ is a continuous function and with $a > 0$. Then,

$$\left| \int_{C_R} f(z) dz \right| \leq \frac{\pi}{a} \max_{\theta \in [0, \pi]} |g(Re^{i\theta})|.$$

Real Integrals

Proof:

$$\int_{C_R} f(z) dz = \int_0^\pi g(Re^{i\theta}) e^{iaR(\cos\theta + i\sin\theta)} iRe^{i\theta} d\theta.$$

Real Integrals

Proof:

$$\int_{C_R} f(z) dz = \int_0^\pi g(Re^{i\theta}) e^{iaR(\cos\theta + i\sin\theta)} iRe^{i\theta} d\theta.$$

Therefore,

$$\left| \int_{C_R} f(z) dz \right| \leq R \int_0^\pi \left| g(Re^{i\theta}) e^{aR(i\cos\theta - \sin\theta)} i e^{i\theta} \right| d\theta$$

Real Integrals

Proof:

$$\int_{C_R} f(z) dz = \int_0^\pi g(Re^{i\theta}) e^{iaR(\cos\theta + i\sin\theta)} iRe^{i\theta} d\theta.$$

Therefore,

$$\begin{aligned} \left| \int_{C_R} f(z) dz \right| &\leq R \int_0^\pi \left| g(Re^{i\theta}) e^{aR(i\cos\theta - \sin\theta)} i e^{i\theta} \right| d\theta \\ &= R \int_0^\pi \left| g(Re^{i\theta}) \right| e^{-aR\sin\theta} d\theta \end{aligned}$$

Real Integrals

Proof:

$$\int_{C_R} f(z) dz = \int_0^\pi g(Re^{i\theta}) e^{iaR(\cos\theta + i\sin\theta)} iRe^{i\theta} d\theta.$$

Therefore,

$$\begin{aligned} \left| \int_{C_R} f(z) dz \right| &\leq R \int_0^\pi \left| g(Re^{i\theta}) e^{aR(i\cos\theta - \sin\theta)} i e^{i\theta} \right| d\theta \\ &= R \int_0^\pi \left| g(Re^{i\theta}) \right| e^{-aR\sin\theta} d\theta \\ &\leq 2RM_R \int_0^{\frac{\pi}{2}} e^{-aR\sin\theta} d\theta \quad \text{where } M_R = \sup |g(Re^{i\theta})| \end{aligned}$$

Real Integrals

Proof:

$$\int_{C_R} f(z) dz = \int_0^\pi g(Re^{i\theta}) e^{iaR(\cos\theta + i\sin\theta)} iRe^{i\theta} d\theta.$$

Therefore,

$$\begin{aligned} \left| \int_{C_R} f(z) dz \right| &\leq R \int_0^\pi \left| g(Re^{i\theta}) e^{aR(i\cos\theta - \sin\theta)} i e^{i\theta} \right| d\theta \\ &= R \int_0^\pi \left| g(Re^{i\theta}) \right| e^{-aR\sin\theta} d\theta \\ &\leq 2RM_R \int_0^{\frac{\pi}{2}} e^{-aR\underline{\sin\theta}} d\theta \quad \text{where } M_R = \sup |g(Re^{i\theta})| \\ &\leq 2RM_R \int_0^{\frac{\pi}{2}} e^{\frac{-2aR\theta}{\pi}} d\theta = \frac{\pi}{a} (1 - e^{-aR}) M_R \leq \frac{\pi}{a} M_R, \end{aligned}$$

Real Integrals

Proof:

$$\int_{C_R} f(z) dz = \int_0^\pi g(Re^{i\theta}) e^{iaR(\cos\theta + i\sin\theta)} iRe^{i\theta} d\theta.$$

Therefore,

$$\begin{aligned} \left| \int_{C_R} f(z) dz \right| &\leq R \int_0^\pi \left| g(Re^{i\theta}) e^{aR(i\cos\theta - \sin\theta)} i e^{i\theta} \right| d\theta \\ &= R \int_0^\pi \left| g(Re^{i\theta}) \right| e^{-aR\sin\theta} d\theta \\ &\leq 2RM_R \int_0^{\frac{\pi}{2}} e^{-aR\sin\theta} d\theta \quad \text{where } M_R = \sup |g(Re^{i\theta})| \\ &\leq 2RM_R \int_0^{\frac{\pi}{2}} e^{\frac{-2aR\theta}{\pi}} d\theta = \frac{\pi}{a} (1 - e^{-aR}) M_R \leq \frac{\pi}{a} M_R, \end{aligned}$$

since $\sin\theta \geq \frac{2\theta}{\pi}$ for $\theta \in [0, \frac{\pi}{2}]$.

Example

Compute $\int_0^\infty \frac{\sin x}{x} dx$.

Example

Compute $\int_0^\infty \frac{\sin x}{x} dx$.

We'll consider the function

$$f(z) = \frac{e^{iz}}{z}.$$

Let γ be the boundary of the upper part of the annulus $A(0; r, R)$.
Then, $\int_\gamma f(z) dz = 0$, by Cauchy's theorem.

Example

Compute $\int_0^\infty \frac{\sin x}{x} dx$.

We'll consider the function

$$f(z) = \frac{e^{iz}}{z}.$$

Let γ be the boundary of the upper part of the annulus $A(0; r, R)$.
Then, $\int_\gamma f(z) dz = 0$, by Cauchy's theorem.

Example

But,

$$\int_{\gamma} f(z) dz = \int_r^R \frac{e^{ix}}{x} dx + \int_{\gamma_R} \frac{e^{iz}}{z} dz + \int_{-R}^{-r} \frac{e^{ix}}{x} dx + \int_{\gamma_r} \frac{e^{iz}}{z} dz.$$

Now,

$$\int_r^R \frac{\sin x}{x} dx = \frac{1}{2i} \int_r^R \frac{e^{ix} - e^{-ix}}{x} dx$$

Example

But,

$$\int_{\gamma} f(z) dz = \int_r^R \frac{e^{iz}}{x} dx + \int_{\gamma_R} \frac{e^{iz}}{z} dz + \int_{-R}^{-r} \frac{e^{iz}}{x} dx + \int_{\gamma_r} \frac{e^{iz}}{z} dz.$$

Now,

$$\begin{aligned} \int_r^R \frac{\sin x}{x} dx &= \frac{1}{2i} \int_r^R \frac{e^{ix} - e^{-ix}}{x} dx \\ &= \frac{1}{2i} \int_r^R \frac{e^{ix}}{x} dx + \frac{1}{2i} \int_{-R}^{-r} \frac{e^{ix}}{x} dx. \end{aligned}$$

Example

But,

$$\int_{\gamma} f(z) dz = \int_r^R \frac{e^{ix}}{x} dx + \int_{\gamma_R} \frac{e^{iz}}{z} dz + \int_{-R}^{-r} \frac{e^{ix}}{x} dx + \int_{\gamma_r} \frac{e^{iz}}{z} dz.$$

Now,

$$\begin{aligned} \int_r^R \frac{\sin x}{x} dx &= \frac{1}{2i} \int_r^R \frac{e^{ix} - e^{-ix}}{x} dx \\ &= \frac{1}{2i} \int_r^R \frac{e^{ix}}{x} dx + \frac{1}{2i} \int_{-R}^{-r} \frac{e^{ix}}{x} dx. \end{aligned}$$

Thus, we only need to compute

$$\lim_{R \rightarrow \infty} \int_{\gamma_R} \frac{e^{iz}}{z} dz \quad \& \quad \lim_{r \rightarrow 0} \int_{\gamma_r} \frac{e^{iz}}{z} dz.$$

Example

But,

$$\int_{\gamma} f(z) dz = \int_r^R \frac{e^{ix}}{x} dx + \int_{\gamma_R} \frac{e^{iz}}{z} dz + \int_{-R}^{-r} \frac{e^{ix}}{x} dx + \int_{\gamma_r} \frac{e^{iz}}{z} dz.$$

Now,

$$\begin{aligned} \int_r^R \frac{\sin x}{x} dx &= \frac{1}{2i} \int_r^R \frac{e^{ix} - e^{-ix}}{x} dx \\ &= \frac{1}{2i} \int_r^R \frac{e^{ix}}{x} dx + \frac{1}{2i} \int_{-R}^{-r} \frac{e^{ix}}{x} dx. \end{aligned}$$

Thus, we only need to compute

$$\lim_{R \rightarrow \infty} \int_{\gamma_R} \frac{e^{iz}}{z} dz \quad \& \quad \lim_{r \rightarrow 0} \int_{\gamma_r} \frac{e^{iz}}{z} dz.$$

Example

Now,

$$\lim_{R \rightarrow \infty} \int_{\gamma_R} \frac{e^{iz}}{z} dz = 0,$$

by Jordan's lemma.

Example

Now,

$$\lim_{R \rightarrow \infty} \int_{\gamma_R} \frac{e^{iz}}{z} dz = 0,$$

by Jordan's lemma. On the other hand, note that $\frac{e^{iz}-1}{z}$ has a removable singularity at 0.

Example

Now,

$$\lim_{R \rightarrow \infty} \int_{\gamma_R} \frac{e^{iz}}{z} dz = 0,$$

by Jordan's lemma. On the other hand, note that $\frac{e^{iz}-1}{z}$ has a removable singularity at 0. Thus, there is $M > 0$ such that

$$\left| \frac{e^{iz} - 1}{z} \right| \leq M,$$

for $|z| \leq 1$.

Example

Now,

$$\lim_{R \rightarrow \infty} \int_{\gamma_R} \frac{e^{iz}}{z} dz = 0,$$

by Jordan's lemma. On the other hand, note that $\frac{e^{iz}-1}{z}$ has a removable singularity at 0. Thus, there is $M > 0$ such that

$$\left| \frac{e^{iz} - 1}{z} \right| \leq M,$$

for $|z| \leq 1$. Thus,

$$\lim_{r \rightarrow 0} \int_{\gamma_r} \frac{e^{iz} - 1}{z} dz = 0,$$

by appealing to ML inequality.

Example

Therefore,

$$\lim_{r \rightarrow 0} \int_{\gamma_r} \frac{e^{iz}}{z} dz = -\pi i.$$

Example

Therefore,

$$\lim_{r \rightarrow 0} \int_{\gamma_r} \frac{e^{iz}}{z} dz = -\pi i.$$

Thus,

$$\int_0^\infty \frac{\sin x}{x} dx = \frac{\pi}{2}.$$

Example

Therefore,

$$\lim_{r \rightarrow 0} \int_{\gamma_r} \frac{e^{iz}}{z} dz = -\pi i.$$

Thus,

$$\int_0^\infty \frac{\sin x}{x} dx = \frac{\pi}{2}.$$

Example

Show that $\int_0^\infty \frac{\log x}{1+x^2} dx = 0$.

Example

Show that $\int_0^\infty \frac{\log x}{1+x^2} dx = 0$.

We'll work with γ as in the previous problem. We take

$$f(z) = \frac{\log z}{1+z^2},$$

Example

Show that $\int_0^\infty \frac{\log x}{1+x^2} dx = 0$.

We'll work with γ as in the previous problem. We take

$$f(z) = \frac{\log z}{1+z^2},$$

where $\log z$ is a branch of the logarithm which is defined on the x -axis, so that \int_r^R and \int_{-R}^{-r} make sense.

Example

Show that $\int_0^\infty \frac{\log x}{1+x^2} dx = 0$.

We'll work with γ as in the previous problem. We take

$$f(z) = \frac{\log z}{1+z^2},$$

where $\log z$ is a branch of the logarithm which is defined on the x -axis, so that \int_r^R and \int_{-R}^{-r} make sense. For instance, we can take the branch with negative y -axis as the branch cut. Then,

Example

Show that $\int_0^\infty \frac{\log x}{1+x^2} dx = 0$.

We'll work with γ as in the previous problem. We take

$$f(z) = \frac{\log z}{1+z^2},$$

where $\log z$ is a branch of the logarithm which is defined on the x -axis, so that \int_r^R and \int_{-R}^{-r} make sense. For instance, we can take the branch with negative y -axis as the branch cut. Then,

$$\log x = \begin{cases} \log x & \text{if } x > 0, \\ \log |x| + i\pi & \text{if } x < 0. \end{cases}$$

Example

Now,

$$\int_{\gamma} \frac{\log z}{1+z^2} dz = \int_r^R \frac{\log x}{1+x^2} dx + \int_{\gamma_R} \frac{\log z}{1+z^2} dz$$

Example

Now,

$$\begin{aligned}\int_{\gamma} \frac{\log z}{1+z^2} dz &= \int_r^R \frac{\log x}{1+x^2} dx + \int_{\gamma_R} \frac{\log z}{1+z^2} dz \\ &+ \int_{-R}^{-r} \frac{\log |x| + i\pi}{1+x^2} dx + \int_{\gamma_r} \frac{\log z}{1+z^2} dz.\end{aligned}$$

Example

Now,

$$\begin{aligned}\int_{\gamma} \frac{\log z}{1+z^2} dz &= \int_r^R \frac{\log x}{1+x^2} dx + \int_{\gamma_R} \frac{\log z}{1+z^2} dz \\ &\quad + \int_{-R}^{-r} \frac{\log |x| + i\pi}{1+x^2} dx + \int_{\gamma_r} \frac{\log z}{1+z^2} dz.\end{aligned}$$

Lhs is $2\pi i \cdot \text{Res}(f; i)$ which is $2\pi i \cdot \frac{\log i}{2i} = \frac{\pi^2 i}{2}$. Also,

$$\int_r^R \frac{\log x}{1+x^2} dx + \int_{-R}^{-r} \frac{\log |x| + i\pi}{1+x^2} dx = 2 \int_r^R \frac{\log x}{1+x^2} dx + i\pi \int_r^R \frac{dx}{1+x^2}$$

Example

Now,

$$\begin{aligned}\int_{\gamma} \frac{\log z}{1+z^2} dz &= \int_r^R \frac{\log x}{1+x^2} dx + \int_{\gamma_R} \frac{\log z}{1+z^2} dz \\ &\quad + \int_{-R}^{-r} \frac{\log |x| + i\pi}{1+x^2} dx + \int_{\gamma_r} \frac{\log z}{1+z^2} dz.\end{aligned}$$

Lhs is $2\pi i \cdot \text{Res}(f; i)$ which is $2\pi i \cdot \frac{\log i}{2i} = \frac{\pi^2 i}{2}$. Also,

$$\begin{aligned}\int_r^R \frac{\log x}{1+x^2} dx + \int_{-R}^{-r} \frac{\log |x| + i\pi}{1+x^2} dx &= 2 \int_r^R \frac{\log x}{1+x^2} dx + i\pi \int_r^R \frac{dx}{1+x^2} \\ &= 2 \int_r^R \frac{\log x}{1+x^2} dx + \frac{\pi^2 i}{2}.\end{aligned}$$

(In the Limit)

Thus,

$$\int_r^R \frac{\log x}{1+x^2} dx = -\frac{1}{2} \left[\int_{\gamma_R} \frac{\log z}{1+z^2} dz + \int_{\gamma_r} \frac{\log z}{1+z^2} dz \right].$$

Thus,

$$\int_r^R \frac{\log x}{1+x^2} dx = -\frac{1}{2} \left[\int_{\gamma_R} \frac{\log z}{1+z^2} dz + \int_{\gamma_r} \frac{\log z}{1+z^2} dz \right].$$

Note that

$$\left| \int_{\gamma_\rho} \frac{\log z}{1+z^2} dz \right| = \left| \rho \int_0^\pi \frac{\log \rho + i\theta}{1+\rho^2 e^{i\theta}} e^{i\theta} d\theta \right|$$

Thus,

$$\int_r^R \frac{\log x}{1+x^2} dx = -\frac{1}{2} \left[\int_{\gamma_R} \frac{\log z}{1+z^2} dz + \int_{\gamma_r} \frac{\log z}{1+z^2} dz \right].$$

Note that

$$\begin{aligned} \left| \int_{\gamma_\rho} \frac{\log z}{1+z^2} dz \right| &= \left| \rho \int_0^\pi \frac{\log \rho + i\theta}{1+\rho^2 e^{i\theta}} e^{i\theta} d\theta \right| \\ &\leq \frac{\rho |\log \rho|}{|1-\rho^2|} \int_0^\pi d\theta + \frac{\rho}{|1-\rho^2|} \int_0^\pi \theta d\theta \end{aligned}$$

Thus,

$$\int_r^R \frac{\log x}{1+x^2} dx = -\frac{1}{2} \left[\int_{\gamma_R} \frac{\log z}{1+z^2} dz + \int_{\gamma_r} \frac{\log z}{1+z^2} dz \right].$$

Note that

$$\begin{aligned} \left| \int_{\gamma_\rho} \frac{\log z}{1+z^2} dz \right| &= \left| \rho \int_0^\pi \frac{\log \rho + i\theta}{1+\rho^2 e^{i\theta}} e^{i\theta} d\theta \right| \\ &\leq \frac{\rho |\log \rho|}{|1-\rho^2|} \int_0^\pi d\theta + \frac{\rho}{|1-\rho^2|} \int_0^\pi \theta d\theta \\ &= \frac{\pi \rho |\log \rho|}{|1-\rho^2|} + \frac{\rho \pi^2}{2|1-\rho^2|}. \end{aligned}$$

Thus,

$$\int_r^R \frac{\log x}{1+x^2} dx = -\frac{1}{2} \left[\int_{\gamma_R} \frac{\log z}{1+z^2} dz + \int_{\gamma_r} \frac{\log z}{1+z^2} dz \right].$$

Note that

$$\begin{aligned} \left| \int_{\gamma_\rho} \frac{\log z}{1+z^2} dz \right| &= \left| \rho \int_0^\pi \frac{\log \rho + i\theta}{1+\rho^2 e^{i\theta}} e^{i\theta} d\theta \right| \\ &\leq \frac{\rho |\log \rho|}{|1-\rho^2|} \int_0^\pi d\theta + \frac{\rho}{|1-\rho^2|} \int_0^\pi \theta d\theta \\ &= \frac{\pi \rho |\log \rho|}{|1-\rho^2|} + \frac{\rho \pi^2}{2|1-\rho^2|}. \end{aligned}$$

This is zero in the limit if $\rho \rightarrow 0+$ or $\rho \rightarrow \infty$. Thus, the given integral is zero.

Maximum Modulus Theorem

An important theorem in Complex Analysis states that a non-constant holomorphic function on an open connected domain never attains its maximum modulus at any point in the domain. This is called the maximum modulus theorem. Once again, this is vastly different from what happens to real differentiable functions; in fact even for real analytic functions. Real analytic functions can achieve maximum anywhere inside the interval. We'll use CIF and the identity theorem to prove MMT.

Maximum Modulus Theorem

Proof: Suppose there is $z_0 \in \Omega$ such that $|f(z_0)| \geq |f(z)|$ for all $z \in \Omega$. Then we'll prove that f is a constant.

Maximum Modulus Theorem

Proof: Suppose there is $z_0 \in \Omega$ such that $|f(z_0)| \geq |f(z)|$ for all $z \in \Omega$. Then we'll prove that f is a constant. Let γ be a small circle around z_0 with radius r such that the closed disc with boundary γ is contained in Ω . CIF gives,

$$f(z_0) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{z - z_0} dz = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{i\theta}) d\theta.$$

Maximum Modulus Theorem

Proof: Suppose there is $z_0 \in \Omega$ such that $|f(z_0)| \geq |f(z)|$ for all $z \in \Omega$. Then we'll prove that f is a constant. Let γ be a small circle around z_0 with radius r such that the closed disc with boundary γ is contained in Ω . CIF gives,

$$f(z_0) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{z - z_0} dz = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{i\theta}) d\theta.$$

Hence,

$$|f(z_0)| \leq \frac{1}{2\pi} \int_0^{2\pi} |f(z_0 + re^{i\theta})| d\theta \leq |f(z_0)|,$$

since $|f(z_0)|$ is assumed to be the maximum value.

Maximum Modulus Theorem

Thus,

$$\int_0^{2\pi} \left[|f(z_0)| - |f(z_0 + re^{i\theta})| \right] dt = 0.$$

Note that the integrand is non-negative. Therefore it has to be zero; i.e., $|f(z_0)| = |f(z_0 + re^{i\theta})|$ for all θ . Since this is true for each small r , we see that $|f(z)|$ is a constant on a small disc around z_0 . This means that $f(z)$ is a constant, say c , on this small disc. (Why?) This implies that $f \equiv c$ on Ω by the identity theorem, since a disc has limit points!

A nice consequence of the Maximum modulus principle is the following lemma of Schwartz.

Schwarz Lemma : Let $\mathbb{D} = \{z : |z| < 1\}$ be the open unit disk and let $f : \mathbb{D} \rightarrow \mathbb{C}$ be a holomorphic map such that $f(0) = 0$ and $|f(z)| \leq 1$ on \mathbb{D} .

Then, $|f(z)| \leq |z| \forall z \in \mathbb{D}$ and $|f'(0)| \leq 1$.

Moreover, if $|f(z)| = |z|$ for some non-zero z or $|f'(0)| = 1$, then $f(z) = az$ for some $a \in \mathbb{C}$ with $|a| = 1$.