

MA 205 Complex Analysis: Gamma Function and Harmonic Functions

September 5, 2017

Now, let's introduce ourselves to a very important function; the Gamma function. It's defined as:

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt,$$

for $\operatorname{Re}(z) > 0$.

Now, let's introduce ourselves to a very important function; the Gamma function. It's defined as:

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt,$$

for $\operatorname{Re}(z) > 0$. One checks that this integral exists and defines a holomorphic function in the right half plane. $\Gamma(1) = 1$.

Integration by parts in the right half plane gives:

$$\Gamma(z+1) = z\Gamma(z).$$

Integration by parts in the right half plane gives:

$$\Gamma(z+1) = z\Gamma(z).$$

Thus,

$$\Gamma(n+1) = n\Gamma(n) = n(n-1)\Gamma(n-1) \dots \Gamma(1) = n!.$$

The Gamma function interpolates the factorial function!

Real Integrals

We would like to define the Gamma function throughout \mathbb{C} . Right now, it's defined only on the right half plane.

We would like to define the Gamma function throughout \mathbb{C} . Right now, it's defined only on the right half plane. For this, use the identity

$$\Gamma(z+1) = z\Gamma(z).$$

We would like to define the Gamma function throughout \mathbb{C} . Right now, it's defined only on the right half plane. For this, use the identity

$$\Gamma(z+1) = z\Gamma(z).$$

Plug in $z = 0$, and we get:

$$0 \cdot \Gamma(0) = 1.$$

We would like to define the Gamma function throughout \mathbb{C} . Right now, it's defined only on the right half plane. For this, use the identity

$$\Gamma(z+1) = z\Gamma(z).$$

Plug in $z = 0$, and we get:

$$0 \cdot \Gamma(0) = 1.$$

Thus 0 must be a pole for the extended Gamma function.

We would like to define the Gamma function throughout \mathbb{C} . Right now, it's defined only on the right half plane. For this, use the identity

$$\Gamma(z+1) = z\Gamma(z).$$

Plug in $z = 0$, and we get:

$$0 \cdot \Gamma(0) = 1.$$

Thus 0 must be a pole for the extended Gamma function. Similarly, all negative integers are also poles. And these are the only poles.

Real Integrals

We would like to define the Gamma function throughout \mathbb{C} . Right now, it's defined only on the right half plane. For this, use the identity

$$\Gamma(z+1) = z\Gamma(z).$$

Plug in $z = 0$, and we get:

$$0 \cdot \Gamma(0) = 1.$$

negative



Thus 0 must be a pole for the extended Gamma function. Similarly, all negative integers are also poles. And these are the only poles.

Exercise: Check that these poles are simple and

$$\operatorname{Res}(\Gamma; -n) = \frac{(-1)^n}{n!}.$$

Real Integrals

What is $\Gamma(x)\Gamma(y)$?

$$\Gamma(x)\Gamma(y) = \int_0^\infty \int_0^\infty e^{-u-v} u^{x-1} v^{y-1} du dv.$$

Put

$$u = zt; \quad v = z(1 - t),$$

and apply the change of variables formula from MA 105.

Real Integrals

What is $\Gamma(x)\Gamma(y)$?

$$\Gamma(x)\Gamma(y) = \int_0^\infty \int_0^\infty e^{-u-v} u^{x-1} v^{y-1} du dv.$$

Put

$$u = zt; \quad v = z(1 - t),$$

and apply the change of variables formula from MA 105. Check:

$$\Gamma(x)\Gamma(y) = \Gamma(x + y)B(x, y),$$

where

$$B(x, y) = \int_0^1 t^{x-1} (1 - t)^{y-1} dt.$$

Real Integrals

What is $\Gamma(x)\Gamma(y)$?

$$\Gamma(x)\Gamma(y) = \int_0^\infty \int_0^\infty e^{-u-v} u^{x-1} v^{y-1} du dv.$$

Put

$$u = zt; \quad v = z(1-t),$$

and apply the change of variables formula from MA 105. Check:

$$\Gamma(x)\Gamma(y) = \Gamma(x+y)B(x,y), \quad \text{ye ho kya raha hai}$$

where

$$B(x,y) = \int_0^1 t^{x-1}(1-t)^{y-1} dt.$$

Put $t = \frac{s}{s+1}$ to get:

$$B(1-c, c) = \int_0^\infty \frac{s^{-c}}{1+s} ds, \quad \text{← } x+y=1$$

for $0 < c < 1$.

Thus, for $0 < x < 1$,

$$\Gamma(x)\Gamma(1-x) = \int_0^\infty \frac{t^{-x}}{1+t} dt = \frac{\pi}{\sin \pi x}.$$

Thus, for $0 < x < 1$,

$$\Gamma(x)\Gamma(1-x) = \int_0^\infty \frac{t^{-x}}{1+t} dt = \frac{\pi}{\sin \pi x}.$$

In particular,

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}.$$

By identity theorem,

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin \pi z},$$

for all $z \in \mathbb{C}$

Real Integrals

Thus, for $0 < x < 1$,

$$\Gamma(x)\Gamma(1-x) = \int_0^\infty \frac{t^{-x}}{1+t} dt = \frac{\pi}{\sin \pi x}.$$

In particular,

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}.$$

By identity theorem,

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin \pi z},$$

for all $z \in \mathbb{C}$

domain minus pts of pole

Introduce meromorphic functions. Why is the identity theorem valid for meromorphic functions?



“A mathematician is one to whom $\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}$ is as obvious as that twice two makes four is to you. ” - Lord Kelvin

Harmonic Functions

We now revisit a topic we studied at the beginning of the course, namely harmonic functions. Recall that a function $u(x, y)$ of real variables is said to be harmonic if it is twice differentiable and $u_{xx} + u_{yy} = 0$. It turns out that harmonic functions share many properties similar to holomorphic functions. We'll see some of them.

Harmonic Functions

We now revisit a topic we studied at the beginning of the course, namely harmonic functions. Recall that a function $u(x, y)$ of real variables is said to be harmonic if it is twice differentiable and $u_{xx} + u_{yy} = 0$. It turns out that harmonic functions share many properties similar to holomorphic functions. We'll see some of them.

Recall that if u is a harmonic function, then a harmonic conjugate of u is another harmonic function v such that $u + iv$ is holomorphic. We saw some examples of computing harmonic conjugates and that time I commented that if the domain is “nice”, then a harmonic conjugate always exists. The mathematical notion that replaces “nice” is simply connectedness.

Harmonic Functions

We now revisit a topic we studied at the beginning of the course, namely harmonic functions. Recall that a function $u(x, y)$ of real variables is said to be harmonic if it is twice differentiable and $u_{xx} + u_{yy} = 0$. It turns out that harmonic functions share many properties similar to holomorphic functions. We'll see some of them.

Recall that if u is a harmonic function, then a harmonic conjugate of u is another harmonic function v such that $u + iv$ is holomorphic. We saw some examples of computing harmonic conjugates and that time I commented that if the domain is “nice”, then a harmonic conjugate always exists. The mathematical notion that replaces “nice” is simply connectedness.

Theorem: Let U be a simply-connected domain in \mathbb{C} and let u be a harmonic function on U . Then u admits exactly one harmonic conjugate up to a constant.

Harmonic Functions

Proof: Let's dismiss the uniqueness first. Suppose u has a harmonic conjugate v . Let $f(z) = u + iv$. By CR equations, v_x and v_y are determined and hence v is determined upto a constant. To prove existence, let $g(z) = u_x - iu_y$. Then by CR equations, $g(z)$ is holomorphic. Now fix $z_0 \in U$, and define f to be the anti-derivative of g :

Harmonic Functions

Proof: Let's dismiss the uniqueness first. Suppose u has a harmonic conjugate v . Let $f(z) = u + iv$. By CR equations, v_x and v_y are determined and hence v is determined upto a constant. To prove existence, let $g(z) = u_x - iu_y$. Then by CR equations, $g(z)$ is holomorphic. Now fix $z_0 \in U$, and define f to be the anti-derivative of g :

$f(z) = u(z_0) + \int_{z_0}^z g(z)dz$ with the integral being along a path in U connecting z_0 and z . As U is simply-connected, this function is well-defined. By construction, f is holomorphic and $f' = g = u_x - iu_y$.

Harmonic Functions

Proof: Let's dismiss the uniqueness first. Suppose u has a harmonic conjugate v . Let $f(z) = u + iv$. By CR equations, v_x and v_y are determined and hence v is determined upto a constant. To prove existence, let $g(z) = u_x - iu_y$. Then by CR equations, $g(z)$ is holomorphic. Now fix $z_0 \in U$, and define f to be the anti-derivative of g :

$f(z) = u(z_0) + \int_{z_0}^z g(z)dz$ with the integral being along a path in U connecting z_0 and z . As U is simply-connected, this function is well-defined. By construction, f is holomorphic and $f' = g = u_x - iu_y$. Letting $\tilde{u} = \operatorname{Re}(f)$, by the Cauchy-Riemann for f , $f' = \tilde{u}_x - i\tilde{u}_y$. On comparing the two equations, one concludes that $\tilde{u}_x = u_x$ and $\tilde{u}_y = u_y$.

Harmonic Functions

Proof: Let's dismiss the uniqueness first. Suppose u has a harmonic conjugate v . Let $f(z) = u + iv$. By CR equations, v_x and v_y are determined and hence v is determined upto a constant. To prove existence, let $g(z) = u_x - iu_y$. Then by CR equations, $g(z)$ is holomorphic. Now fix $z_0 \in U$, and define f to be the anti-derivative of g :

$f(z) = u(z_0) + \int_{z_0}^z g(z)dz$ with the integral being along a path in U connecting z_0 and z . As U is simply-connected, this function is well-defined. By construction, f is holomorphic and $f' = g = u_x - iu_y$. Letting $\tilde{u} = \operatorname{Re}(f)$, by the Cauchy-Riemann for f , $f' = \tilde{u}_x - i\tilde{u}_y$. On comparing the two equations, one concludes that $\tilde{u}_x = u_x$ and $\tilde{u}_y = u_y$. Therefore $u - \tilde{u}$ is a constant. Since u and \tilde{u} are equal at z_0 , they are equal throughout.

Mean Value Property

Corollary: Harmonic functions are infinitely differentiable. (Why ?)

Mean Value Property

Corollary: Harmonic functions are infinitely differentiable. (Why ?)

Let us now derive an expression analogous to the CIF.

Mean Value Property

Corollary: Harmonic functions are infinitely differentiable. (Why ?)

Let us now derive an expression analogous to the CIF.

Theorem: (Mean-Value Property): Let u be a harmonic function on a disc of radius R . Then for any $r < R$, we have,

$$u(w) = \frac{1}{2\pi} \int_0^{2\pi} u(w + re^{i\theta}) d\theta.$$

In particular, u does not attain its maximum at any interior point unless it is constant.

Mean Value Property

Corollary: Harmonic functions are infinitely differentiable. (Why ?)

Let us now derive an expression analogous to the CIF.

Theorem: (Mean-Value Property): Let u be a harmonic function on a disc of radius R . Then for any $r < R$, we have,

$$u(w) = \frac{1}{2\pi} \int_0^{2\pi} u(w + re^{i\theta}) d\theta.$$

In particular, u does not attain its maximum at any interior point unless it is constant.

Proof: Can assume $w = 0$ without loss of generality. Since u is harmonic and the domain is simply-connected, there exists a holomorphic function $f(z)$ such that $\operatorname{Re}(f) = u$. By Cauchy's integral formula,

$$f(0) = \frac{1}{2\pi i} \int_{|z|=r} \frac{f(z)}{z} dz$$

Identity Principle

Hence

$$f(0) = \frac{1}{2\pi} \int_0^{2\pi} f(re^{i\theta}) d\theta$$

Taking real parts on both sides, we get the desired result.

Identity Principle

Hence

$$f(0) = \frac{1}{2\pi} \int_0^{2\pi} f(re^{i\theta}) d\theta$$

Taking real parts on both sides, we get the desired result.

Note that this in particular implies that if u is a harmonic function on a domain Ω and $z \in \Omega$. Suppose $u(z) \geq u(w)$ for all w in a neighborhood of z , then u is a constant. By considering $-u$ instead of u , the same holds for \geq replaced by \leq .

Identity Principle

Hence

$$f(0) = \frac{1}{2\pi} \int_0^{2\pi} f(re^{i\theta}) d\theta$$

Taking real parts on both sides, we get the desired result.

Note that this in particular implies that if u is a harmonic function on a domain Ω and $z \in \Omega$. Suppose $u(z) \geq u(w)$ for all w in a neighborhood of z , then u is a constant. By considering $-u$ instead of u , the same holds for \geq replaced by \leq .

Identity Principle: Let u be a harmonic function on a domain $\Omega \in \mathbb{C}$. If $u = 0$ on a non-empty open subset $U \subseteq \Omega$, then $u = 0$ throughout Ω .

Identity Principle

Hence

$$f(0) = \frac{1}{2\pi} \int_0^{2\pi} f(re^{i\theta}) d\theta$$

Taking real parts on both sides, we get the desired result.

Note that this in particular implies that if u is a harmonic function on a domain Ω and $z \in \Omega$. Suppose $u(z) \geq u(w)$ for all w in a neighborhood of z , then u is a constant. By considering $-u$ instead of u , the same holds for \geq replaced by \leq .

Identity Principle: Let u be a harmonic function on a domain $\Omega \in \mathbb{C}$. If $u = 0$ on a non-empty open subset $U \subseteq \Omega$, then $u = 0$ throughout Ω .

Proof: Set $f = u_x - iu_y$. Then as before, f is holomorphic on Ω . Since $u = 0$ on U then so is f . Hence, by the Identity Principle for the holomorphic functions $f = 0$ on Ω , and consequently, $u_x = x_y = 0$ on Ω . Therefore u is constant on Ω , and as it is zero on U , it must be zero on Ω .

Identity Theorem

Remark: Recall that the identity theorem for holomorphic functions is stronger; namely if a holomorphic function vanishes on a set of points having a limit point, then it is identically zero. **This is not true for harmonic functions.** The function $\operatorname{Re}(z)$ vanishes identically on imaginary axis but is non-zero elsewhere. However the following stronger identity theorem holds:

Remark: Recall that the identity theorem for holomorphic functions is stronger; namely if a holomorphic function vanishes on a set of points having a limit point, then it is identically zero. **This is not true for harmonic functions.** The function $Re(z)$ vanishes identically on imaginary axis but is non-zero elsewhere. However the following stronger identity theorem holds:

Theorem: Let Ω be a domain in \mathbb{C} and u_1 and u_2 are two harmonic functions that extend continuously to the boundary $\partial\Omega$ of Ω . If $u_1 = u_2$ on $\partial\Omega$ then these two functions are equal throughout Ω .

Dirichlet Problem

A very important problem in Mathematics and Mathematical Physics is the Dirichlet Problem. It asks if Ω is a domain with boundary $\partial\Omega$, and f is a continuous real function on the boundary, does there exist a function u on $\bar{\Omega}$ which is harmonic on Ω and equals f on the boundary. The problem has a positive answer if $\partial\Omega$ is “sufficiently smooth”. (the precise condition is more technical). Many mathematicians have contributed to the solution - Green, Gauss, Kelvin, Dirichlet (who solved it for the ball), Riemann, Poincare, Hilbert

Dirichlet Problem

A very important problem in Mathematics and Mathematical Physics is the Dirichlet Problem. It asks if Ω is a domain with boundary $\partial\Omega$, and f is a continuous real function on the boundary, does there exist a function u on $\bar{\Omega}$ which is harmonic on Ω and equals f on the boundary. The problem has a positive answer if $\partial\Omega$ is “sufficiently smooth”. (the precise condition is more technical). Many mathematicians have contributed to the solution - Green, Gauss, Kelvin, Dirichlet (who solved it for the ball), Riemann, Poincare, Hilbert Riemann gave a physical argument roughly as follows: any charge distribution on the boundary should, by the laws of electrostatics, determine an electrical potential as solution. **I confess I don't understand this !** In case of the disc in \mathbb{C} , the solution can be simply given as :

Dirichlet Problem

A very important problem in Mathematics and Mathematical Physics is the Dirichlet Problem. It asks if Ω is a domain with boundary $\partial\Omega$, and f is a continuous real function on the boundary, does there exist a function u on $\bar{\Omega}$ which is harmonic on Ω and equals f on the boundary. The problem has a positive answer if $\partial\Omega$ is “sufficiently smooth”. (the precise condition is more technical). Many mathematicians have contributed to the solution - Green, Gauss, Kelvin, Dirichlet (who solved it for the ball), Riemann, Poincaré, Hilbert Riemann gave a physical argument roughly as follows: any charge distribution on the boundary should, by the laws of electrostatics, determine an electrical potential as solution. I confess I don't understand this ! In case of the disc in \mathbb{C} , the solution can be simply given as :

$$u(z) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\psi}) \frac{(1 - |z|^2)}{|z - e^{i\psi}|^2} \, d\psi \text{ on } \mathbb{D}$$

and equal to $f(z)$ on $\partial\mathbb{D}$.

Exercise: Show that the function $\log(\sqrt{X^2 + Y^2})$ on \mathbb{C}^* admits no harmonic conjugate.