

# MA 205 Complex Analysis: Laurent Series and Examples

August 19, 2017

Last time we looked at some examples of contour integration. We then began discussing singularities. There are two of singularities: isolated and non-isolated singularities.

Last time we looked at some examples of contour integration. We then began discussing singularities. There are two of singularities: isolated and non-isolated singularities. Consider  $\tan(1/z)$ . Is the singularity at 0 isolated or non-isolated ?

Last time we looked at some examples of contour integration. We then began discussing singularities. There are two of singularities: isolated and non-isolated singularities. Consider  $\tan(1/z)$ . Is the singularity at 0 isolated or non-isolated ?

Isolated singularities are of 3 types: Removable singularity, Pole and Essential Singularity.

Last time we looked at some examples of contour integration. We then began discussing singularities. There are two of singularities: isolated and non-isolated singularities. Consider  $\tan(1/z)$ . Is the singularity at 0 isolated or non-isolated ?

Isolated singularities are of 3 types: Removable singularity, Pole and Essential Singularity.

A singularity at  $z_0$  is **removable** if  $\lim_{z \rightarrow z_0} f(z)$  exists. In particular  $f(z)$  is bounded in a neighborhood of  $z_0$ .

A singularity at  $z_0$  is a **pole** if  $f(z) \rightarrow \infty$  as  $z \rightarrow z_0$ . In particular the function takes unbounded values in any punctured neighborhood of  $z_0$ .

A singularity at  $z_0$  is an **essential singularity** if it is neither a removable singularity nor a pole.

# Casorati-Weierstrass Theorem

We also discussed the Casorati Weierstrass theorem which said that restricted to any punctured neighborhood of an essential singularity, the image of  $f(z)$  is dense, i.e, comes arbitrarily close to every complex number.

# Casorati-Weierstrass Theorem

We also discussed the Casorati Weierstrass theorem which said that restricted to any punctured neighborhood of an essential singularity, the image of  $f(z)$  is dense, i.e, comes arbitrarily close to every complex number.

We will see little later that this is a special case of a much more general and very beautiful theorem. I will keep the suspense for now !

# Casorati-Weierstrass Theorem

We also discussed the Casorati Weierstrass theorem which said that restricted to any punctured neighborhood of an essential singularity, the image of  $f(z)$  is dense, i.e, comes arbitrarily close to every complex number.

We will see little later that this is a special case of a much more general and very beautiful theorem. I will keep the suspense for now !

We also discussed an example :  $e^{1/z}$  has an essential singularity at 0.

Let us discuss some more examples (on the board).



# Laurent Series

Recall how we derived the power series representation of a holomorphic function on a disc centered around  $z_0$ . We used

$$f(z) = \frac{1}{2\pi i} \int_{|w-z_0|=r} \frac{f(w)}{w-z} dw,$$

and manipulated  $\frac{1}{w-z}$  as

$$\frac{1}{w-z_0} \cdot \frac{1}{1 - \frac{z-z_0}{w-z_0}}.$$

# Laurent Series

Now suppose  $z_0$  is an isolated singularity for  $f$ . Consider an annulus with radii  $R > r$  centered at  $z_0$  such that  $f$  is holomorphic there. CIF takes the form:

Now suppose  $z_0$  is an isolated singularity for  $f$ . Consider an annulus with radii  $R > r$  centered at  $z_0$  such that  $f$  is holomorphic there. CIF takes the form:

$$f(z) = \frac{1}{2\pi i} \int_{|w-z_0|=R} \frac{f(w)}{w-z} dw - \frac{1}{2\pi i} \int_{|w-z_0|=r} \frac{f(w)}{w-z} dw.$$

# Laurent Series

Now suppose  $z_0$  is an isolated singularity for  $f$ . Consider an annulus with radii  $R > r$  centered at  $z_0$  such that  $f$  is holomorphic there. CIF takes the form:

$$f(z) = \frac{1}{2\pi i} \int_{|w-z_0|=R} \frac{f(w)}{w-z} dw - \frac{1}{2\pi i} \int_{|w-z_0|=r} \frac{f(w)}{w-z} dw.$$

The first integral gives rise to  $\sum_{n=1}^{\infty} a_n (z - z_0)^n$  with

$$a_n = \frac{1}{2\pi i} \int_{|z-z_0|=R} \frac{f(z)}{(z-z_0)^{n+1}} dz,$$

exactly as before.

# Laurent Series

In the second integral, write

$$\frac{-1}{w - z} = \frac{1}{z - z_0} \cdot \frac{1}{1 - \frac{w - z_0}{z - z_0}},$$

Note that  $\left| \frac{w - z_0}{z - z_0} \right| < 1$  for all  $w$  with  $|w - z_0| = r$ .

Expand to get  $\sum_{n=1}^{\infty} \frac{b_n}{(z - z_0)^n}$  with :

# Laurent Series

In the second integral, write

$$\frac{-1}{w - z} = \frac{1}{z - z_0} \cdot \frac{1}{1 - \frac{w - z_0}{z - z_0}},$$

Note that  $\left| \frac{w - z_0}{z - z_0} \right| < 1$  for all  $w$  with  $|w - z_0| = r$ .

Expand to get  $\sum_{n=1}^{\infty} \frac{b_n}{(z - z_0)^n}$  with :

$$b_n = \frac{1}{2\pi i} \int_{|z - z_0| = r} \frac{f(z)}{(z - z_0)^{-n+1}} dz.$$

# Laurent Series

In the second integral, write

$$\frac{-1}{w - z} = \frac{1}{z - z_0} \cdot \frac{1}{1 - \frac{w - z_0}{z - z_0}},$$

Note that  $|\frac{w - z_0}{z - z_0}| < 1$  for all  $w$  with  $|w - z_0| = r$ .

Expand to get  $\sum_{n=1}^{\infty} \frac{b_n}{(z - z_0)^n}$  with :

$$b_n = \frac{1}{2\pi i} \int_{|z - z_0|=r} \frac{f(z)}{(z - z_0)^{-n+1}} dz.$$

We write both together as  $\sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$ . This is the Laurent series around the isolated singularity  $z_0$ . The negative part is called the principal part of the Laurent series.

If  $z_0$  is an isolated singularity of  $f$ , then  $f$  is holomorphic in an annulus  $0 < |z - z_0| < R$  for some  $R$ . The corresponding Laurent expansion is called the Laurent expansion around  $z_0$ .



If  $z_0$  is an isolated singularity of  $f$ , then  $f$  is holomorphic in an annulus  $0 < |z - z_0| < R$  for some  $R$ . The corresponding Laurent expansion is called the Laurent expansion around  $z_0$ . Consider the  $-1$ -st coefficient of this Laurent series.

$$a_{-1} = \frac{1}{2\pi i} \int_{\gamma} f(z) dz.$$

If you integrate a Laurent series, only  $a_{-1}$  remains; other terms vanish. What remains is usually called a residue.

$$a_{-1} = \operatorname{Res}(f; z_0).$$

Often  $a_{-1}$  is easy to compute from  $f(z)$  and if that's the case integration has become easy.

# Cauchy Residue Theorem

Suppose  $f$  is given and  $\gamma$  is given.

# Cauchy Residue Theorem

Suppose  $f$  is given and  $\gamma$  is given. Suppose there are finitely many isolated singularities of  $f$  inside  $\gamma$ ; say  $z_1, z_2, \dots, z_n$ .

# Cauchy Residue Theorem

Suppose  $f$  is given and  $\gamma$  is given. Suppose there are finitely many isolated singularities of  $f$  inside  $\gamma$ ; say  $z_1, z_2, \dots, z_n$ . What's  $\int_{\gamma} f(z) dz$  ?


# Cauchy Residue Theorem

Suppose  $f$  is given and  $\gamma$  is given. Suppose there are finitely many isolated singularities of  $f$  inside  $\gamma$ ; say  $z_1, z_2, \dots, z_n$ . What's  $\int_{\gamma} f(z) dz$  ?

## Theorem (Cauchy Residue Theorem)

$$\int_{\gamma} f(z) dz = 2\pi i \cdot \sum_{i=1}^n \text{Res}(f, z_i).$$

Proof : We have already seen the proof in the previous lectures. The following figure should remind you of the proof. (Here the case of 1 singularity is considered; similarly one can handle more singular points)




# Cauchy Residue Theorem

Suppose  $f$  is given and  $\gamma$  is given. Suppose there are finitely many isolated singularities of  $f$  inside  $\gamma$ ; say  $z_1, z_2, \dots, z_n$ . What's  $\int_{\gamma} f(z) dz$  ?

## Theorem (Cauchy Residue Theorem)

$$\int_{\gamma} f(z) dz = 2\pi i \cdot \sum_{i=1}^n \text{Res}(f, z_i).$$

Proof : We have already seen the proof in the previous lectures. The following figure should remind you of the proof. (Here the case of 1 singularity is considered; similarly one can handle more singular points)



Why did the mathematician name his dog Cauchy?

Why did the mathematician name his dog Cauchy?





Why did the mathematician name his dog Cauchy?




B'coz it left a residue at every pole!

What's the contour integral over Western Europe?

What's the contour integral over Western Europe?



What's the contour integral over Western Europe?



Zero. All the poles are in Eastern Europe.

Modification: Actually there are poles in Western Europe but they are all removable !!

# Principal Part of the Laurent Series

If  $\sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$  is the Laurent expansion around  $z_0$ , then its principal part is

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n.$$

# Principal Part of the Laurent Series

If  $\sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$  is the Laurent expansion around  $z_0$ , then its principal part is

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n.$$

Note that:

- removable iff principal part is zero.
- pole iff principal part is finite.
- essential iff principal part is infinite.

Proof (Easy exercise using previous slides).

# Residue at a Pole

If the isolated singularity is removable, then the residue is trivial. If the isolated singularity is a pole, then the residue is trivial to compute. If  $z_0$  is a pole, can write

$$f(z) = \frac{a_{-m}}{(z - z_0)^m} + \dots + \frac{a_{-1}}{(z - z_0)} + a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \dots$$



# Residue at a Pole

If the isolated singularity is removable, then the residue is trivial. If the isolated singularity is a pole, then the residue is trivial to compute. If  $z_0$  is a pole, can write

$$f(z) = \frac{a_{-m}}{(z-z_0)^m} + \dots + \frac{a_{-1}}{(z-z_0)} + a_0 + a_1(z-z_0) + a_2(z-z_0)^2 + \dots$$

Thus,

$$g(z) = (z-z_0)^m f(z) = a_{-m} + \dots + a_{-1}(z-z_0)^{m-1} + a_0(z-z_0)^m + \dots$$

# Residue at a Pole

If the isolated singularity is removable, then the residue is trivial. If the isolated singularity is a pole, then the residue is trivial to compute. If  $z_0$  is a pole, can write

$$f(z) = \frac{a_{-m}}{(z-z_0)^m} + \dots + \frac{a_{-1}}{(z-z_0)} + a_0 + a_1(z-z_0) + a_2(z-z_0)^2 + \dots$$

Thus,

$$g(z) = (z-z_0)^m f(z) = a_{-m} + \dots + a_{-1}(z-z_0)^{m-1} + a_0(z-z_0)^m + \dots$$

Thus,  $g$  is holomorphic and

$$a_{-1} = \frac{g^{(m-1)}(z_0)}{(m-1)!}.$$

# Example

Find the Laurent Series expansion of  $\frac{1}{(z-1)(z-2)}$  in the annulus  $1 < |z| < 2$ .

# Example

Find the Laurent Series expansion of  $\frac{1}{(z-1)(z-2)}$  in the annulus  $1 < |z| < 2$ .

$$\frac{1}{(z-1)(z-2)} = \frac{1}{z-2} - \frac{1}{z-1}.$$

For the first term,  $\frac{1}{z-2} = -\frac{1}{2}\left(\frac{1}{1-\frac{z}{2}}\right) = -\frac{1}{2} \sum_{n=0}^{\infty} \left(\frac{z}{2}\right)^n.$

For the second term,  $\frac{1}{z-1} = \frac{1}{z}\left(\frac{1}{1-1/z}\right) = \sum_{n=1}^{\infty} \frac{1}{z^n}.$

# Example

Find the Laurent Series expansion of  $\frac{1}{(z-1)(z-2)}$  in the annulus  $1 < |z| < 2$ .

$$\frac{1}{(z-1)(z-2)} = \frac{1}{z-2} - \frac{1}{z-1}.$$

For the first term,  $\frac{1}{z-2} = -\frac{1}{2}\left(\frac{1}{1-\frac{z}{2}}\right) = -\frac{1}{2} \sum_{n=0}^{\infty} \left(\frac{z}{2}\right)^n.$

For the second term,  $\frac{1}{z-1} = \frac{1}{z}\left(\frac{1}{1-1/z}\right) = \sum_{n=1}^{\infty} \frac{1}{z^n}.$

Putting the two together we get the desired Laurent Series Expansion.

# Example

Determine the Laurent series of  $e^{1/z}$  around the point 0.

$$\begin{aligned} e^{1/z} &= \sum_0^{\infty} \frac{1}{n!} \frac{1}{z^n} \\ &= \sum_{-\infty}^0 \frac{z^n}{(-n)!} \end{aligned}$$

# Example

Determine the Laurent series of  $\sin\left(\frac{1}{z-1}\right)$  in the range  $1 < |z| < 2$ .

$\sin\left(\frac{1}{z-1}\right)$

# Example

Determine the Laurent series of  $\sin\left(\frac{1}{z-1}\right)$  in the range  $1 < |z| < 2$ .

$$\begin{aligned} & \sin\left(\frac{1}{z-1}\right) \\ &= \sin\left(\frac{1}{z(1-\frac{1}{z})}\right) \end{aligned}$$



# Example

Determine the Laurent series of  $\sin(\frac{1}{z-1})$  in the range  $1 < |z| < 2$ .

$$\sin(\frac{1}{z-1})$$

$$= \sin(\frac{1}{z(1-\frac{1}{z})})$$

$$= \sin(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots)$$

# Example

Determine the Laurent series of  $\sin(\frac{1}{z-1})$  in the range  $1 < |z| < 2$ .

$$\sin(\frac{1}{z-1})$$

$$= \sin(\frac{1}{z(1-\frac{1}{z})})$$

$$= \sin(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots)$$

$$= (\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots) - \frac{1}{3!}(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots)^3 + \frac{1}{5!}(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \cdots)^5 + \cdots$$

# Example

Determine the Laurent series of  $\sin(\frac{1}{z-1})$  in the range  $1 < |z| < 2$ .

$$\sin(\frac{1}{z-1})$$

$$= \sin(\frac{1}{z(1-\frac{1}{z})})$$

$$= \sin(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \dots)$$

$$= (\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \dots) - \frac{1}{3!}(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \dots)^3 + \frac{1}{5!}(\frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} \dots)^5 + \dots$$

$$= \frac{1}{z} + \frac{1}{z^2} + \frac{5}{6} \frac{1}{z^3} + \frac{1}{2} \frac{1}{z^4} + \dots$$