

CS/ECE/ME532 Assignment 5

1. Here we continue the problem studied in Activity 11. Let a 4-by-2 matrix \mathbf{X} have

$$\text{SVD } \mathbf{X} = \mathbf{U}\mathbf{S}\mathbf{V}^T \text{ where } \mathbf{U} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix}, \mathbf{S} = \begin{bmatrix} 1 & 0 \\ 0 & \gamma \end{bmatrix}, \text{ and } \mathbf{V} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$\text{Let } \mathbf{y} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}.$$

- a) The ratio of the largest to the smallest singular values is termed the condition number of \mathbf{X} . Find the condition number if $\gamma = 0.1$, and $\gamma = 10^{-8}$. Solve $\mathbf{X}\mathbf{w} = \mathbf{y}$ for \mathbf{w} and find $\|\mathbf{w}\|_2^2$ for these two values of γ .

- b) A system of linear equations with a large condition number is said to be “ill-conditioned”. One consequence of an ill-conditioned system of equations is solutions with large norms as you found in the previous part of this problem. A second consequence is that the solution is very sensitive to small errors in \mathbf{y} such as may

result from measurement error or numerical error. Suppose $\mathbf{y} = \begin{bmatrix} 1 + \epsilon \\ 0 \\ 0 \\ 1 \end{bmatrix}$. Write

$\mathbf{w} = \mathbf{w}_o + \mathbf{w}_\epsilon$ where \mathbf{w}_o is the solution for arbitrary γ when $\epsilon = 0$ and \mathbf{w}_ϵ is the perturbation in that solution due to some error $\epsilon \neq 0$. How does the norm of the perturbation due to $\epsilon \neq 0$, $\|\mathbf{w}_\epsilon\|_2^2$, depend on the condition number? Find $\|\mathbf{w}_\epsilon\|_2^2$ for $\epsilon = 0.01$ and $\gamma = 0.1$ and $\gamma = 10^{-8}$.

- c) Now consider a “low-rank” inverse. Instead of writing

$$(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T = \sum_{i=1}^p \frac{1}{\sigma_i} \mathbf{v}_i \mathbf{u}_i^T$$

where p is the number of columns of \mathbf{X} (assumed less than the number of rows), we approximate

$$(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \approx \sum_{i=1}^r \frac{1}{\sigma_i} \mathbf{v}_i \mathbf{u}_i^T$$

In this approximation we only invert the largest r singular values, and ignore all of them smaller than σ_r . Use $r = 1$ in the low-rank inverse to find $\mathbf{w} = \mathbf{w}_o + \mathbf{w}_\epsilon$

where $\mathbf{y} = \begin{bmatrix} 1 + \epsilon \\ 0 \\ 0 \\ 1 \end{bmatrix}$ as in part b). Compare the results to part b).