

### **Overview**

#### **Burden classification**

#### The Global Nutrition Report classifies this country as experiencing three forms of malnutrition – overweight, anaemia and stunting

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: Thresholds for a country having the form or not: stunting in children aged under 5 years  $\geq$ 20%; anaemia in women of reproductive age  $\geq$ 20%; overweight (body mass index  $\geq$ 25) in adult women aged  $\geq$ 18 years  $\geq$ 35%.

#### Progress against global nutrition targets 2018



Under-5 stunting



Under-5 wasting
On course



Under-5 overweight



WRA anaemia





Exclusive breastfeeding
On course



Adult female obesity
No progress or worsening



Adult male obesity
No progress or worsening



Adult female diabetes
No progress or worsening



Adult male diabetes

No progress or worsening


INO progress or worsering

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF global databases: Infant and Young Child Feeding, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates. See Appendix 1 of the 2018 Global Nutrition Report for details of the methods and sources used to assess progress towards global nutrition targets.

## **Economics and demography**

#### Poverty rates (%) and GDP (PPP\$)



**Source**: World Bank 2018. **Note**: GDP = gross domestic product. PPP = purchasing power

#### Under-5 mortality (per 000 live births)



Source: UN Inter-agency Group for Child Mortality Estimation 2018.

#### Income inequality

Gini index score <sup>1</sup>		Gini index rank <sup>2</sup>	Year
	38	84	2010

Source: World Bank 2018.

**Notes**: 0 = perfect equality, 100 = perfect inequality. Countries are ranked from most equal (1) to most unequal (155). 2

#### **Population**


Population (000)	276	2017
Under-5 population (000)	35	2018
Rural (%)	75	2017
≥65 years (000)	13	2018

Source: UN Population Division 2017.


# **Underlying determinants**

#### Food supply

Source: FAOSTAT 2018.



#### Drinking water coverage (% population)



#### **Gender-related determinants**

Early childbearing: births by age 18 (%) <sup>1</sup>	13	2013
Gender Inequality Index (score*) <sup>2</sup>	NA	NA
Gender Inequality Index (country rank) <sup>2</sup>	NA	NA

Sources: UNICEF 2018; UNDP 2018.<sup>2</sup>


**Note**: \*0 = low inequality, 1 = high inequality.

# Population density of health workers per 1,000 people

Physicians	0.19	2012
Nurses and midwives	2.22	2012
Community health workers	0.94	2008


**Source**: WHO's Global Health Workforce Statistics, OECD, supplemented by country data.

# Female secondary education




Source: UNESCO Institute for Statistics 2018.

#### Sanitation coverage (% population)




#### Government expenditures (% total)




1

# Children (under 5) nutrition status


#### Under 5 by gender: wasting (%)



#### Under 5 by gender: stunting (%)




#### Under 5 by gender: overweight (%)



#### Under 5 by household income: wasting (%)

malnutrition estimates.




#### Under 5 by household income: stunting (%)



#### Under 5 by household income: overweight (%)






#### Under 5 by household location: wasting (%) | Under 5 by household location: stunting (%)




#### Under 5 by household location: overweight (%)



#### Under-5 coexistence of wasting, stunting and overweight





# **Child feeding practices**




Sources: UNICEF, Division of Data Research and Policy (2018), Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2018.


# Children and adolescent (aged 5-19) nutrition status









## **Adult nutrition status**









# Male Female 25 23.4 24.2 20 15 10 1999 2003 2007 2011 2015


| Source: NCD Risk Factor Collaboration.


Adult by gender: raised blood pressure (%)





## **Dietary needs**





**Source**: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: Men and women aged 25 and older. TMREL = theoretical minimum risk exposure level.

# Financial resources and policy, legislation and institutional arrangements

#### **Development assistance** Basic nutrition ODA received -% of total ODA 1.00 suggest of the state of the sta 0.6 % ᅌ 0.4 6 0.00 0.0 2014 2016 2008 2010 2012

**Sources**: Development Initiatives based on OECD Development Assistance Committee (DAC) Creditor Reporting System (CRS).

**Notes**: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2016 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

#### **National policies**

Mandatory legislation for salt iodisation	No
Sugar-sweetened beverage tax	Yes
Food-based dietary guidelines	NA
Multisectoral comprehensive nutrition plan	Yes

**Sources**: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

#### Targets included in national (nutrition or other) plan

Stunting	Anaemia	Low birth weight	Child overweight	Exclusive breastfeeding	Wasting	Salt intake	Overweight adults and adolescents
Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes

Sources: WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

# Intervention coverage

Coverage/practice indicator	%	Male	Female	Year
Children 0–59 months with diarrhoea who received zinc treatment	NA	NA	NA	NA
Children 6–59 months who received vitamin A supplements in last 6 months	NA	NA	NA	NA
Children 6–59 months given iron supplements in past 7 days	NA	NA	NA	NA
Women with a birth in last five years who received iron and folic acid during their most recent pregnancy	NA		NA	NA
Household consumption of any iodised salt	NA	NA	NA	NA

Sources: Kothari M. and Huestis A., based on 2016 Global Nutrition Report and UNICEF global databases, 2018.

Notes: Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005–2017.