

## **Overview**

#### **Burden classification**

#### The Global Nutrition Report classifies this country as experiencing two forms of malnutrition – overweight and anaemia

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: Thresholds for a country having the form or not: stunting in children aged under 5 years  $\geq$ 20%; anaemia in women of reproductive age  $\geq$ 20%; overweight (body mass index  $\geq$ 25) in adult women aged  $\geq$ 18 years  $\geq$ 35%.

## Progress against global nutrition targets 2018



**Under-5 stunting** 



**Under-5 wasting** 



**Under-5 overweight** 



WRA anaemia

No progress or worsening



**Exclusive breastfeeding** 



Adult female obesity No progress or worsening



Adult male obesity No progress or worsening



Adult female diabetes No progress or worsening




Adult male diabetes

No progress or worsening

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF global databases: Infant and Young Child Feeding, NCD Risk Factor Collaboration, WHO Global Health Observatory. Notes: The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates. See Appendix 1 of the 2018 Global Nutrition Report for details of the methods and sources used to assess progress towards global nutrition targets.

## **Economics and demography**

## Poverty rates (%) and GDP (PPP\$)



Source: World Bank 2018.

**Note**: GDP = gross domestic product. PPP = purchasing power

## Under-5 mortality (per 000 live births)



Source: UN Inter-agency Group for Child Mortality Estimation 2018.

## Income inequality

| Gini index score <sup>1</sup> | Gini index rank <sup>2</sup> | Year |
|-------------------------------|------------------------------|------|
| NA                            | NA                           | NA   |

Source: World Bank 2018.

**Notes**: 0 = perfect equality, 100 = perfect inequality. Countries are ranked from most equal (1) to most unequal (155).

## **Population**


| Population (000)         | 286 | 2017 |
|--------------------------|-----|------|
| Under-5 population (000) | 17  | 2018 |
| Rural (%)                | 69  | 2017 |
| ≥65 years (000)          | 44  | 2018 |

Source: UN Population Division 2017.


# **Underlying determinants**

## Food supply

Source: FAOSTAT 2018.



## Drinking water coverage (% population)



## **Gender-related determinants**

| Early childbearing: births by age 18 (%) <sup>1</sup> | 7    | 2012 |
|---|------|------|
| Gender Inequality Index (score*) <sup>2</sup>         | 0.28 | 2017 |
| Gender Inequality Index (country rank) <sup>2</sup>   | 60   | 2017 |

Sources: UNICEF 2018: UNDP 2018.<sup>2</sup>


Note: \*0 = low inequality, 1 = high inequality.

## Population density of health workers per 1,000 people

| Physicians               | 1.81 | 2010 |
|--------------------------|------|------|
| Nurses and midwives      | 4.86 | 2010 |
| Community health workers | NA   | NA   |


Source: WHO's Global Health Workforce Statistics, OECD, supplemented by country data.

## Female secondary education enrolment (net, % population)




Source: UNESCO Institute for Statistics 2018.

## Sanitation coverage (% population)




## Government expenditures (% total)




# Children (under 5) nutrition status


#### Under 5 by gender: wasting (%)



Under 5 by gender: stunting (%)




#### Under 5 by gender: overweight (%)




Under 5 by household income: wasting (%)


malnutrition estimates.




Under 5 by household income: stunting (%)




Under 5 by household income: overweight (%)







Under 5 by household location: wasting (%) | Under 5 by household location: stunting (%)



Under 5 by household location: overweight (%)




Under-5 coexistence of wasting, stunting and overweight



Sources: UNICEF, Division of Data Research and Policy (2018), Global UNICEF Global Databases: Overlapping Stunting, Wasting and Overweight, New York, May 2018.


Notes: Percentage of children under 5 years of age who experience different and overlapping forms of malnutrition.


## **Child feeding practices**




Sources: UNICEF, Division of Data Research and Policy (2018), Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2018.


## Children and adolescent (aged 5-19) nutrition status









## **Adult nutrition status**









# Male Female 30 29.1 27 23.4 21.7 20 10 1999 2003 2007 2011 2015

Source: NCD Risk Factor Collaboration.


Adult by gender: raised blood pressure (%)






## **Dietary needs**

Consumption of food groups and components, 2016






Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation. Notes: Men and women aged 25 and older. TMREL = theoretical minimum risk exposure level.

# Financial resources and policy, legislation and institutional arrangements

## **Development assistance**



**Notes**: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2016 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

## **National policies**

| Mandatory legislation for salt iodisation  | NA  |
|--|-----|
| Sugar-sweetened beverage tax               | Yes |
| Food-based dietary guidelines              | Yes |
| Multisectoral comprehensive nutrition plan | Yes |

**Sources**: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

## Targets included in national (nutrition or other) plan

| Stunting | Anaemia | Low birth weight | Child overweight | Exclusive<br>breastfeeding | Wasting | Salt intake | Overweight adults and adolescents |
|----------|---------|------------------|------------------|----------------------------|---------|-------------|-----------------------------------|
| No       | No      | No               | Yes              | Yes                        | No      | Yes         | No                                |

# Intervention coverage

| Coverage/practice indicator   | %  | Male | Female | Year |
|---|----|------|--------|------|
| Children 0–59 months with diarrhoea who received zinc treatment   | NA | NA   | NA     | NA   |
| Children 6–59 months who received vitamin A supplements in last 6 months                                  | NA | NA   | NA     | NA   |
| Children 6–59 months given iron supplements in past 7 days  | NA | NA   | NA     | NA   |
| Women with a birth in last five years who received iron and folic acid during their most recent pregnancy | NA |      | NA     | NA   |
| Household consumption of any iodised salt   | NA | NA   | NA     | NA   |

Sources: Kothari M. and Huestis A., based on 2016 Global Nutrition Report and UNICEF global databases, 2018. Notes: Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005–2017.