Chile



Overview

Burden classification

The Global Nutrition Report classifies this country as experiencing one form of malnutrition – overweight only

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: Thresholds for a country having the form or not: stunting in children aged under 5 years \geq 20%; anaemia in women of reproductive age \geq 20%; overweight (body mass index \geq 25) in adult women aged \geq 18 years \geq 35%.

Progress against global nutrition targets 2018



Under-5 stunting
On course



Under-5 wasting
On course



Under-5 overweightOn course



WRA anaemia

No progress or worsening



Exclusive breastfeeding

NA

nû

Adult female obesity
No progress or worsening



Adult male obesity

No progress or worsening



Adult female diabetes
No progress or worsening



Adult male diabetes


No progress or worsening

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF global databases: Infant and Young Child Feeding, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates. See Appendix 1 of the 2018 Global Nutrition Report for details of the methods and sources used to assess progress towards global nutrition targets.

Economics and demography

Poverty rates (%) and GDP (PPP\$)



Source: World Bank 2018.

Note: GDP = gross domestic product. PPP = purchasing power

Under-5 mortality (per 000 live births)



Source: UN Inter-agency Group for Child Mortality Estimation 2018.

Income inequality

Gini index score ¹		Gini index rank ²	Year		
	48	137	2015		

Source: World Bank 2018.

Notes: 0 = perfect equality, 100 = perfect inequality. Countries are ranked from most equal (1) to most unequal (155). 2

Population


Population (000)	18,055	2017
Under-5 population (000)	1,179	2018
Rural (%)	13	2017
>65 years (000)	2,084	2018

Source: UN Population Division 2017.


Underlying determinants

Food supply

Source: FAOSTAT 2018.



Drinking water coverage (% population)



Gender-related determinants

Early childbearing: births by age 18 (%) ¹	NA	NA
Gender Inequality Index (score*) ²	0.32	2016
Gender Inequality Index (country rank) ²	70	2016

Sources: UNICEF 2018; UNDP 2018. 2


Note: *0 = low inequality, 1 = high inequality.

Population density of health workers per 1,000 people

Physicians	1.03	2009
Nurses and midwives	0.14	2009
Community health workers	NA	NA


Source: WHO's Global Health Workforce Statistics, OECD, supplemented by country data.

Female secondary education




Source: UNESCO Institute for Statistics 2018.

Sanitation coverage (% population)



Government expenditures (% total)




Chile

Children (under 5) nutrition status


Under 5 by gender: wasting (%)

Girls
Boys
Both



Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.

Under 5 by gender: stunting (%)



Under 5 by gender: overweight (%)



No data

Under 5 by household income: wasting (%) | Under 5 by household income: stunting (%) |

Under 5 by household income: overweight (%)

No data

No data

Sources: UNICEF/WHO/World Bank Group: Joint child

malnutrition estimates.

Under 5 by household location: wasting (%) | Under 5 by household location: stunting (%) |

Ourban Rural 9.6 7.6

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.

No data

Under 5 by household location: overweight (%)

No data

Under-5 coexistence of wasting, stunting and overweight

No data


Child feeding practices

No data


Sources: UNICEF, Division of Data Research and Policy (2018), Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2018.

Children and adolescent (aged 5-19) nutrition status

Ages 5-19 by gender: underweight (%)



Ages 5-19 by gender: overweight (%)




Ages 5-19 by gender: obesity (%)




Adult nutrition status


Adult by gender: diabetes (%)




Adult by gender: overweight (%)




Adult by gender: obesity (%)




Adult by gender: raised blood pressure (%) |



Adult by gender: anaemia in WRA (%)




Adult: sodium intake (grams per day)




Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation.



Dietary needs





Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation. Notes: Men and women aged 25 and older. TMREL = theoretical minimum risk exposure level.

Financial resources and policy, legislation and institutional arrangements

Development assistance Basic nutrition ODA received -% of total ODA 1.00 1.00 suggest of the state of the sta 0.50 0.25 0.00 0.00 2016

2012

2014

Sources: Development Initiatives based on OECD Development Assistance Committee (DAC) Creditor Reporting System (CRS).

Notes: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2016 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

National policies

Mandatory legislation for salt iodisation	Yes
Sugar-sweetened beverage tax	Yes
Food-based dietary guidelines	Yes
Multisectoral comprehensive nutrition plan	Yes

Sources: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

Targets included in national (nutrition or other) plan

2010

Stunting	Anaemia	Low birth weight	Child overweight	Exclusive breastfeeding	Wasting	Salt intake	Overweight adults and adolescents
Yes	No	Yes	Yes	Yes	Yes	Yes	Yes

Intervention coverage

2008

Coverage/practice indicator	%	Male	Female	Year
Children 0–59 months with diarrhoea who received zinc treatment	NA	NA	NA	NA
Children 6–59 months who received vitamin A supplements in last 6 months	NA	NA	NA	NA
Children 6–59 months given iron supplements in past 7 days	NA	NA	NA	NA
Women with a birth in last five years who received iron and folic acid during their most recent pregnancy	NA		NA	NA
Household consumption of any iodised salt	NA	NA	NA	NA

Sources: Kothari M. and Huestis A., based on 2016 Global Nutrition Report and UNICEF global databases, 2018. Notes: Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005–2017.