## Canada



### **Overview**

### **Burden classification**

#### There is insufficient data for the Global Nutrition Report to classify this country's burden

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: Thresholds for a country having the form or not: stunting in children aged under 5 years  $\geq$ 20%; anaemia in women of reproductive age  $\geq$ 20%; overweight (body mass index  $\geq$ 25) in adult women aged  $\geq$ 18 years  $\geq$ 35%.

### Progress against global nutrition targets 2018



**Under-5 stunting** 



**Under-5** wasting



**Under-5 overweight** 



WRA anaemia

No progress or worsening



**Exclusive breastfeeding** 



Adult female obesity No progress or worsening



Adult male obesity No progress or worsening



Adult female diabetes On course




Adult male diabetes

No progress or worsening

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF global databases: Infant and Young Child Feeding, NCD Risk Factor Collaboration, WHO Global Health Observatory. Notes: The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates. See Appendix 1 of the 2018 Global Nutrition Report for details of the methods and sources used to assess progress towards global nutrition targets.

## **Economics and demography**

### Poverty rates (%) and GDP (PPP\$)



Source: World Bank 2018. Note: GDP = gross domestic product. PPP = purchasing power

### **Under-5 mortality (per 000 live births)**



Source: UN Inter-agency Group for Child Mortality Estimation 2018.

### Income inequality

Gini index score <sup>1</sup>	Gini index rank <sup>2</sup>	Year
34	52	2013

Source: World Bank 2018.

**Notes**: 0 = perfect equality, 100 = perfect inequality. Countries are ranked from most equal (1) to most unequal (155).


#### **Population**

Population (000)	36,624	2017	
Under-5 population (000)	1,958	2018	
Rural (%)	19	2017	
≥65 years (000)	6,430	2018	
· · · · · · · · · · · · · · · · · · ·			

Source: UN Population Division 2017.

## **Underlying determinants**

### Food supply



Source: FAOSTAT 2018.

### **Gender-related determinants**

Early childbearing: births by age 18 (%) <sup>1</sup>	NA	NA
Gender Inequality Index (score*) <sup>2</sup>	0.09	2017
Gender Inequality Index (country	20	2017

Sources: UNICEF 2018; UNDP 2018.<sup>2</sup>


**Note**: \*0 = low inequality, 1 = high inequality.

### Population density of health workers per 1,000 people

Physicians	2.54	2015
Nurses and midwives	9.84	2015
Community health workers	NA	NA


Source: WHO's Global Health Workforce Statistics, OECD, supplemented by country data

### Female secondary education enrolment (net, % population)




Source: UNESCO Institute for Statistics 2018.


### **Drinking water coverage (% population)**



### Sanitation coverage (% population)



### Government expenditures (% total)



# Canada

Under 5 by gender: wasting (%)	Under 5 by gender: stunting (%)	Under 5 by gender: overweight (%)
No data  Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.	No data	2004
Inder 5 by household income: wasting (%	b)   Under 5 by household income: stunting (%)	)   Under 5 by household income: overweight (%
No data  Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.	No data	No data
nder 5 by household location: wasting (%	b)   Under 5 by household location: stunting (%)	)   Under 5 by household location: overweight (9
No data	No data	No data
<b>Sources</b> : UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.		
Under-5 coexistence of wasting, stunting	g and overweight	
	No data	

Sources: UNICEF, Division of Data Research and Policy (2018), Global UNICEF Global Databases: Overlapping Stunting, Wasting and Overweight, New York, May 2018.


Notes: Percentage of children under 5 years of age who experience different and overlapping forms of malnutrition.


## **Child feeding practices**


No data

Sources: UNICEF, Division of Data Research and Policy (2018), Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2018.

## Children and adolescent (aged 5-19) nutrition status







### **Adult nutrition status**


2007


2011


2015

2003


| Source: NCD Risk Factor Collaboration.








### Adult by gender: raised blood pressure (%)



### Adult: anaemia in WRA (%)



### Adult: sodium intake (grams per day)




**Source**: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

## Canada

## **Dietary needs**

### Consumption of food groups and components, 2016






Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation. Notes: Men and women aged 25 and older. TMREL = theoretical minimum risk exposure level.

## Financial resources and policy, legislation and institutional arrangements

## **Development assistance** -% of total ODA



Sources: Development Initiatives based on OECD Development Assistance Committee (DAC)

Creditor Reporting System (CRS).

**Notes**: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2016 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

### **National policies**

Mandatory legislation for salt iodisation	Yes
Sugar-sweetened beverage tax	No
Food-based dietary guidelines	Yes
Multisectoral comprehensive nutrition plan	Yes

**Sources**: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

### Targets included in national (nutrition or other) plan

Stunting	Anaemia	Low birth weight	Child overweight	Exclusive breastfeeding	Wasting	Salt intake	Overweight adults and adolescents
No	No	No	Yes	Yes	No	Yes	Yes

## Intervention coverage

Coverage/practice indicator	%	Male	Female	Year
Children 0–59 months with diarrhoea who received zinc treatment		NA	NA	NA
Children 6–59 months who received vitamin A supplements in last 6 months		NA	NA	NA
Children 6–59 months given iron supplements in past 7 days		NA	NA	NA
Women with a birth in last five years who received iron and folic acid during their most recent pregnancy			NA	NA
Household consumption of any iodised salt	NA	NA	NA	NA

Sources: Kothari M. and Huestis A., based on 2016 Global Nutrition Report and UNICEF global databases, 2018. Notes: Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005–2017.