


Overview

Burden classification

The Global Nutrition Report classifies this country as experiencing three forms of malnutrition – overweight, anaemia and stunting

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.  
Notes: Thresholds for a country having the form or not: stunting in children aged under 5 years ≥20%; anaemia in women of reproductive age ≥20%; overweight (body mass index ≥25) in adult women aged ≥18 years ≥35%.


Progress against global nutrition targets 2018



Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF global databases: Infant and Young Child Feeding, NCD Risk Factor Collaboration, WHO Global Health Observatory.  
Notes: The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates. See Appendix 1 of the 2018 Global Nutrition Report for details of the methods and sources used to assess progress towards global nutrition targets.


Economics and demography

Poverty rates (%) and GDP (PPP\$)



Source: World Bank 2018.  
Note: GDP = gross domestic product. PPP = purchasing power parity.

Under-5 mortality (per 000 live births)



Source: UN Inter-agency Group for Child Mortality Estimation 2018.

Income inequality

| Gini index score <sup>1</sup> | Gini index rank <sup>2</sup> | Year |
|-------------------------------|------------------------------|------|
| 34                            | 48                           | 2012 |

Source: World Bank 2018.  
Notes: 0 = perfect equality, 100 = perfect inequality.<sup>1</sup> Countries are ranked from most equal (1) to most unequal (155).<sup>2</sup>


Population

|                          |        |      |
|--------------------------|--------|------|
| Population (000)         | 12,717 | 2017 |
| Under-5 population (000) | 2,052  | 2018 |
| Rural (%)                | 64     | 2017 |
| ≥65 years (000)          | 415    | 2018 |

Source: UN Population Division 2017.

Underlying determinants

Food supply



Source: FAOSTAT 2018.

Gender-related determinants

|   |    |      |
|---|----|------|
| Early childbearing: births by age 18 (%) <sup>1</sup> | 40 | 2012 |
| Gender Inequality Index (score*) <sup>2</sup>         | NA | NA   |
| Gender Inequality Index (country rank) <sup>2</sup>   | NA | NA   |


Sources: UNICEF 2018;<sup>1</sup> UNDP 2018.<sup>2</sup>  
Note: \*0 = low inequality, 1 = high inequality.

Population density of health workers per 1,000 people

|                          |      |      |
|--------------------------|------|------|
| Physicians               | 0.07 | 2016 |
| Nurses and midwives      | 0.37 | 2016 |
| Community health workers | 0.52 | 2016 |


Source: WHO's Global Health Workforce Statistics, OECD, supplemented by country data.

Female secondary education enrolment (net, % population)




Source: UNESCO Institute for Statistics 2018.

Drinking water coverage (% population)



Source: WHO/UNICEF Joint Monitoring Programme 2017.

Sanitation coverage (% population)



Source: WHO/UNICEF Joint Monitoring Programme 2017.


Government expenditures (% total)



Source: IFPRI 2015.


Children (under 5) nutrition status

Under 5 by gender: wasting (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.


Under 5 by gender: stunting (%)



Under 5 by gender: overweight (%)




Under 5 by household income: wasting (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.


Under 5 by household income: stunting (%)



Under 5 by household income: overweight (%)




Under 5 by household location: wasting (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.


Under 5 by household location: stunting (%)



Under 5 by household location: overweight (%)




Under-5 coexistence of wasting, stunting and overweight




Sources: UNICEF, Division of Data Research and Policy (2018), Global UNICEF Global Databases: Overlapping Stunting, Wasting and Overweight, New York, May 2018.  
Notes: Percentage of children under 5 years of age who experience different and overlapping forms of malnutrition.

Child feeding practices





Sources: UNICEF, Division of Data Research and Policy (2018), Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2018.

Children and adolescent (aged 5–19) nutrition status



Adult nutrition status



Dietary needs


Consumption of food groups and components, 2016



Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation.  
Notes: Men and women aged 25 and older. TMREL = theoretical minimum risk exposure level.

Financial resources and policy, legislation and institutional arrangements

Development assistance



Sources: Development Initiatives based on OECD Development Assistance Committee (DAC) Creditor Reporting System (CRS).  
Notes: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2016 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

National policies

|  |     |
|--|-----|
| Mandatory legislation for salt iodisation  | Yes |
| Sugar-sweetened beverage tax               | No  |
| Food-based dietary guidelines              | NA  |
| Multisectoral comprehensive nutrition plan | Yes |

Sources: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

Targets included in national (nutrition or other) plan

|          |         |                  |                  |                         |         |             |                                   |
|----------|---------|------------------|------------------|-------------------------|---------|-------------|-----------------------------------|
| Stunting | Anaemia | Low birth weight | Child overweight | Exclusive breastfeeding | Wasting | Salt intake | Overweight adults and adolescents |
| Yes      | Yes     | Yes              | Yes              | Yes                     | Yes     | Yes         | Yes                               |

Sources: WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

Intervention coverage

| Coverage/practice indicator   | Total (%) | Male (%) | Female (%) | Year |
|---|-----------|----------|------------|------|
| Children 0–59 months with diarrhoea who received zinc treatment   | 0         | NA       | NA         | 2012 |
| Children 6–59 months who received vitamin A supplements in last 6 months                                  | 41        | 42       | 40         | 2012 |
| Children 6–59 months given iron supplements in past 7 days  | 12        | 12       | 11         | 2012 |
| Women with a birth in last five years who received iron and folic acid during their most recent pregnancy | 81        |          | 81         | 2012 |
| Household consumption of any iodised salt   | 64        | NA       | NA         | 2012 |

Sources: Kothari M. and Huestis A., based on 2016 Global Nutrition Report and UNICEF global databases, 2018.  
Notes: Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005–2017.