




Southern Asia

Overview


Progress against global nutrition targets 2018



Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF global databases: Infant and Young Child Feeding, NCD Risk Factor Collaboration, WHO Global Health Observatory.
Notes: The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates. See Appendix 1 of the 2018 Global Nutrition Report for details of the methods and sources used to assess progress towards global nutrition targets.


Economics and demography

Poverty rates (%) and GDP (PPP\$)



Source: World Bank 2018.
Note: GDP = gross domestic product. PPP = purchasing power parity. n = 8 (poverty); n = 9 (GDP).

Under-5 mortality (per 000 live births)



Source: UN Inter-agency Group for Child Mortality Estimation 2018.


Population

| | | |
|-------------------------------|-------|------|
| Population (millions) | 1,869 | 2017 |
| Under-5 population (millions) | 177 | 2018 |
| Rural (%) | 65 | 2017 |
| ≥65 years (millions) | 111 | 2018 |

Source: UN Population Division 2017.
Note: n = 9 (total); n = 9 (under-5); n = 9 (rural); n = 9 (over 65)

Underlying determinants

Food supply



Source: FAOSTAT 2018.


Gender-related determinants

| | | |
|---|------|------|
| Early childbearing: births by age 18 (%) | 19 | 2015 |
| Population density of health workers per 1,000 people | | |
| Physicians | 0.78 | 2016 |
| Nurses and midwives | 1.72 | 2016 |
| Community health workers | 0.51 | 2016 |


Source: UNICEF 2018.
Note: n = 9.

Source: WHO's Global Health Workforce Statistics, OECD, supplemented by country data. n = 9


Female secondary education enrolment (net, % population)




Drinking water coverage (% population)



Sanitation coverage (% population)




Government expenditures (% total)



Southern Asia


Children (under 5) nutrition status

Under 5: wasting (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.

Under 5: stunting (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.

Under 5: overweight (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.

Under 5 by household income: wasting (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates. n = NA

Under 5 by household income: stunting (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates. n = 4

Under 5 by household income: overweight (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates. n = 4

Under 5 by household location: wasting (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates. n = NA

Under 5 by household location: stunting (%)




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates. n = 4

Under 5 by household location: overweight (%)



Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates. n = 4

Under-5 coexistence of wasting, stunting and overweight




Sources: UNICEF, Division of Data Research and Policy (2018), Global UNICEF Global Databases: Overlapping Stunting, Wasting and Overweight, New York, May 2018.

Notes: Percentage of children under 5 years of age who experience different and overlapping forms of malnutrition. n = 6

Southern Asia


Child feeding practices



Sources: UNICEF, Division of Data Research and Policy (2018), Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2018. n = 6


Children and adolescent (aged 5–19) nutrition status

Ages 5–19 by gender: underweight (%)




Source: NCD Risk Factor Collaboration. n = 9

Ages 5–19 by gender: overweight (%)



Source: NCD Risk Factor Collaboration. n = 9


Ages 5–19 by gender: obesity (%)



Source: NCD Risk Factor Collaboration. n = 9


Adult nutrition status

Adult by gender: diabetes (%)




Source: NCD Risk Factor Collaboration. n = 9

Adult by gender: overweight (%)




Source: NCD Risk Factor Collaboration. n = 9

Adult by gender: obesity (%)




Source: NCD Risk Factor Collaboration. n = 9

Adult by gender: raised blood pressure (%)




Source: NCD Risk Factor Collaboration. n = 9

Adult: anaemia in WRA (%)



Source: WHO Global Health Observatory.
Note: WRA = women of reproductive age. n = 9


Adult: sodium intake (grams per day)



Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation. n = 9

Dietary needs


Consumption of food groups and components, 2016



Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation.
Notes: Men and women aged 25 and older. TMREL = theoretical minimum risk exposure level. n = 9

Financial resources and policy, legislation and institutional arrangements

Development assistance



Sources: Development Initiatives based on OECD Development Assistance Committee (DAC) Creditor Reporting System (CRS).
Notes: ODA = official development assistance. Amounts based on gross ODA per capita, constant 2016 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

National policies

| | |
|--|-----|
| Mandatory legislation for salt iodisation | 5/9 |
| Sugar-sweetened beverage tax | 2/9 |
| Food-based dietary guidelines | 6/9 |
| Multisectoral comprehensive nutrition plan | 7/9 |

Sources: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

Targets included in national (nutrition or other) plan

| | | | | | | | |
|----------|---------|------------------|------------------|-------------------------|---------|-------------|-----------------------------------|
| Stunting | Anaemia | Low birth weight | Child overweight | Exclusive breastfeeding | Wasting | Salt intake | Overweight adults and adolescents |
| 8/9 | 9/9 | 8/9 | 7/9 | 9/9 | 9/9 | 8/9 | 8/9 |

Sources: WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.