Western Europe

Subregional overview

Malnutrition burden

In the Western Europe subregion, there has been some progress towards achieving global nutrition targets. The global target for female diabetes has seven countries on course to meet it, while male diabetes has one country on course. However, not a single country in the subregion is on course to meet the targets for under-five overweight, under-five stunting, under-five wasting, infant exclusive breastfeeding, anaemia in women of reproductive age, low birth weight, male obesity, and female obesity. Nine countries in the subregion have insufficient data to comprehensively assess their progress towards these global targets.

The Western Europe subregion has no prevalence data available for under-five overweight, stunting, or wasting.

There is also insufficient data on exclusive breastfeeding among infants, while the subregion's average low birth weight prevalence of 6.9% is less than the global average of 14.6%.

The Western Europe subregion's adult population face a malnutrition burden. An average of 17% of women of reproductive age have anaemia, and 6.4% of adult men have diabetes, compared to 3.9% of women. Meanwhile, 22.9% of men and 20.4% of women have obesity.

Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF/WHO Low birthweight estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: Data on the adult indicators are based on modelled estimates.

Progress against global nutrition targets 2019



Under-five stunting

- 0 On course
- 0 Off course
- 9 No data



Low birthweight

- 0 On course
- 8 Off course
- 1 No data



Adult male obesity

- 0 On course
- 7 Off course
- 2 No data



- 0 On course
- 7 Off course



Under-five wasting

- 0 On course
- 0 Off course
- 9 No data



Exclusive breastfeeding

- 0 On course
- 0 Off course
- 9 No data



Adult female diabetes

- 7 On course
- 0 Off course
- 2 No data



Under-five overweight

- 0 On course
- 0 Off course
- 9 No data



Adult female obesity

- 0 On course
- 7 Off course
- 2 No data



Adult male diabetes

- 1 On course
- 6 Off course
- 2 No data



WRA anaemia

- 2 No data

Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory and Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: WRA = Women of a reproductive age; NA = not applicable. The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates.

Child (under-five) nutrition status

Coexistence of wasting, stunting and overweight

No data


Sources: UNICEF, Division of Data Research and Policy (2019).

UNICEF Global Databases: Overlapping Stunting, Wasting and

Overweight, January 2019, New York.

Notes: Percentage of children under-five years of age who experience different and overlapping forms of malnutrition.

Low birth weight



Source: UNICEF/WHO Low birthweight estimates, 2019 edition.

Notes: Based on population weighted means of 8 countries.

Child (under-five) nutrition status over time

Child (under-five) nutrition status over time				
Wasting by gender	Stunting by gender	Overweight by gender		
No data	No data	No data		
Wasting by location	Stunting by location	Overweight by location		
No data	No data	No data		
Wasting by income	Stunting by income	Overweight by income		
No data	No data	No data		

Wasting by mother's education	Stunting by mother's education	Overweight by mother's education
No data	No data	No data
Wasting by age	Stunting by age	Overweight by age
No data	No data	No data

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.

Notes: Disaggregated data (coloured lines/bars in charts) is based on population weighted means. Estimates are presented only where available data represents at least 50% of the regional population.

Infant and young child feeding over time

Exclusive Continued Minimum Intro. to solid, breastfeeding by breastfeeding at 1 semi-solid, soft acceptable diet by foods by gender gender year by gender gender No data No data No data No data Exclusive Continued Minimum Intro. to solid, breastfeeding by breastfeeding at 1 semi-solid, soft acceptable diet by foods by location year by location location location No data No data No data No data Exclusive Continued Minimum Intro. to solid, breastfeeding by breastfeeding at 1 acceptable diet by semi-solid, soft foods by income year by income income income No data No data No data No data

Exclusive	Continued breastfeeding at 1 year by mother's education	Minimum	Intro. to solid,
breastfeeding by		acceptable diet by	semi-solid, soft
mother's		mother's	foods by mother's
education		education	education
No data	No data	No data	No data
Exclusive	Continued breastfeeding at 1 year by age	Minimum	Intro. to solid,
breastfeeding by		acceptable diet by	semi-solid, soft
age		age	foods by age
No data	No data	No data	No data

Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2019.

Notes: Disaggregated data (coloured lines/bars in charts) is based on population weighted means. Estimates are presented only where available data represents at least 50% of the regional population.

No data No data No data


Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, Predominant breastfeeding, New York, May 2019.

Child and adolescent (aged 5-19) nutrition status

Underweight by gender


Overweight by gender

Obesity by gender




Sources: NCD Risk Factor Collaboration.

Notes: Based on population weighted means of 7 countries.



Notes: Based on population weighted means of 7 countries.



Notes: Based on population weighted means of 7 countries.

Adult nutrition status


Diabetes by gender

Male Female 5.9 4.1 3.9 2002 2006 2010 2014

Sources: NCD Risk Factor Collaboration.


Notes: Based on population weighted means of 7 countries.

Overweight by gender




Notes: Based on population weighted means of 7 countries.

Obesity by gender



Notes: Based on population weighted means of 7 countries.


Raised blood pressure by gender



Sources: NCD Risk Factor Collaboration.

Notes: Based on population weighted means of 7 countries.


Anaemia in WRA



Source: WHO Global Health Observatory.

Notes: WRA = women of reproductive age. Based on population weighted means of 7 countries.

Salt intake (grams per day)





Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: Based on population weighted means of 7 countries.

Dietary needs

Consumption of food groups and components, 2016





Sources: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: TMREL = theoretical minimum risk of exposure level. Men and women aged 25 and older. Based on population weighted means of 7 countries.


Intervention coverage

Coverage/practice indicator	Total (%)	Boy (%)	Girl (%)	Year
Children 0-59 months with diarrhoea who received zinc treatment	No	No	No	No
	data	data	data	data
Children 6-59 months who received vitamin A supplements in last 6 months	No	No	No	No
	data	data	data	data
Children 6-59 months given iron supplements in past 7 days	No	No	No	No
	data	data	data	data
Women with a live birth in the five years preceding the survey who received iron tablets or syrup during antenatal care	No data	NA	NA	No data
Household consumption of any iodised salt	No data	NA	NA	No data


Notes: NA = not applicable. Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005-2018.

Determinants

Undernourishment



Food supply




Notes: *0 = low inequality, 1 = high inequality.

Sources: ¹ UNICEF 2018; ² UNDP 2018.

Female secondary education enrolment (net,

% population)




Source: UNESCO Institute for Statistics

Notes: Based on population weighted means of between 6 and 7 countries.

Drinking water coverage (% population)

Source: FAOSTAT 2018.




Source: WHO/UNICEF Joint Monitoring Programme 2019.

Notes: Based on population weighted means of 9 countries.

Gender-related determinants

Early childbearing births by age 18 (%) ¹	NA	NA
Gender Inequality Index (score *) ²	NA	NA
Gender Inequality Index (country rank) ²	NA	NA

Sanitation coverage (% population)



Source: WHO/UNICEF Joint Monitoring Programme 2019.

Notes: Based on population weighted means of 9 countries.

Resources, policies and targets

Development assistance



Sources: Development Initiatives based on OECD Development Assistance Committee (DAC) Creditor Reporting System (CRS).

Notes: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2017 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

National policies

Mandatory legislation for salt iodisation	1/8
Sugar-sweetened beverage tax	3/8
Food-based dietary guidelines	6/8
Policy to reduce salt consumption	4/8
Operational policy, strategy or action plan to reduce unhealthy diet related to NCDs	8/8
Operational, multisectoral national NCD policy, strategy or action plan	4/8
Operational policy, strategy or action plan for diabetes	7/8
Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt	5/8
Policy to limit saturated fatty acids and virtually eliminate industrially produced trans-fats	6/8

Sources: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review, WHO Global Health Observatory.

Notes: Value refers to the number of countries with policy. NA = not applicable; NCD = non-communicable disease.

Targets included in national (nutrition or other) plan


Stunting	Anaemia
0/8	0/8
Low birth weight	Child overweight
1/8	6/8
Exclusive breastfeeding	Wasting
1/8	0/8
Salt intake	Overweight adults and adolescents
2/8	7/8
Multisectoral comprehensive nutrition plan	
3/8	

Sources: WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

Notes: Value refers to the number of countries with target.

Economics and demography

Poverty rates (%) and GDP (PPP\$)



Sources: World Bank 2019, IMF World Economic Outlook Database 2019.

Notes: PPP = purchasing power parity.

Based on population weighted means of
7 countries.

Under-five mortality (per 1,000 live births)



Source: UN Inter-agency Group for Child Mortality Estimation 2018.

Notes: Based on population weighted means of 8 countries.

Government revenues (\$m)



Sources: IMF Article IV staff reports (country specific) and IMF World Economic Outlook Database (April 2019).

Income inequality

Gini index score ¹	Gini index rank ²	Year
NA	NA	NA

Sources: World Bank 2019.

Notes: ¹ 0 = perfect equality, 100 = perfect inequality. ² Countries are ranked from most equal (1) to most unequal (159).

Population

Population (thousands)	196,616	2018
Under-five population (thousands)	10,084	2019
Rural (%)	20	2018
>65 years (thousands)	40,267	2019

Sources: World Bank 2019, UN Population Division Department of Economic and Social Affairs 2019.

Notes: Based on population weighted means of 9 countries.

Population density of health workers per 1,000 people

Physicians	3.78	2016
Nurses and midwives	12.2	2016
Community health workers	No data	No data

Sources: WHO's Global Health Workforce Statistics, OECD, supplemented by country data.

Notes: Based on population weighted means of 8 countries.