#### Myanmar

#### **Country overview**

#### Malnutrition burden

Myanmar is on course to meet the global targets for under-five overweight and infant exclusive breastfeeding, but is off course to meet the targets for all other indicators analysed with adequate data. There is insufficient target data to assess Myanmar's progress for male obesity.

Although it performs relatively well against other developing countries, Myanmar still experiences a malnutrition burden among its under-five population. As of 2016, the national prevalence of under-five overweight is 1.5%, which has decreased slightly from 2.6% in 2009. The national prevalence of under-five stunting is 29.4%, which is greater than the developing country average of 25%. Conversely, Myanmar's under-five wasting prevalence of 6.6% is less than the developing country average of 8.9%.

In Myanmar, 51.2% of infants under 6 months are exclusively breastfed, this is well above the South-eastern Asia average of 43.7%. Myanmar's 2015 low birth weight prevalence of 12.3% has decreased slightly from 13.9% in 2000.

Myanmar's adult population also face a malnutrition burden. 46.3% of women of reproductive age have anaemia, and 7.9% of adult women have diabetes, compared to 6.9% of men. Meanwhile, 7.3% of women and 4% of men have obesity.

Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF/WHO Low birthweight estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: Data on the adult indicators are based on modelled estimates.

#### Progress against global nutrition targets 2019

worsening




Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory and Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: WRA = Women of a reproductive age; NA = not applicable. The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates.

.

#### Child (under-five) nutrition status

## Coexistence of wasting, stunting and overweight



Sources: UNICEF, Division of Data Research and Policy (2019).
UNICEF Global Databases: Overlapping Stunting, Wasting and
Overweight, January 2019, New York.

Notes: Percentage of children under-five years of age who experience different and overlapping forms of malnutrition.

#### Low birth weight





Source: UNICEF/WHO Low birthweight estimates, 2019 edition.


#### Prevalence of under-five stunting

Stunting at subnational level

Stunting at 5km level







Source: Kinyoki, D.K. et al. Mapping child growth failure across low- and middle-income countries. Nature 577, 231–234 (2020) doi:10.1038/s41586-019-1878-8.


Notes: 5 km level map shows prevalence at the  $5 \times 5$ -km resolution. Prevalence is the 2017 estimated prevalence, based on a model using a range of surveys between 1998-2018. See source paper for full methods.

#### Child (under-five) nutrition status over time


Wasting by sex

Stunting by sex

Overweight by sex




29.4 27.5





Wasting by location

Stunting by location

Overweight by location









Wasting by income

Stunting by income

Overweight by income










## Wasting by mother's education

## Stunting by mother's education

## Overweight by mother's education









Wasting by age

Stunting by age

Overweight by age









Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.

#### Infant and young child feeding over time


Exclusive breastfeeding by sex




Continued breastfeeding at 1 year by sex



Minimum acceptable diet by sex




Intro. to solid, semi-solid, soft foods by sex





Exclusive breastfeeding by location





Minimum acceptable diet by location













Exclusive breastfeeding by income





Minimum acceptable diet by income

Intro. to solid, semi-solid, soft foods by income












# Exclusive breastfeeding by mother's education




#### Continued breastfeeding at 1 year by mother's education




Minimum acceptable diet by mother's education




Intro. to solid, semi-solid, soft foods by mother's education



Exclusive breastfeeding by age




Continued breastfeeding at 1 year by age



Minimum acceptable diet by age




Intro. to solid, semi-solid, soft foods by age



Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2019.

#### Infant and young child feeding




Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, Predominant breastfeeding, New York, May 2019.


#### Child and adolescent (aged 5-19) nutrition status


Underweight by sex

Overweight by sex

Obesity by sex







Sources: NCD Risk Factor Collaboration.


#### **Adult nutrition status**

#### Diabetes by sex


# Male Female 7,9 6,9 4,8

Sources: NCD Risk Factor Collaboration.


#### Overweight by sex



#### Obesity by sex




## Raised blood pressure by sex



Sources: NCD Risk Factor Collaboration.

#### Anaemia in WRA




Source: WHO Global Health

Observatory.

Notes: WRA = women of reproductive age.


## Sodium intake (grams per day)




Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

#### **Dietary needs**

#### Consumption of food groups and components, 2016





Sources: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: TMREL = theoretical minimum risk of exposure level. Men and women aged 25 and older.

#### Intervention coverage


| Coverage/practice indicator   | Total<br>(%) | Boy<br>(%) | Girl<br>(%) | Year |
|---|--------------|------------|-------------|------|
| Children 0-59 months with diarrhoea who received zinc treatment   | 8            | 6          | 11          | 2015 |
| Children 6-59 months who received vitamin A supplements in last 6 months  | 54           | 53         | 56          | 2015 |
| Children 6-59 months given iron supplements in past 7 days  | 8            | 8          | 9           | 2015 |
| Women with a live birth in the five years preceding the survey who received iron tablets or syrup during antenatal care | 87           | NA         | NA          | 2015 |
| Household consumption of any iodised salt   | 82           | NA         | NA          | 2015 |

Sources: Huestis A. and Kothari M., based on 2016 Global Nutrition Report.


Notes: NA = not applicable. Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005-2018.

#### **Determinants**

#### Undernourishment



#### Food supply




Source: FAOSTAT 2018.

## Gender-related determinants

| Early childbearing<br>births by age 18 (%) <sup>1</sup>       | 5    | 2016 |
|---|------|------|
| Gender Inequality<br>Index (score <sup>*</sup> ) <sup>2</sup> | 0.46 | 2017 |
| Gender Inequality Index (country rank) <sup>2</sup>           | 106  | 2017 |


Sources: <sup>1</sup> UNICEF 2018; <sup>2</sup> UNDP 2018. Notes: \*0 = low inequality, 1 = high inequality.

## Female secondary education enrolment (net, % population)




Source: UNESCO Institute for Statistics 2018.

## Drinking water coverage (% population)



Source: WHO/UNICEF Joint Monitoring Programme 2019.

## Sanitation coverage (% population)




Source: WHO/UNICEF Joint Monitoring Programme 2019.

#### Resources, policies and targets

#### Development assistance

- Basic nutrition ODA received
- % of total ODA



Sources: Development Initiatives based on OECD Development Assistance Committee (DAC) Creditor Reporting System (CRS).

Notes: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2017 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

#### National policies

| Mandatory legislation for salt iodisation  | Yes        |
|--|------------|
| Sugar-sweetened beverage tax   | No         |
| Food-based dietary guidelines  | No<br>data |
| Policy to reduce salt consumption  | No         |
| Operational policy, strategy or action plan to reduce unhealthy diet related to NCDs   | No         |
| Operational, multisectoral national NCD policy, strategy or action plan  | No         |
| Operational policy, strategy or action plan for diabetes   | No         |
| Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt | No         |
| Policy to limit saturated fatty acids and virtually eliminate industrially produced trans-fats   | No         |

Sources: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review, WHO Global Health Observatory.

Notes: NA = not applicable; NCD = non-communicable disease.


#### Targets included in national (nutrition or other) plan

| Stunting                                   | Anaemia                           |
|--|-----------------------------------|
| Yes  | Yes                               |
| Low birth weight                           | Child overweight                  |
| Yes  | Yes                               |
| Exclusive breastfeeding                    | Wasting                           |
| Yes  | Yes                               |
| Salt intake                                | Overweight adults and adolescents |
| Yes  | Yes                               |
| Multisectoral comprehensive nutrition plan |                                   |
| Yes  |                                   |

Sources: WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

#### **Economics and demography**


## Poverty rates (%) and GDP (PPP\$)



Sources: World Bank 2019, IMF World Economic Outlook Database 2019.


Notes: PPP = purchasing power parity.

## Under-five mortality (per 1,000 live births)



Source: UN Inter-agency Group for Child Mortality Estimation 2018.

## Government revenues (\$m)



Sources: IMF Article IV staff reports (country specific) and IMF World Economic Outlook Database (April 2019).

#### Income inequality

| Gini index<br>score <sup>1</sup> | Gini index<br>rank <sup>2</sup> | Year |
|----------------------------------|---------------------------------|------|
| 38                               | 91                              | 2015 |

Sources: World Bank 2019.

Notes: <sup>1</sup> 0 = perfect equality, 100 = perfect inequality. <sup>2</sup> Countries are ranked from most equal (1) to most unequal (159).

#### Population

| Population<br>(thousands)               | 53,708 | 2018 |
|---|--------|------|
| Under-five<br>population<br>(thousands) | 4,511  | 2019 |
| Rural (%)                               | 69     | 2018 |
| >65 years<br>(thousands)                | 3,249  | 2019 |

Sources: World Bank 2019, UN Population Division Department of Economic and Social Affairs 2019.

#### Population density of health workers per 1,000 people

| Physicians               | 0.57 | 2012 |
|--------------------------|------|------|
| Nurses and midwives      | 0.93 | 2012 |
| Community health workers | 0.2  | 2012 |

Sources: WHO's Global Health Workforce Statistics, OECD, supplemented by country data.