

## Subregional overview

### Malnutrition burden

In the Northern America subregion, there has been some progress towards achieving global nutrition targets. The global targets for under-five stunting, under-five wasting, and female diabetes each have one country on course to meet them. However, not a single country in the subregion is on course to meet the targets for under-five overweight, infant exclusive breastfeeding, anaemia in women of reproductive age, low birth weight, male diabetes, male obesity, and female obesity. One country in the subregion has insufficient data to comprehensively assess its progress towards these global targets.

The Northern America subregion experiences a malnutrition burden among its under-five population. The average prevalence of overweight in under-fives is 8.8%, this is greater than the global average of 5.9%. Conversely, the Northern America subregion's prevalence of stunting in under-fives of 2.6% is significantly less than the global average of 21.9%. The prevalence of wasting in under-fives is 0.4%.


Some 34.7% of infants under 6 months in the Northern America subregion are exclusively breastfed, while the subregion's average low birth weight prevalence of 7.9% is less than the global average of 14.6%.


The Northern America subregion's adult population also face a malnutrition burden. An average of 12.9% of women of reproductive age have anaemia, and 8% of adult men have diabetes, compared to 6.3% of women. Meanwhile, 36.2% of women and 34.9% of men have obesity.

Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF/WHO Low birthweight estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: Data on the adult indicators are based on modelled estimates.

### Progress against global nutrition targets 2018






Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory and Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: WRA = Women of a reproductive age; NA = not applicable. The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates.

## Child (under-five) nutrition status


### Coexistence of wasting, stunting and overweight



Sources: UNICEF, Division of Data Research and Policy (2019). UNICEF Global Databases: Overlapping Stunting, Wasting and Overweight, January 2019, New York.

Notes: Percentage of children under-five years of age who experience different and overlapping forms of malnutrition.


### Low birth weight




Source: UNICEF/WHO Low birthweight estimates, 2019 edition.

# Child (under-five) nutrition status over time


Wasting by gender




Stunting by gender




Overweight by gender




Wasting by location




Stunting by location




Overweight by location




Wasting by income




Stunting by income




Overweight by income




## Wasting by mother's education




## Stunting by mother's education




## Overweight by mother's education




## Wasting by age



## Stunting by age



## Overweight by age




Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.


Notes: Regional trends (grey line in charts) refer to estimates from UNICEF/WHO/World Bank Group: Joint child malnutrition estimates. Stunting and wasting based only on United States data. Disaggregated data (coloured lines/bars in charts) is based on population weighted means. Estimates are presented only where available data represents at least 50% of the regional population.

# Infant and young child feeding over time

Exclusive breastfeeding by gender



Continued breastfeeding at 1 year by gender




Minimum acceptable diet by gender




Intro. to solid, semi-solid, soft foods by gender



Exclusive breastfeeding by location



Continued breastfeeding at 1 year by location




Minimum acceptable diet by location




Intro. to solid, semi-solid, soft foods by location



Exclusive breastfeeding by income



Continued breastfeeding at 1 year by income



Minimum acceptable diet by income



Intro. to solid, semi-solid, soft foods by income






Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2019.

Notes: Regional trends (grey line in charts) for exclusive breastfeeding and continued breastfeeding at 1 year refer to estimates from UNICEF/WHO/World Bank Group: Joint child malnutrition estimates. Disaggregated data (coloured lines/bars in charts) is based on population weighted means. Estimates are presented only where available data represents at least 50% of the regional population.


## Infant and young child feeding



Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, Predominant breastfeeding, New York, May 2019.


# Child and adolescent (aged 5-19) nutrition status

## Underweight by gender




Sources: NCD Risk Factor Collaboration.  
Notes: Based on population weighted means of 2 countries.

## Overweight by gender



Notes: Based on population weighted means of 2 countries.


## Obesity by gender



Notes: Based on population weighted means of 2 countries.


# Adult nutrition status

## Diabetes by gender




Sources: NCD Risk Factor Collaboration.  
Notes: Based on population weighted means of 2 countries.

## Overweight by gender




Notes: Based on population weighted means of 2 countries.

## Obesity by gender




Notes: Based on population weighted means of 2 countries.

## Raised blood pressure by gender




Sources: NCD Risk Factor Collaboration.  
Notes: Based on population weighted means of 2 countries.

## Anaemia in WRA



Source: WHO Global Health Observatory.  
Notes: WRA = women of reproductive age. Based on population weighted means of 2 countries.

## Salt intake (grams per day)



Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation.  
Notes: Based on population weighted means of 2 countries.



# Dietary needs

## Consumption of food groups and components, 2016



Sources: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: TMREL = theoretical minimum risk of exposure level. Men and women aged 25 and older. Based on population weighted means of 2 countries.


# Intervention coverage

Coverage/practice indicator	Total (%)	Boy (%)	Girl (%)	Year
Children 0-59 months with diarrhoea who received zinc treatment	No data	No data	No data	No data
Children 6-59 months who received vitamin A supplements in last 6 months	No data	No data	No data	No data
Children 6-59 months given iron supplements in past 7 days	No data	No data	No data	No data
Women with a live birth in the five years preceding the survey who received iron tablets or syrup during antenatal care	No data	NA	NA	No data
Household consumption of any iodised salt	No data	NA	NA	No data

Notes: NA = not applicable. Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005-2018.


# Determinants

## Undernourishment



Source: FAOSTAT 2018.

## Food supply



Source: FAOSTAT 2018.


## Gender-related determinants

Early childbearing births by age 18 (%) <sup>1</sup>	NA	NA
Gender Inequality Index (score <sup>*</sup> ) <sup>2</sup>	NA	NA
Gender Inequality Index (country rank) <sup>2</sup>	NA	NA

Sources: <sup>1</sup> UNICEF 2018; <sup>2</sup> UNDP 2018.

Notes: <sup>\*</sup> 0 = low inequality, 1 = high inequality.


## Female secondary education enrolment (net, % population)



Source: UNESCO Institute for Statistics 2018.

Notes: Based on population weighted means of between 1 and 2 countries.


## Drinking water coverage (% population)



Source: WHO/UNICEF Joint Monitoring Programme 2019.

Notes: Based on population weighted means of 2 countries.

## Sanitation coverage (% population)



Source: WHO/UNICEF Joint Monitoring Programme 2019.

Notes: Based on population weighted means of 2 countries.

# Resources, policies and targets

## Development assistance



Sources: Development Initiatives based on OECD Development Assistance Committee (DAC) Creditor Reporting System (CRS).

Notes: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2017 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

## National policies

Mandatory legislation for salt iodisation	1/2
Sugar-sweetened beverage tax	0/2
Food-based dietary guidelines	2/2
Policy to reduce salt consumption	2/2
Operational policy, strategy or action plan to reduce unhealthy diet related to NCDs	2/2
Operational, multisectoral national NCD policy, strategy or action plan	2/2
Operational policy, strategy or action plan for diabetes	2/2
Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt	2/2
Policy to limit saturated fatty acids and virtually eliminate industrially produced trans-fats	2/2

Sources: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review, WHO Global Health Observatory.

Notes: Value refers to the number of countries with policy. NA = not applicable; NCD = non-communicable disease.

Targets included in national (nutrition or other) plan


Stunting	Anaemia
0/2	1/2
Low birth weight	Child overweight
1/2	2/2
Exclusive breastfeeding	Wasting
2/2	0/2
Salt intake	Overweight adults and adolescents
2/2	2/2
Multisectoral comprehensive nutrition plan	
0/2	

Sources: WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

Notes: Value refers to the number of countries with target.

# Economics and demography


## Poverty rates (%) and GDP (PPP\$)



Sources: World Bank 2019, IMF World Economic Outlook Database 2019.

Notes: PPP = purchasing power parity. Based on population weighted means of 2 countries.

## Under-five mortality (per 1,000 live births)



Source: UN Inter-agency Group for Child Mortality Estimation 2018.

Notes: Based on population weighted means of 2 countries.

## Government revenues (\$m)



Sources: IMF Article IV staff reports (country specific) and IMF World Economic Outlook Database (April 2019).

## Income inequality

Gini index score <sup>1</sup>	Gini index rank <sup>2</sup>	Year
NA	NA	NA

Sources: World Bank 2019.

Notes: <sup>1</sup> 0 = perfect equality, 100 = perfect inequality.<sup>2</sup> Countries are ranked from most equal (1) to most unequal (159).

## Population

Population (thousands)	364,226	2018
Under-five population (thousands)	21,590	2019
Rural (%)	18	2018
>65 years (thousands)	59,962	2019

Sources: World Bank 2019, UN Population Division Department of Economic and Social Affairs 2019.

Notes: Based on population weighted means of 2 countries.

## Population density of health workers per 1,000 people

Physicians	2.57	2016
Nurses and midwives	9.88	2016
Community health workers	No data	No data

Sources: WHO's Global Health Workforce Statistics, OECD, supplemented by country data.

Notes: Based on population weighted means of 2 countries.