### **United States of America**

### **Country overview**

#### Malnutrition burden

The United States is on course to meet the global targets for under-five stunting and under-five wasting, but is off course to meet the targets for all other indicators analysed with adequate data.

Although it performs relatively well against other countries, the United States still experiences a malnutrition burden among its under-five population. As of 2016, the national prevalence of under-five overweight is 9.4%, which has increased slightly from 8.2% in 2014. The national prevalence of under-five stunting is 3.5%, which is significantly less than the global average of 21.9%. the United States' under-five wasting prevalence of 0.4% is also less than the global average of 7.3%.

In the United States, 34.7% of infants under 6 months are exclusively breastfed. The United States' 2015 low birth weight prevalence of 8% has increased slightly from 7.5% in 2000.

The United States' adult population also face a malnutrition burden. 13.3% of women of reproductive age have anaemia, and 8.2% of adult men have diabetes, compared to 6.4% of women. Meanwhile, 37% of women and 35.5% of men have obesity.

Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF/WHO Low birthweight estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: Data on the adult indicators are based on modelled estimates.

#### Progress against global nutrition targets 2018

No progress or

worsening



Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory and Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: WRA = Women of a reproductive age; NA = not applicable. The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates.

### Child (under-five) nutrition status

### Coexistence of wasting, stunting and overweight




Sources: UNICEF, Division of Data Research and Policy (2019).

UNICEF Global Databases: Overlapping Stunting, Wasting and

Overweight, January 2019, New York.

Notes: Percentage of children under-five years of age who experience different and overlapping forms of malnutrition.

### Low birth weight




Source: UNICEF/WHO Low birthweight estimates, 2019 edition.


### Child (under-five) nutrition status over time


### Wasting by gender

#### Stunting by gender

### Overweight by gender










Wasting by location

Stunting by location

Overweight by location










Wasting by income

Stunting by income

Overweight by income










### Wasting by mother's education

### Stunting by mother's education

### Overweight by mother's education










Wasting by age

Stunting by age

Overweight by age







Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.

### Infant and young child feeding over time


**Exclusive** breastfeeding by gender



Continued breastfeeding at 1 year by gender

Minimum acceptable diet by gender

Intro. to solid, semi-solid, soft foods by gender



No data




**Exclusive** breastfeeding by location

Continued breastfeeding at 1 year by location

Minimum acceptable diet by location

Intro. to solid, semi-solid, soft foods by location









**Exclusive** breastfeeding by income



Minimum acceptable diet by income

Intro. to solid, semi-solid, soft foods by income












### **Exclusive** breastfeeding by mother's education



### Continued breastfeeding at 1 year by mother's education



### Minimum acceptable diet by mother's education



### Intro. to solid, semi-solid, soft foods by mother's education




### **Exclusive** breastfeeding by age




breastfeeding at 1

Continued






Intro. to solid, semi-solid, soft foods by age








Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2019.

### Infant and young child feeding



Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, Predominant breastfeeding, New York, May 2019.


### Child and adolescent (aged 5-19) nutrition status


Underweight by gender

Overweight by gender

Obesity by gender







Sources: NCD Risk Factor Collaboration.


#### **Adult nutrition status**

### Diabetes by gender


# Male Female % and a second point of the secon

Sources: NCD Risk Factor Collaboration.


### Overweight by gender



### Obesity by gender




### Raised blood pressure by gender



Sources: NCD Risk Factor Collaboration.


### Anaemia in WRA



Source: WHO Global Health Observatory.

Notes: WRA = women of reproductive age.


### Salt intake (grams per day)




Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

### **Dietary needs**

#### Consumption of food groups and components, 2016





Sources: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: TMREL = theoretical minimum risk of exposure level. Men and women aged 25 and older.

### Intervention coverage

Coverage/practice indicator	Total (%)	Boy (%)	Girl (%)	Year
Children 0-59 months with diarrhoea who received zinc treatment	No	No	No	No
	data	data	data	data
Children 6-59 months who received vitamin A supplements in last 6 months	No	No	No	No
	data	data	data	data
Children 6-59 months given iron supplements in past 7 days	No	No	No	No
	data	data	data	data
Women with a live birth in the five years preceding the survey who received iron tablets or syrup during antenatal care	No data	NA	NA	No data
Household consumption of any iodised salt	No data	NA	NA	No data

Sources: Huestis A. and Kothari M., based on 2016 Global Nutrition Report.


Notes: NA = not applicable. Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005-2018.

#### **Determinants**

#### Undernourishment



#### Food supply




Source: FAOSTAT 2018.

### Gender-related determinants

Early childbearing births by age 18 (%) <sup>1</sup>	No data	No data
Gender Inequality Index (score <sup>*</sup> ) <sup>2</sup>	0.19	2017
Gender Inequality Index (country rank) <sup>2</sup>	41	2017


Sources: <sup>1</sup> UNICEF 2018; <sup>2</sup> UNDP 2018. Notes: \*0 = low inequality, 1 = high inequality.

## Female secondary education enrolment (net, % population)




Source: UNESCO Institute for Statistics 2018.

### Drinking water coverage (% population)



Source: WHO/UNICEF Joint Monitoring Programme 2019.

### Sanitation coverage (% population)




Source: WHO/UNICEF Joint Monitoring Programme 2019.

### Resources, policies and targets

### Development assistance

- % of total ODA
- Basic nutrition ODA disbursed



Sources: Development Initiatives based on OECD Development Assistance Committee (DAC) Creditor Reporting System (CRS).

Notes: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2017 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

### National policies

Mandatory legislation for salt iodisation	No
Sugar-sweetened beverage tax	No
Food-based dietary guidelines	Yes
Policy to reduce salt consumption	Yes
Operational policy, strategy or action plan to reduce unhealthy diet related to NCDs	Yes
Operational, multisectoral national NCD policy, strategy or action plan	Yes
Operational policy, strategy or action plan for diabetes	Yes
Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt	Yes
Policy to limit saturated fatty acids and virtually eliminate industrially produced trans-fats	Yes

Sources: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review, WHO Global Health Observatory.

Notes: NA = not applicable; NCD = non-communicable disease.


### Targets included in national (nutrition or other) plan

Stunting	Anaemia
No	Yes
Low birth weight	Child overweight
Yes	Yes
Exclusive breastfeeding	Wasting
Yes	No
Salt intake	Overweight adults and adolescents
Yes	Yes
Multisectoral comprehensive nutrition plan	
No	

Sources: WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

### **Economics and demography**


### Poverty rates (%) and GDP (PPP\$)



Sources: World Bank 2019, IMF World Economic Outlook Database 2019.

Notes: PPP = purchasing power parity.

### Under-five mortality (per 1,000 live births)



Source: UN Inter-agency Group for Child Mortality Estimation 2018.

### Government revenues (\$m)



Sources: IMF Article IV staff reports (country specific) and IMF World Economic Outlook Database (April 2019).

#### Income inequality

Gini index score <sup>1</sup>	Gini index rank <sup>2</sup>	Year
42	111	2016

Sources: World Bank 2019.

Notes: <sup>1</sup> 0 = perfect equality, 100 = perfect inequality. <sup>2</sup> Countries are ranked from most equal (1) to most unequal (159).

#### **Population**

Population (thousands)	327,167	2018
Under-five population (thousands)	19,604	2019
Rural (%)	18	2018
>65 years (thousands)	53,340	2019

Sources: World Bank 2019, UN Population Division Department of Economic and Social Affairs 2019.

### Population density of health workers per 1,000 people

Physicians	2.57	2014
Nurses and midwives	9.88	2005
Community health workers	No data	No data

Sources: WHO's Global Health Workforce Statistics, OECD, supplemented by country data.