Guinea

Country overview

Malnutrition burden

Guinea is on course to meet the global target for infant exclusive breastfeeding, but is off course to meet the targets for all other indicators analysed with adequate data. There is insufficient target data to assess Guinea's progress for low birth weight.

Although it performs relatively well against other developing countries, Guinea still experiences a malnutrition burden among its under-five population. As of 2016, the national prevalence of under-five overweight is 4%, which has increased slightly from 3.8% in 2012. The national prevalence of under-five stunting is 32.4%, which is greater than the developing country average of 25%. Conversely, Guinea's under-five wasting prevalence of 8.1% is less than the developing country average of 8.9%.

In Guinea, 34.2% of infants under 6 months are exclusively breastfed. There is insufficient data on low birth weight.

Guinea's adult population also face a malnutrition burden. 50.6% of women of reproductive age have anaemia, and 6.8% of adult men have diabetes, compared to 6.1% of women. Meanwhile, 11.5% of women and 3.7% of men have obesity.

Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF/WHO Low birthweight estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: Data on the adult indicators are based on modelled estimates.

Progress against global nutrition targets 2019

No progress or

worsening




Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory and Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: WRA = Women of a reproductive age; NA = not applicable. The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates.

Child (under-five) nutrition status

Coexistence of wasting, stunting and overweight



Sources: UNICEF, Division of Data Research and Policy (2019).

UNICEF Global Databases: Overlapping Stunting, Wasting and

Overweight, January 2019, New York.

Notes: Percentage of children under-five years of age who experience different and overlapping forms of malnutrition.

Low birth weight




Source: UNICEF/WHO Low birthweight estimates, 2019 edition.

Child (under-five) nutrition status over time


Wasting by gender

Stunting by gender

Overweight by gender




8 39.2 39.2 34.5 32.4 30.3




Wasting by location

Stunting by location

Overweight by location







Wasting by income

Stunting by income

Overweight by income









Wasting by mother's education

Stunting by mother's education

Overweight by mother's education







Wasting by age

Stunting by age

Overweight by age








Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.


Infant and young child feeding over time


Exclusive breastfeeding by gender





Continued breastfeeding at 1 year by gender













Exclusive breastfeeding by location


Continued breastfeeding at 1 year by location


Minimum acceptable diet by location

Intro. to solid, semi-solid, soft foods by location











Exclusive breastfeeding by income


Continued breastfeeding at 1 year by income


Minimum acceptable diet by income

Intro. to solid, semi-solid, soft foods by income











Exclusive breastfeeding by mother's education



Continued breastfeeding at 1 year by mother's education




breastfeeding at 1


Continued

year by age

Minimum acceptable diet by mother's education




Intro. to solid, semi-solid, soft foods by mother's education



Intro. to solid,

semi-solid, soft

Exclusive breastfeeding by age



Minimum acceptable diet by age





Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding, New York, May 2019.

Infant and young child feeding




Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, Predominant breastfeeding, New York, May 2019.


Child and adolescent (aged 5-19) nutrition status


Underweight by gender

Overweight by gender

Obesity by gender







Sources: NCD Risk Factor Collaboration.

Adult nutrition status

Diabetes by gender

Male Female 6,8 6,1

Sources: NCD Risk Factor Collaboration.

Overweight by gender




5,7 11,5

2008

2016


Obesity by gender

Raised blood pressure by gender



Sources: NCD Risk Factor Collaboration.


Anaemia in WRA



Source: WHO Global Health Observatory.

Notes: WRA = women of reproductive age.


Salt intake (grams per day)




Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Dietary needs

Consumption of food groups and components, 2016





Sources: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: TMREL = theoretical minimum risk of exposure level. Men and women aged 25 and older.

Intervention coverage

| Coverage/practice indicator | Total (%) | Boy (%) | Girl (%) | Year |
|---|--------------|------------|-------------|------|
| Children 0-59 months with diarrhoea who received zinc treatment | 0 | 0 | 1 | 2012 |
| Children 6-59 months who received vitamin A supplements in last 6 months | 41 | 42 | 40 | 2012 |
| Children 6-59 months given iron supplements in past 7 days | 12 | 12 | 11 | 2012 |
| Women with a live birth in the five years preceding the survey who received iron tablets or syrup during antenatal care | 81 | NA | NA | 2012 |
| Household consumption of any iodised salt | 64 | NA | NA | 2012 |

Sources: Huestis A. and Kothari M., based on 2016 Global Nutrition Report.


Notes: NA = not applicable. Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005-2018.

Determinants

Undernourishment



Food supply



Source: FAOSTAT 2018.


Gender-related determinants

| Early childbearing births by age 18 (%) ¹ | 40 | 2012 |
|---|------------|------------|
| Gender Inequality Index (score [*]) ² | No data | No data |
| Gender Inequality Index (country rank) ² | No data | No data |

Sources: 1 UNICEF 2018; 2 UNDP 2018.


Notes: *0 = low inequality, 1 = high inequality.

Female secondary education enrolment (net, % population)



Source: UNESCO Institute for Statistics 2018.

Drinking water coverage (% population)



Source: WHO/UNICEF Joint Monitoring Programme 2019.

Sanitation coverage (% population)




Source: WHO/UNICEF Joint Monitoring Programme 2019.

Resources, policies and targets

Development assistance

- Basic nutrition ODA received
- % of total ODA



Sources: Development Initiatives based on OECD Development Assistance Committee (DAC) Creditor Reporting System (CRS).

Notes: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2017 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

National policies

| Mandatory legislation for salt iodisation | Yes |
|--|------------|
| Sugar-sweetened beverage tax | No |
| Food-based dietary guidelines | No data |
| Policy to reduce salt consumption | No |
| Operational policy, strategy or action plan to reduce unhealthy diet related to NCDs | No |
| Operational, multisectoral national NCD policy, strategy or action plan | No |
| Operational policy, strategy or action plan for diabetes | Yes |
| Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt | No |
| Policy to limit saturated fatty acids and virtually eliminate industrially produced trans-fats | No |

Sources: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review, WHO Global Health Observatory.

Notes: NA = not applicable; NCD = non-communicable disease.


Targets included in national (nutrition or other) plan

| Stunting | Anaemia |
|--|-----------------------------------|
| Yes | Yes |
| Low birth weight | Child overweight |
| Yes | Yes |
| Exclusive breastfeeding | Wasting |
| Yes | Yes |
| Salt intake | Overweight adults and adolescents |
| Yes | Yes |
| Multisectoral comprehensive nutrition plan | |
| Yes | |

Sources: WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

Economics and demography


Poverty rates (%) and GDP (PPP\$)



Sources: World Bank 2019, IMF World Economic Outlook Database 2019.


Notes: PPP = purchasing power parity.

Under-five mortality (per 1,000 live births)



Source: UN Inter-agency Group for Child Mortality Estimation 2018.

Government revenues (\$m)



Sources: IMF Article IV staff reports (country specific) and IMF World Economic Outlook Database (April 2019).

Income inequality

| Gini index score ¹ | Gini index rank ² | Year |
|----------------------------------|---------------------------------|------|
| 34 | 49 | 2012 |

Sources: World Bank 2019.

Notes: ¹ 0 = perfect equality, 100 = perfect inequality. ² Countries are ranked from most equal (1) to most unequal (159).

Population

| Population (thousands) | 12,414 | 2018 |
|---|--------|------|
| Under-five population (thousands) | 2,060 | 2019 |
| Rural (%) | 64 | 2018 |
| >65 years (thousands) | 376 | 2019 |

Sources: World Bank 2019, UN Population Division Department of Economic and Social Affairs 2019.

Population density of health workers per 1,000 people

| Physicians | 0.07 | 2016 |
|--------------------------|------|------|
| Nurses and midwives | 0.37 | 2016 |
| Community health workers | 0.52 | 2016 |

Sources: WHO's Global Health Workforce Statistics, OECD, supplemented by country data.