Lao People's Democratic Republic

Overview

Malnutrition status

Lao PDR is off course to meet the global targets for infant exclusive breastfeeding, anaemia in women of reproductive age, low birth weight, male diabetes, and female diabetes. There is insufficient target data to assess Lao PDR's progress for under-five overweight, under-five stunting, under-five wasting, male obesity, and female obesity.

Although it performs relatively well against other developing countries, Lao PDR still experiences a malnutrition burden among its under-five population. As of 2011, the national prevalence of under-five overweight is 2%, which has increased slightly from 1.3% in 2006. The national prevalence of under-five stunting is 44.2%, which is significantly greater than the developing country average of 25%. Conversely, Lao PDR's underfive wasting prevalence of 6.4% is less than the developing country average of 8.9%.

In Lao PDR, 44.4% of infants under 23 months are exclusively breastfed, this is well above the South-eastern Asia average of 38.8%. Lao PDR's 2015 low birth weight prevalence of 17.3% has decreased slightly from 17.4% in 2014.


Lao PDR's adult population also face a malnutrition burden. While there is no prevalence data available for anaemia among women of reproductive age, 7.7% of adult men suffer from diabetes, compared to 7.6% of women. Meanwhile, 6.7% of women and 3.7% of men suffer from obesity.

Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, UNICEF/WHO Low birthweight estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory.

Notes: Data on the adult indicators are based on modelled estimates.

Progress against global nutrition targets 2018

worsening




Sources: UNICEF global databases Infant and Young Child Feeding, UNICEF/WHO/World Bank Group: Joint child malnutrition estimates, NCD Risk Factor Collaboration, WHO Global Health Observatory and Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: WRA = Women of a reproductive age; NA = not applicable. The methodologies for tracking differ between targets. Data on the adult indicators are based on modelled estimates.

Infant and child (under-five) status


Under-five coexistence of wasting, stunting and overweight



Sources: UNICEF, Division of Data Research and Policy (2019). UNICEF Global Databases: Overlapping Stunting, Wasting and Overweight, January 2019, New York.

Notes: Percentage of children under-five years of age who experience different and overlapping forms of malnutrition.

Low birth weight




Source: UNICEF/WHO Low birthweight estimates, 2019 edition.

Children (under-five) nutrition status


Wasting by gender

Stunting by gender


Overweight by gender




Wasting by location




Stunting by location




Overweight by location




Wasting by income





Stunting by income





Overweight by income

















Wasting by age

Stunting by age

Overweight by age









Sources: UNICEF/WHO/World Bank Group: Joint child malnutrition estimates.

Infant and child (under-five) feeding over time


Exclusive breastfeeding by gender




Continued breastfeeding at 1 year by gender




Minimum acceptable diet by gender




Introduction to solid, semi-solid or soft foods by gender




Exclusive breastfeeding by location



Continued breastfeeding at 1 year by location



Minimum acceptable diet by location




Introduction to solid, semi-solid or soft foods by location




Exclusive breastfeeding by income




Continued breastfeeding at 1 year by income




Minimum acceptable diet by income




Introduction to solid, semi-solid or soft foods by income




Exclusive breastfeeding by mother's education




Continued breastfeeding at 1 year by mother's education




Minimum acceptable diet by mother's education



Introduction to solid, semi-solid or soft foods by mother's education




Exclusive breastfeeding by age




Continued breastfeeding at 1 year by age



Minimum acceptable diet by age




Introduction to solid, semi-solid or soft foods by age



Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, Predominant breastfeeding, New York, May 2019.

Infant and young child feeding



Sources: UNICEF, Division of Data Research and Policy (2019). Global UNICEF Global Databases: Infant and Young Child Feeding: Exclusive breastfeeding, Predominant breastfeeding, New York, May 2019.


Children and adolescent (aged 5-19) nutrition status


Ages 5-19 by gender: underweight (%)

Ages 5-19 by gender: overweight (%)

Ages 5-19 by gender: obesity (%)







Sources: NCD Risk Factor


Collaboration.


Adult nutrition status


Adult by gender: diabetes (%)


Adult by gender: overweight (%)

Adult by gender: obesity (%)










Collaboration.


Adult by gender: raised blood pressure (%)

Adult: anaemia in WRA (%)

Adult: sodium intake (grams per day)




Sources: NCD Risk Factor Collaboration.



Source: WHO Global Health Observatory.


Notes: WRA = women of reproductive




Source: Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Dietary needs

Consumption of food groups and components, 2016





Sources: TMREL = theoretical minimum risk of exposure level. Global Burden of Disease, the Institute for Health Metrics and Evaluation.

Notes: Men and women aged 25 and older.

Intervention coverage

Coverage/practice indicator	Total (%)	Boy (%)	Girl (%)	Year
Children 0-59 months with diarrhoea who received zinc treatment	NA	NA	NA	NA

Coverage/practice indicator	Total (%)	Boy (%)	Girl (%)	Year
Children 6-59 months who received A supplements in last 6 months	NA	NA	NA	NA
Children 6-59 months given iron supplements in past 7 days	NA	NA	NA	NA
Women with a live birth in the five years preceding the survey who received iron tablets or syrup during antenatal care	NA			NA
Household consumption of any iodised salt	NA	NA	NA	NA

Sources: Huestis A. and Kothari M., based on 2016 Global Nutrition Report and UNICEF global databases, 2019.

Notes: NA = not applicable. Data is compiled using STATcompiler and taken from country Demographic and Health Surveys for 2005-2018.

Determinants

Undernourishment (% population)

Food supply

Gender-related determinants






Early childbearing: births by age 18 (%) ¹	18	2012
Gender Inequality Index (score*) ²	0.46	2017
Gender Inequality Index (country rank) ²	110	2017

Sources: ¹ UNICEF 2018; ² UNDP 2018. Notes: *0 = low inequality, 1 = high inequality.


Female secondary education enrolment (net, % population)

Drinking water coverage (% population)


Sanitation coverage (% population)



Source: UNESCO Institute for Statistics 2018.



Source: WHO/UNICEF Joint Monitoring Programme 2019.




Source: WHO/UNICEF Joint Monitoring Programme 2019.

Resources, policies and targets

Development assistance

- Basic nutrition ODA received
- % of total ODA



Sources: Development Initiatives based on OECD Development Assistance Committee (DAC) Creditor Reporting System (CRS).

Notes: ODA = official development assistance. Amounts based on gross ODA disbursements, constant 2017 prices. Figure includes ODA grants and loans, but excludes other official flows and private grants.

National policies

Sugar-sweetened beverage tax Food-based dietary guidelines Policy to reduce salt consumption Operational policy, strategy or action plan to reduce unhealthy diet related to NCDs Operational, multisectoral national NCD policy, strategy or action plan Operational policy, strategy or action plan for diabetes Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt		
Food-based dietary guidelines Policy to reduce salt consumption Operational policy, strategy or action plan to reduce unhealthy diet related to NCDs Operational, multisectoral national NCD policy, strategy or action plan Operational policy, strategy or action plan for diabetes Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt	ndatory legislation for salt iodisation	Yes
Policy to reduce salt consumption Operational policy, strategy or action plan to reduce unhealthy diet related to NCDs Operational, multisectoral national NCD policy, strategy or action plan Operational policy, strategy or action plan for diabetes Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt	ar-sweetened beverage tax	No
Operational policy, strategy or action plan to reduce unhealthy diet related to NCDs Operational, multisectoral national NCD policy, strategy or action plan Operational policy, strategy or action plan for diabetes Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt	d-based dietary guidelines	NA
Operational, multisectoral national NCD policy, strategy or action plan Operational policy, strategy or action plan for diabetes Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt	cy to reduce salt consumption	No
Operational policy, strategy or action plan for diabetes Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt	erational policy, strategy or action plan to reduce unhealthy diet related to NCDs	Yes
Policy to reduce the impact on children of marketing of foods and beverages high in saturated fats, trans-fatty acids, free sugars or salt	erational, multisectoral national NCD policy, strategy or action plan	Yes
free sugars or salt	erational policy, strategy or action plan for diabetes	Yes
Policy to limit saturated fatty acids and virtually eliminate industrially produced trans-fats		No
	icy to limit saturated fatty acids and virtually eliminate industrially produced trans-fats	No

Sources: Global Fortification Data Exchange 2018; Sugar-sweetened data prepared using data from the NOURISHING database, academic references and market reports; FAO 2018; WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review, WHO Global Health Observatory.

Notes: Add note: NA = not applicable; NCD = non-communicable disease.

Targets included in national (nutrition or other) plan

Stunting	Anaemia
Yes	Yes
Low birth weight	Child overweight
No	Yes
Exclusive breastfeeding	Wasting
Yes	No
Salt intake	Overweight adults and adolescents
No	Yes
Multisectoral comprehensive nutrition plan	
No	


Sources: WHO Global database on the Implementation of Nutrition Action (GINA), 2nd Global Nutrition Policy Review.

Economics and demography

Poverty rates (%) and GDP (PPP\$)

Under-five mortality (per 1,000 live births)

Government revenues (\$m)




Sources: World Bank 2019, IMF World Economic Outlook Database 2019.

Notes: PPP = purchasing power parity.



Source: UN Inter-agency Group for Child Mortality Estimation 2018.



Sources: IMF Article IV staff reports (country specific) and IMF World Economic Outlook Database (April 2019).

Income inequality

Gini index score ¹	Gini index rank ²	Year
36	75	2012

Sources: World Bank 2019.

Notes: ¹ 0 = perfect equality, 100 = perfect inequality. ² Countries are ranked from most equal (1) to most unequal (120).

Population

Population (000)	7,061,507	2018
Under-five population (000)	794	2019
Rural (%)	65	2018
>65 years (000)	299	2019

Sources: World Bank 2019, UN Population Division Department of Economic and Social Affairs 2019.

Population density of health workers per 1,000 people

Physicians	0.49	2014
Nurses and midwives	0.96	2014
Community health workers	NA	NA

Sources: WHO's Global Health Workforce Statistics, OECD, supplemented by country data.