AN UPPER BOUND FOR THE EUCLIDEAN TRAVELLING SALESMAN CONSTANT

DEVIN O'KEEFE

Abstract

By a theorem of Beardwood, Halton and Hammersley, the length of a shortest tour through n independent uniform random points in the unit square is asymptotic to $\beta\sqrt{n}$ for a constant β, which numerical estimates place near 0.712. We prove $\beta < 0.8618$. The best earlier upper bound, $\beta \le 0.8810$, is in a recent preprint of Dong and Cao. They cut the square into strips, take the points of each strip in blocks of six, and choose the order of visits within a block by a rule that is constant on the cells of a finite partition of the space of blocks. In each cell they bound the expected length of each segment of the path by the square root of its second moment. We change this last step: the length of a segment is a convex function of the positions of the points, and we bound its expectation by the Edmundson–Madansky inequality, a standard bound for convex functions, on a mesh much finer than the partition. For blocks of six points, $4.8 \cdot 10^7$ cells then give a better bound than their $1.6 \cdot 10^{13}$ cells. This makes blocks of ten points feasible, and these give the bound 0.8618. The proof ends in a computation in integer arithmetic, which three programs in different languages carry out with identical results.

1. Introduction

1.1. **The constant.** Let $X_1, X_2, \dots$ be independent random points, uniformly distributed in the unit square, and let L_n be the length of a shortest closed tour through $X_1, \dots, X_n$. Beardwood, Halton and Hammersley [1] proved that there is a constant β such that

$$\lim_{n\to\infty} \frac{L_n}{\sqrt{n}} = \beta \qquad \text{almost surely.}$$

The value of β is not known. Numerical estimates are 0.7124 ± 0.0002 [13] and 0.7120 ± 0.0002 [17]. The proved lower and upper bounds leave a much wider interval (Table 1.1). The constant is number 12 in the online list of optimisation constants [4].

All upper bounds for β so far come from the strip construction of [1]: cut the square into horizontal strips of height $h/\sqrt{n}$ and visit the points of each strip in the order of their first coordinates, one strip after another. With $h = \sqrt{3}$ this gives $\beta \le 0.92116\dots$. Few [8] had used a very similar argument for arbitrary points, as Steinerberger notes [19, Section 2.2].[1] Exchanging two consecutive points of a strip in certain configurations improves the bound by a small amount [19]. The same paper proposes to optimise the order of visits within groups of consecutive points; for disjoint groups of six points, with the first and the last point of each group kept in place, a simulation gave about 0.8902 [19, Section 2.5].

2020 *Mathematics Subject Classification.* Primary 60D05; Secondary 90C27, 68V05.

Key words and phrases. travelling salesman problem, Beardwood–Halton–Hammersley theorem, computer-assisted proof, integer arithmetic.

[1] Sections, tables and remarks of [19] and [9] are cited by their numbers in the arXiv versions; for the article [3] we cite its preprint [20], with the numbers of the preprint.

Yu and Carlsson [20] were the first to turn this proposal into a proved bound. Take the points of a strip in blocks of $k+1$ consecutive points, each block sharing its last point with the next, and visit the $k-1$ interior points of each block in the best order. This gives

$$\beta \le \frac{1}{kh}\,\mathbb{E}\min_{\pi} L_\pi, \tag{1.1}$$

where L_π is the length of the path that visits the interior points of a random block in the order π, measured after a magnification by the factor h (hence the h in the denominator); precise definitions are in Section 2.

The right-hand side is an integral over $2k+1$ variables of a minimum of $(k-1)!$ functions, and the difficulty is to bound it rigorously. For $k=4$ this is done in [20], with a partition of the domain found by a decision tree, one order for each part, and interval arithmetic; the published article [3] gives $\beta < 0.9038$.[2] Gaudio and Guan [9] improved the bound to 0.90367 by allowing a point near the boundary of a strip to be visited from the adjacent strip. They also estimated the right-hand side of (1.1) by simulation: for $h=\sqrt{3.75}\approx 1.94$ it is about 0.8748 for $k=5$, 0.8650 for $k=7$ and 0.8622 for $k=8$ [9, Table 2].

The best bound so far, $\beta \le 0.8810$, was proved with $k=5$ and $h=2$ by Dong and Cao in a preprint of 2 October 2026 [6]. They show first that, for blocks of six points, every rule that chooses the order of visits from the positions of the points gives an upper bound for β [6, Theorem 4]. The space of blocks is then divided into the cells of a product grid, with one order for each cell, and in each cell the expected length of a segment is bounded by the square root of its second moment.

They also give a statistical estimate: a bound computed from a random sample, which holds with probability at least $1-10^{-4}$ and is 0.8749 for their sample [6, Section 4.3]; it is not a proved bound, and Table 1.1 does not list it. Their conclusion names finer partitions and larger blocks as sources of further improvement.

TABLE 1.1. Proved bounds for the constant β. The upper bound of [1] is given as corrected in [19]; the value 0.92037 printed in [1] is a numerical error. The lower bound of [19] is given as corrected in [10], to $5/8 + 19/10368$. The last column gives the number k of (1.1) for the bounds that come from blocks.

	Bound	Source	k
Lower	0.625	Beardwood, Halton, Hammersley [1]	
	0.6268	Steinerberger [19]	
	0.6277	Gaudio, Jaillet [10]	
	0.6421	Dong, Cao [6], preprint	
Upper	$0.92116\ldots$	Beardwood, Halton, Hammersley [1]	1
	$0.92116\ldots - \frac{9}{16}\cdot 10^{-6}$	Steinerberger [19]	
	0.9038	Carlsson, Yu [3]	4
	0.90367	Gaudio, Guan [9]	4
	0.8810	Dong, Cao [6], preprint	5
	0.8618	Theorem 1.1	9

[2]The preprint [20] stated 0.90304, which is also the value in the list [4] at the time of writing; Table 1.1 follows the published articles, and the value of the article is quoted from its abstract.

1.2. **The result.**

Theorem 1.1. *The constant β of Beardwood, Halton and Hammersley satisfies $\beta < 0.8618$.*

The proof uses blocks of ten points ($k = 9$) and strips with $h = 11/5$. With smaller blocks the same method gives weaker bounds, starting from $\beta < 0.8789$ for $k = 5$ and $h = 2$ (Theorem 5.1).

1.3. **The method.** We take two things from [6]: rules that are constant on the cells of a product grid, and the theorem that every rule gives an upper bound for β. Since that preprint is recent and treats $k = 5$, we prove the theorem in full for general k (Theorem 2.2). What we change is how the expected length of a segment between two points of a block is bounded in a cell.

The square root of the second moment exceeds the expected length by an amount that depends on how much the length varies within the cell, so a good bound needs small cells. The bound 0.8810 was obtained with a grid of $12^5 \cdot 20^6 \approx 1.6 \cdot 10^{13}$ cells, reduced by symmetry [6].

We use instead the fact that the length of a segment is a convex function of the gaps of the block (the horizontal distances between consecutive points) and of the heights of its points. On an interval, the expectation of a convex function is at most the expectation of its chord; this is the Edmundson–Madansky inequality [7, 16]. We apply it to one variable at a time, on a mesh much finer than the cells (Proposition 3.5). A cell confines each gap and each height to an interval; where the interval of a gap is unbounded, we first truncate the gap and add the expectation of the excess (Lemma 3.3).

The accuracy of the bound for a segment is then set by the mesh and by the level of the truncation, which can be changed without changing the cells. The cells matter only because the rule chooses the same order throughout a cell. As with the second-moment bound of [6], our bound for a segment depends on the cell only through a few integers (Section 3.3), so that for $k = 9$ all of its values fit in a table with 4284 entries.

With $4^5 \cdot 6^6 \approx 4.8 \cdot 10^7$ cells we obtain 0.8789 for $k = 5$ and $h = 2$, against 0.8810 from the grid of $1.6 \cdot 10^{13}$ cells. This leaves room for larger blocks. The computation for $k = 9$ has $4^9 \cdot 6^{10} \approx 1.6 \cdot 10^{13}$ cells, about as many as that grid, though a cell takes more work for larger k.

The proof of Theorem 1.1 ends in a finite computation. Every quantity in it is rounded to an integer at a fixed scale, in the direction that makes the bound larger, so that only integer arithmetic is used (Section 4). We carried out the computation with three programs, in Python, C and Rust, which share no code and agree on every entry of the table of bounds for the segments, on all 262,144 cell sums of Section 4.1 and on the total (Section 5).

The total for $k = 9$ was also confirmed in the proof assistant Lean, by an evaluation in compiled code that the Lean kernel does not check. The agreement of these computations is evidence against errors of implementation, not against an error in the proofs, and the proofs of Sections 2 to 4 are not formalised. Section 5.2 says what these checks can and cannot show.

The separate steps of the method are not new. Applying the Edmundson–Madansky inequality to one independent variable at a time, on each interval of a partition, is a standard way to bound the expectation of a convex function in stochastic programming [14, 12], and one integral is bounded from above with chords in [20, Remark 6]. A program that Carlsson published in 2024 as a complement to [3, 20] already uses convexity on a grid: it bounds the length of

the path of an order on a cell by the largest of its values at the corners of the cell, and is designed to take for each cell the order for which this bound is smallest.[3] Rigorous arithmetic has been used before (intervals in [20], rational numbers in [6]), and two of the three identities by which our programs shorten the sum over the cells are used in [6] (Section 5.2). We add the combination of these steps and the scheme of integer arithmetic of Section 4.

The bound is still far from the numerical estimates near 0.712. For given k and h no rule gives less than the right-hand side of (1.1), which for $k = 9$ and $h = 11/5$ is about 0.854 by simulation (Table 5.1), and in the simulations of Section 6 the gain from each further point in a block is small.

2. The block bound

We prove that every rule for choosing the order of visits within a block gives an upper bound for β (Theorem 2.2).

2.1. **The constant in terms of a Poisson process.** For a finite set $A \subset \mathbb{R}^2$ let $L(A)$ be the length of a shortest closed tour through the points of A, with $L(A) = 0$ if A has fewer than two points. In this notation the L_n of Section 1 is $L(\{X_1, \dots, X_n\})$. We use the following standard consequence of the theorem of Beardwood, Halton and Hammersley [1, 18]. It is stated in this form in [6, equation (5)]; the short proof follows the sketch in [19, Section 2.1], attributed there to Steele.

Lemma 2.1. *Let Π be a Poisson process of intensity 1 in the plane, and let $\Lambda_W = [0, W]^2$. Then*

$$\lim_{W \to \infty} \frac{\mathbb{E}\, L(\Pi \cap \Lambda_W)}{W^2} = \beta.$$

Proof. Put $\ell_n = \mathbb{E}\, L_n$. By a theorem of Few [8], any $n \geq 1$ points of the unit square lie on a path of length at most $\sqrt{2n} + 1.75$. Closing the path adds at most $\sqrt{2}$, so $L(A) \leq c\sqrt{n}$ with $c = 2\sqrt{2} + 1.75$ for every set A of $n \geq 1$ points in the unit square, and $\ell_n/\sqrt{n} \to \beta$ by bounded convergence. Hence for every $\varepsilon > 0$ there is a constant C_ε with $|\ell_n - \beta\sqrt{n}| \leq \varepsilon\sqrt{n} + C_\varepsilon$ for all $n \geq 0$.

Let N be the number of points of Π in Λ_W. It has the Poisson distribution with mean W^2, and given $N = n$ the points are independent and uniform in Λ_W. Scaling by W gives $\mathbb{E}\, L(\Pi \cap \Lambda_W) = W\, \mathbb{E}\, \ell_N$, and therefore

$$\left|\mathbb{E}\, L(\Pi \cap \Lambda_W) - \beta W\, \mathbb{E}\sqrt{N}\right| \leq \varepsilon W\, \mathbb{E}\sqrt{N} + C_\varepsilon W.$$

By Jensen's inequality, $\mathbb{E}\sqrt{N} \leq W$. In the other direction, $\sqrt{N} - W = (N - W^2)/(\sqrt{N} + W) \geq -|N - W^2|/W$ and $\mathbb{E}|N - W^2| \leq (\operatorname{Var} N)^{1/2} = W$, so $\mathbb{E}\sqrt{N} \geq W - 1$. Dividing by W^2 and letting $W \to \infty$ and then $\varepsilon \to 0$ gives the claim. □

2.2. **Blocks, orders and rules.** Fix an integer $k \geq 1$ and a real number $h > 0$. Let $Z_1, \dots, Z_k$ be exponential random variables with mean 1 and let $U_0, \dots, U_k$ be uniform on $[0, 1]$, all independent. Put

$$P_i = (Z_1 + \cdots + Z_i,\ h^2 U_i), \qquad 0 \leq i \leq k.$$

[3] https://github.com/johngcarlsson-usc/tsp-constant-discretization, accessed 9 October 2026.

We call $(P_0, \dots, P_k)$ a *block*, the Z_i its *gaps* and the U_i its *heights*; the height of a point is its second coordinate divided by h^2. A block is a copy, magnified by the factor h, of $k+1$ consecutive points of a Poisson process of intensity 1 in a horizontal strip of height h: there the horizontal distances between consecutive points are exponential with mean $1/h$ and the vertical coordinates are uniform on an interval of length h. The proof of Theorem 2.2 makes this precise.

An *order* is a permutation π of $\{1, \dots, k-1\}$. It stands for the path that starts at P_0, visits the *interior points* $P_1, \dots, P_{k-1}$ in the order $P_{\pi(1)}, \dots, P_{\pi(k-1)}$, and ends at P_k; we write L_π for the length of this path. A *rule* is a measurable map ρ from $[0,\infty)^k \times [0,1]^{k+1}$ to the set of orders. It chooses an order from the gaps and heights of the block, and we write L_ρ for the length of the path it chooses (Figure 2.1).




FIGURE 2.1. Three consecutive blocks of a strip for $k = 9$ and $h = 11/5$, drawn from a simulation. Open circles mark the first and the last point of each block. The solid line is the path chosen by the rule of Section 5 for Theorem 1.1; where it differs from the order of the first coordinates, that order is shown by a dashed line.

Theorem 2.2 (Block bound)**.** *For every integer $k \geq 1$, every $h > 0$ and every rule ρ,*

$$\beta \leq \frac{\mathbb{E}\, L_\rho}{kh}.$$

We call $\mathbb{E}\, L_\rho/(kh)$ the *value* of the rule ρ. For $k = 1$ there is one order, and the value is the expected distance between consecutive points of a strip; this is the bound of [1], where $h = \sqrt{3}$. For the rule that chooses a shortest path (the first one in a fixed list of the orders, if there are several), the theorem is the bound (1.1) stated in [20, Section 3.1]; we call the value of this rule the *shortest-path value* for k and h, and no rule has a smaller value.

The computation of [20] already uses a rule of another kind, which chooses one order on each part of a partition; this order need not be shortest at every point of the part. For arbitrary rules and $k = 5$ the theorem is Theorem 4 of [6], and the proof below follows the proof given there, with 5 replaced by k and the strips joined in a slightly different way.

Proof. Every path in a block has length at most $k(Z_1 + \dots + Z_k + h^2)$, so $\mathbb{E}\, L_\rho < \infty$.

Let Π be a Poisson process of intensity 1 in the plane, let $q \geq 1$ be an integer and $W = qh$. For $1 \leq j \leq q$ let $S_j = [0,\infty) \times [(j-1)h, jh)$. Almost surely no point of Π has second coordinate W, and the points of Π in S_j have distinct first coordinates; let $P_1^j, P_2^j, \dots$ be these points in increasing order of the first coordinate. Add an auxiliary point P_0^j with first coordinate 0 and second coordinate $(j-1)h + Y_0^j$, where Y_0^j is uniform on $[0,h)$ and independent of everything else. For $i \geq 1$ let Δ_i^j be the first coordinate of P_i^j minus that of P_{i-1}^j, and let $(j-1)h + Y_i^j$ be

the second coordinate of P_i^j. Then the Δ_i^j, $i \geq 1$, are independent and exponential with mean $1/h$, the Y_i^j, $i \geq 0$, are independent and uniform on $[0, h)$, and the two families are independent of each other: the first coordinates of the points in S_j form a Poisson process of intensity h on $[0, \infty)$, and the second coordinates are independent marks [15].

For $b \geq 0$, the points $P_{kb}^j, P_{kb+1}^j, \ldots, P_{k(b+1)}^j$ form the bth block of the strip. Translating them so that P_{kb}^j has first coordinate 0 and the lower edge of the strip has second coordinate 0, and then multiplying by h, gives $k+1$ points with the distribution of a block: the gaps are $h\Delta_{kb+1}^j, \ldots, h\Delta_{kb+k}^j$ and the heights are $Y_{kb}^j/h, \ldots, Y_{kb+k}^j/h$. Let the rule ρ choose an order for it, and let C_b^j be the length of the corresponding path from P_{kb}^j to $P_{k(b+1)}^j$ in the strip. Then C_b^j has the distribution of L_ρ/h.

Let K_j be the number of points of Π in $S_j \cap \Lambda_W$ and $B_j = 1 + \lfloor K_j/k \rfloor$. The paths of the blocks $0, 1, \ldots, B_j - 1$ share endpoints and join to a walk Γ_j from P_0^j to $P_{kB_j}^j$, whose length we denote by $|\Gamma_j|$. Since $kB_j > K_j$, this walk visits every point of Π in $S_j \cap \Lambda_W$.

The bth block is used, that is $b < B_j$, exactly when $kb \leq K_j$. For $b = 0$ this always holds. For $b \geq 1$ it is the event $\Delta_1^j + \cdots + \Delta_{kb}^j \leq W$, which is determined by $\Delta_1^j, \ldots, \Delta_{kb}^j$. The length C_b^j is a function of $\Delta_{kb+1}^j, \ldots, \Delta_{kb+k}^j$ and $Y_{kb}^j, \ldots, Y_{kb+k}^j$, so it is independent of this event; this is the argument of Wald's identity. (The last block that is used contains the gap across the line $x = W$, which tends to be longer than a typical gap. This does not matter: each term of the sum below belongs to a fixed b, and none is conditioned on b being the last index used. Nor need the lengths C_b^j for different b be independent; they are not, since consecutive blocks share a point.) Since all terms are non-negative,

$$\mathbb{E}\,|\Gamma_j| = \sum_{b \geq 0} \mathbb{E}[C_b^j\,;\, b < B_j] = \sum_{b \geq 0} \mathbb{P}(b < B_j)\,\mathbb{E}\,C_b^j = \mathbb{E}B_j \cdot \frac{\mathbb{E}\,L_\rho}{h} \leq \Big(1 + \frac{hW}{k}\Big)\frac{\mathbb{E}\,L_\rho}{h}, \tag{2.1}$$

because $\mathbb{E}K_j = hW$.

The endpoint $P_{kB_j}^j$ of Γ_j has first coordinate $W + R_j$ with $R_j > 0$. Since $K_j < kB_j \leq K_j + k$, it is one of the first k points of Π in S_j to the right of Λ_W, and the kth of these has first coordinate $W + G_j$, where G_j is a sum of k independent exponential variables with mean $1/h$. Hence $R_j \leq G_j$ and $\mathbb{E}\,R_j \leq k/h$.

Now traverse Γ_1 from P_0^1 to its endpoint, go straight to the endpoint of Γ_2, traverse Γ_2 backwards to P_0^2, go straight to P_0^3, and so on through all q strips; finally return straight to P_0^1. The endpoints of Γ_j and Γ_{j+1} are at distance at most $2h + R_j + R_{j+1}$, and two auxiliary points in adjacent strips are at distance at most $2h$. Each R_j occurs in at most one of these segments between adjacent strips. The final segment starts at the endpoint of Γ_q if q is odd and at P_0^q if q is even, and in both cases its length is at most $2W + R_q$. In all, the R_j contribute at most $\sum_j R_j + R_q \leq 2\sum_j R_j$.

This closed walk passes through every point of $\Pi \cap \Lambda_W$. Leaving out all other points does not make it longer, by the triangle inequality. Therefore

$$L(\Pi \cap \Lambda_W) \leq \sum_{j=1}^{q} |\Gamma_j| + 2hq + 2\sum_{j=1}^{q} R_j + 2W,$$

and by (2.1), with $q = W/h$,

$$\mathbb{E}\, L(\Pi \cap \Lambda_W) \le W^2\, \frac{\mathbb{E}\, L_\rho}{kh} + W\Big(\frac{\mathbb{E}\, L_\rho}{h^2} + \frac{2k}{h^2} + 4\Big).$$

Dividing by W^2 and letting $q \to \infty$ gives the theorem, by Lemma 2.1. □

3. Cells and the chord inequality

This section applies Theorem 2.2 to rules that are constant on the cells of a product grid (Proposition 3.1) and bounds the expected length of a segment in a cell (Proposition 3.5). The result is the bound (3.3), which Sections 4 and 5 evaluate.

3.1. **Cells.** Fix a mesh width $\delta = 1/M$ with an integer $M \ge 2$, integers $m \ge 1$ and $n \ge 1$, and integers $0 = b_0 < b_1 < \cdots < b_{m-1}$. Put $t_c = b_c\delta$ for $0 \le c \le m-1$ and $t_m = \infty$. The *gap cells* are the intervals $I_c = [t_c, t_{c+1})$, $0 \le c \le m-1$, and the *height cells* are the intervals $J_a = [a/n, (a+1)/n)$, $0 \le a \le n-1$. (From here on n is the number of height cells and no longer a number of points.) A gap lies in I_c with probability

$$p_c = e^{-t_c} - e^{-t_{c+1}},$$

and a height lies in J_a with probability $1/n$.

A *cell* is a pair $\sigma = (g, a)$ of a gap tuple $g = (g_1, \dots, g_k) \in \{0, \dots, m-1\}^k$ and a height tuple $a = (a_0, \dots, a_k) \in \{0, \dots, n-1\}^{k+1}$. We also write σ for the event that $Z_i \in I_{g_i}$ for $1 \le i \le k$ and $U_i \in J_{a_i}$ for $0 \le i \le k$. Its probability is $p_g\, n^{-(k+1)}$ with $p_g = p_{g_1} \cdots p_{g_k}$, and almost surely exactly one of these events occurs. For $0 \le i < j \le k$ let

$$e_{ij}(\sigma) = \mathbb{E}\big[\, |P_iP_j| \;\big|\; \sigma \,\big]$$

be the conditional expected length of the segment P_iP_j, and put $e_{ji}(\sigma) = e_{ij}(\sigma)$. For an order π, write $\pi(0) = 0$ and $\pi(k) = k$, so that the path of π consists of the segments $P_{\pi(r-1)}P_{\pi(r)}$, $1 \le r \le k$.

Proposition 3.1. *Let $\bar e_{ij}(\sigma) = \bar e_{ji}(\sigma)$ be real numbers with $\bar e_{ij}(\sigma) \ge e_{ij}(\sigma)$ for all cells σ and all $0 \le i < j \le k$. Then*

$$\beta \le \frac{1}{kh} \sum_{\sigma=(g,a)} \frac{p_g}{n^{k+1}}\, \min_\pi \sum_{r=1}^{k} \bar e_{\pi(r-1)\pi(r)}(\sigma),$$

where the minimum is over all orders π.

Proof. For each cell σ let π_σ be an order at which the minimum is attained. Let ρ be the rule that chooses π_σ when the gaps and heights lie in σ, and the identity when some height equals 1, which has probability 0. It is constant on each of finitely many products of intervals, hence measurable, and

$$\mathbb{E}\, L_\rho = \sum_\sigma \mathbb{P}(\sigma)\, \mathbb{E}[L_{\pi_\sigma} \mid \sigma] = \sum_\sigma \mathbb{P}(\sigma) \sum_{r=1}^{k} e_{\pi_\sigma(r-1)\pi_\sigma(r)}(\sigma) \le \sum_\sigma \mathbb{P}(\sigma) \sum_{r=1}^{k} \bar e_{\pi_\sigma(r-1)\pi_\sigma(r)}(\sigma).$$

The last sum over r is the minimum in the statement, by the choice of π_σ, and the claim follows from Theorem 2.2. □

For $k = 5$, with cells of equal probability in place of ours and with $\bar{e}_{ij}(\sigma)$ the square root of the conditional second moment of $|P_iP_j|$, Proposition 3.1 is inequality (39) of [6]; the boundaries of those cells are not multiples of δ, and the proof above does not need them to be. We take for $\bar{e}_{ij}(\sigma)$ the bound of Proposition 3.5 below.

3.2. **The chord inequality.** If f is convex on $[x_0, x_1]$ and X is a random variable with values in $[x_0, x_1]$, then

$$\mathbb{E}\, f(X) \le \frac{x_1 - \mathbb{E}X}{x_1 - x_0}\, f(x_0) + \frac{\mathbb{E}X - x_0}{x_1 - x_0}\, f(x_1), \tag{3.1}$$

because the graph of f lies below its chord. This is the Edmundson–Madansky inequality [7, 16]; we call it the chord inequality. Proposition 3.5 below applies it to one variable at a time, as in the procedure of [14, Sections 3.4.2 and 3.5]: to the gaps of a block on the mesh $\delta\mathbb{Z}$ and to its heights on a mesh that cuts each height cell into equal intervals.

The weights in the next two lemmas are built from the numbers

$$\lambda = 1 - \frac{1 - e^{-\delta}}{\delta}, \qquad \mu = \frac{1 - e^{-\delta}}{\delta} - e^{-\delta};$$

both are positive, and $\lambda + \mu = 1 - e^{-\delta}$.

Lemma 3.2 (Gaps in a bounded gap cell)**.** *Let $N \ge 1$ be an integer, and let ξ be a random variable with density $e^{-x}/(1 - e^{-N\delta})$ on $[0, N\delta)$. For $0 \le i \le N$ put*

$$w_N(i) = \frac{[i < N]\, \lambda e^{-i\delta} + [i \ge 1]\, \mu e^{-(i-1)\delta}}{1 - e^{-N\delta}},$$

where $[\,\cdot\,]$ is 1 if the condition holds and 0 otherwise. Then $\sum_i w_N(i) = 1$, and $\mathbb{E}f(\xi) \le \sum_{i=0}^{N} w_N(i) f(i\delta)$ for every convex function f on $[0, N\delta]$.

Proof. For $0 \le i < N$, the variable ξ lies in $[i\delta, (i+1)\delta)$ with probability

$$q_i = \frac{e^{-i\delta}(1 - e^{-\delta})}{1 - e^{-N\delta}},$$

and given this, $\xi - i\delta$ has density $e^{-x}/(1 - e^{-\delta})$ on $[0, \delta)$ and mean $\theta = 1 - \delta e^{-\delta}/(1 - e^{-\delta})$. By (3.1),

$$\mathbb{E}f(\xi) \le \sum_{i=0}^{N-1} q_i \Big((1 - \tfrac{\theta}{\delta}) f(i\delta) + \tfrac{\theta}{\delta} f((i+1)\delta) \Big).$$

Since $(1 - e^{-\delta})\theta/\delta = \mu$ and $(1 - e^{-\delta})(1 - \theta/\delta) = \lambda$, the right-hand side is $\sum_i w_N(i) f(i\delta)$. For $f = 1$ it is also $\sum_i q_i = 1$, so the weights sum to 1. □

In the last gap cell, which is unbounded, the gap is truncated at a level T, and the expectation of the excess is added; a tail is split off in a similar way in the statistical estimate of [6, Section 4.3].

Lemma 3.3 (Gaps in the last gap cell)**.** *Let ξ be exponential with mean 1, let $N \geq 1$ be an integer and $T = N\delta$. For $0 \leq i \leq N$ put*

$$w_\infty(i) = [i < N]\, \lambda e^{-i\delta} + [i \geq 1]\, \mu e^{-(i-1)\delta} + [i = N]\, e^{-T}.$$

Then $\sum_i w_\infty(i) = 1$, and for every convex function f on $[0,\infty)$ with $|f(x) - f(y)| \leq |x - y|$,

$$\mathbb{E} f(\xi) \leq \sum_{i=0}^{N} w_\infty(i) f(i\delta) + e^{-T}.$$

Proof. We have $f(\xi) \leq f(\min(\xi, T)) + \max(\xi - T, 0)$, and $\mathbb{E}\max(\xi - T, 0) = e^{-T}$. With probability e^{-T} the variable $\min(\xi, T)$ equals T, and given $\xi < T$ it has the distribution of Lemma 3.2. Hence $\mathbb{E} f(\min(\xi, T)) \leq (1 - e^{-T}) \sum_i w_N(i) f(i\delta) + e^{-T} f(N\delta) = \sum_i w_\infty(i) f(i\delta)$. □

Lemma 3.4 (Heights)**.** *Let $\nu \geq 1$ be an integer and let U be uniform on J_a. Put $\eta_u = 1$ for $u \in \{0, \nu\}$ and $\eta_u = 2$ for $0 < u < \nu$. Then for every convex function f on the closure of J_a,*

$$\mathbb{E} f(U) \leq \sum_{u=0}^{\nu} \frac{\eta_u}{2\nu} f\Big(\frac{a\nu + u}{n\nu}\Big).$$

Proof. Cut J_a into ν intervals of equal length and apply (3.1) on each; given that U lies in one of them, its mean is the midpoint. □

3.3. **The bound for a segment.** Fix integers $\nu \geq 1$ and $T \geq 1$ for the rest of the paper: ν is the number of mesh intervals in a height cell, as in Lemma 3.4, with the numbers η_u defined there, and T is the level at which a gap in the last gap cell is truncated, measured from the left end t_{m-1} of that cell.

For a cell $\sigma = (g, a)$ and $0 \leq i < j \leq k$, the *key* of the segment P_iP_j is the vector $\kappa = (\kappa_0, \dots, \kappa_{m-1})$, where κ_c is the number of indices l with $i < l \leq j$ and $g_l = c$. It records how many of the gaps between the two points lie in each gap cell. For the gap cell $c < m - 1$ let $w^{(c)}$ be the sequence w_N of Lemma 3.2 with $N = b_{c+1} - b_c$, and let $w^{(m-1)}$ be the sequence w_∞ of Lemma 3.3 with $N = TM$. For a key κ let v_κ be the convolution of κ_0 copies of $w^{(0)}$, κ_1 copies of $w^{(1)}$, and so on, a sequence with finitely many non-zero terms $v_\kappa(s)$, $s \geq 0$, and let $b_\kappa = \sum_c \kappa_c b_c$. Finally let $\chi(t) = \sum \eta_u \eta_{u'}$ for $-\nu \leq t \leq \nu$, the sum over $0 \leq u, u' \leq \nu$ with $u - u' = t$, so that $\sum_t \chi(t) = 4\nu^2$.

Proposition 3.5. *Let $\sigma = (g, a)$ be a cell, let $0 \leq i < j \leq k$, let κ be the key of P_iP_j and $d = |a_i - a_j|$. Then $e_{ij}(\sigma) \leq \bar{e}(\kappa, d)$, where*

$$\bar{e}(\kappa, d) = \sum_{s \geq 0} v_\kappa(s) \sum_{t=-\nu}^{\nu} \frac{\chi(t)}{4\nu^2} \sqrt{\big((b_\kappa + s)\delta\big)^2 + \Big(\frac{h^2\,(d\nu + t)}{n\nu}\Big)^2} \;+\; \kappa_{m-1} e^{-T}. \tag{3.2}$$

Proof. Given σ, the variables Z_l $(i < l \leq j)$, U_i and U_j are independent. If $g_l = c < m - 1$ then $Z_l - t_c$ has the distribution of Lemma 3.2 with $N = b_{c+1} - b_c$, and if $g_l = m - 1$ then $Z_l - t_{m-1}$ is exponential with mean 1. The heights U_i and U_j are uniform on J_{a_i} and J_{a_j}. The length

$$|P_iP_j| = \sqrt{(Z_{i+1} + \cdots + Z_j)^2 + h^4 (U_i - U_j)^2}$$

is the Euclidean norm of a vector that depends affinely on these variables, so it is a convex function of each of them when the others are fixed, and it changes by at most $|x-y|$ when one gap changes from x to y.

Apply Lemma 3.2, 3.3 or 3.4 to one variable, with the others fixed, and integrate over the others. This replaces the variable by a discrete one with values on the mesh, adds e^{-T} if the variable is a gap in the last gap cell, and does not decrease the expectation. The new integrand is a combination of the old one at several values of the replaced variable, with non-negative weights that sum to 1, plus a constant. So it is again convex in each remaining variable and changes by at most $|x-y|$ when a gap changes from x to y, and the step can be repeated until all variables have been replaced. Therefore

$$e_{ij}(\sigma) \le \mathbb{E}\sqrt{\big(\hat{Z}_{i+1}+\cdots+\hat{Z}_j\big)^2 + h^4\big(\hat{U}_i-\hat{U}_j\big)^2} \;+\; \kappa_{m-1}e^{-T},$$

where the variables with hats are independent, $\hat{Z}_l = (b_{g_l}+\iota_l)\delta$ with $\mathbb{P}(\iota_l = s) = w^{(g_l)}(s)$, and $\hat{U}_i = (a_i\nu+\zeta_i)/(n\nu)$ with $\mathbb{P}(\zeta_i = u) = \eta_u/(2\nu)$, and likewise for $\hat{U}_j$. The sum $\hat{Z}_{i+1}+\cdots+\hat{Z}_j$ equals $(b_\kappa+s)\delta$ with probability $v_\kappa(s)$. The difference $\zeta_i-\zeta_j$ equals t with probability $\chi(t)/(4\nu^2)$ and has a symmetric distribution, so the expectation does not change if a_i-a_j is replaced by d. This gives (3.2). □

The bound $\bar{e}(\kappa,d)$ depends on the cell only through the key and the difference d. The keys are the vectors of m non-negative integers whose sum lies between 1 and k. There are $\binom{m+k}{k}-1$ of them, so all values of $\bar{e}$ fit in a table with $n\big(\binom{m+k}{k}-1\big)$ entries. Propositions 3.1 and 3.5 together give the bound on which the computation rests:

$$\beta \le \frac{1}{kh}\sum_g p_g \,\frac{1}{n^{k+1}}\sum_a \;\min_\pi \sum_{r=1}^{k} \bar{e}\big(\kappa_{\pi(r-1)\pi(r)},\, |a_{\pi(r-1)}-a_{\pi(r)}|\big), \tag{3.3}$$

where $\kappa_{ij} = \kappa_{ji}$ denotes the key of P_iP_j in the cell (g,a).

4. Integer arithmetic

The right-hand side of (3.3) is a finite sum, but its terms involve exponentials and square roots. We bound it from above by a quotient of integers, computed with integer arithmetic alone. This section defines these integers and proves the bound (Theorem 4.1).

The method is to replace each real quantity $x \ge 0$ that enters (3.3) by an integer $\mathsf{X} \ge Sx$, where the *scale* S is a fixed power of two, and to compute with these integers in place of the real numbers. A sum of such integers bounds the sum of the quantities at the same scale. A product of two of them has the product of their scales, and a division by one scale, rounded up, brings it back to the other: if $S, S' > 0$, and $\mathsf{X}_i \ge Sx_i$ and $\mathsf{Y}_i \ge S'y_i$ with real $x_i, y_i \ge 0$, then

$$\Big\lceil \frac{1}{S}\sum_i \mathsf{X}_i\mathsf{Y}_i \Big\rceil \ge S'\sum_i x_iy_i. \tag{4.1}$$

An upper bound for a difference or a quotient needs a lower bound for the term that is subtracted or for the divisor. This occurs only in Steps 1 and 2 below, for $e^{-\delta}$, the factors of its series and its powers, and these are therefore computed with bounds on both sides.

The integers of the computation are set in sans-serif type. Table 4.1 lists them with the real quantities that they bound and with their scales: $Q = 2^{62}$ for the powers of $e^{-\delta}$ and for λ and μ, $D = 2^{48}$ for probabilities and for the weights $w^{(c)}$ and v_κ, and $F = 2^{32}$ for lengths. For real

y, $\lfloor y \rfloor$ is the greatest integer that is at most y and $\lceil y \rceil$ the least integer that is at least y; for an integer $x \geq 0$, $\lceil \sqrt{x}\, \rceil$ is thus the least integer $z \geq 0$ with $z^2 \geq x$.

TABLE 4.1. The integers of Section 4.1, with the step that defines each. Each integer is at least the quantity multiplied by the scale, except lo_1 and $\mathsf{Rlo}[i]$, which are at most that product.

Step	Integer	Quantity	Scale
1	lo_1, up_1	$e^{-\delta}$	Q
	$\mathsf{Rlo}[i]$, $\mathsf{Rup}[i]$	$e^{-i\delta}$	Q
2	C_λ, C_μ	λ, μ	Q
	$\mathsf{W}_c(i)$	$w^{(c)}(i)$	D
	P_c	p_c	D
	E_T	e^{-T}	F
3	$\mathsf{V}_\kappa(s)$	$v_\kappa(s)$	D
4	$\mathsf{Dist}(j,\tau)$	$\sqrt{(j\delta)^2 + \big(h^2\tau/(n\nu)\big)^2}$	F
	$\mathsf{G}(j,d)$	the inner sum in (3.2), with $b_\kappa + s = j$	F
5	$\mathsf{Tab}(\kappa,d)$	$\bar{e}(\kappa,d)$	F
6	S_g	the sum over a in (3.3)	F
7	A_0	the same sum, multiplied by p_g and summed over g	F

The parameters are k, M, m, $b_1 < \cdots < b_{m-1}$, n, ν and T as in Section 3, and the strip height $h = h_1/h_2$, a quotient of positive integers. Put $n_c = b_{c+1} - b_c$ for $c < m-1$ and $n_{m-1} = N_T = TM$; for $m = 1$ the only gap cell, $I_0 = [0,\infty)$, is the last one. The keys are the vectors $\kappa = (\kappa_0, \ldots, \kappa_{m-1})$ of non-negative integers with $1 \leq \kappa_0 + \cdots + \kappa_{m-1} \leq k$, and b_κ, η_u, $\chi(t)$ and κ_{ij} are as in Section 3.

4.1. **Definitions.** The seven steps build the right-hand side of (3.3) from the inside: the powers of $e^{-\delta}$ (Step 1), the weights of a single gap and the probabilities of the gap cells (Step 2), the weights of a sum of gaps (Step 3), the lengths (Step 4), the bound $\bar{e}$ for a segment (Step 5), the sum over the height tuples (Step 6) and the sum over the gap tuples (Step 7).

Step 1: exponentials. Put $\mathsf{lo}_{13} = 0$, $\mathsf{up}_{13} = Q$ and, for $j = 12, 11, \ldots, 1$,

$$\mathsf{up}_j = Q - \lfloor \mathsf{lo}_{j+1}/(Mj) \rfloor, \qquad \mathsf{lo}_j = Q - \lceil \mathsf{up}_{j+1}/(Mj) \rceil.$$

This is the series of $e^{-\delta}$ in the nested form $1 - \frac{\delta}{1}(1 - \frac{\delta}{2}(1 - \frac{\delta}{3}(\cdots)))$, in which the thirteenth factor is replaced by its bounds 0 and 1. The powers of $e^{-\delta}$ follow: put $\mathsf{Rlo}[0] = \mathsf{Rup}[0] = Q$ and, for $1 \leq i \leq \max(b_{m-1}, N_T)$,

$$\mathsf{Rup}[i] = \lceil \mathsf{Rup}[i-1] \cdot \mathsf{up}_1/Q \rceil, \qquad \mathsf{Rlo}[i] = \lfloor \mathsf{Rlo}[i-1] \cdot \mathsf{lo}_1/Q \rfloor.$$

Step 2: weights and probabilities. By Lemmas 3.2 and 3.3, the gap cell c has the weights

$$w^{(c)}(i) = \frac{\omega_c(i)}{1 - e^{-n_c\delta}} \quad \text{for } c < m-1, \qquad w^{(m-1)}(i) = \omega_{m-1}(i) + [i = N_T]\, e^{-T},$$
$$\text{where} \quad \omega_c(i) = [i < n_c]\, \lambda e^{-i\delta} + [i \geq 1]\, \mu e^{-(i-1)\delta} \quad \text{for } 0 \leq i \leq n_c,$$

and $\lambda = 1 - M(1-e^{-\delta})$, $\mu = M(1-e^{-\delta}) - e^{-\delta}$. The integers of this step have the same form. Put $\mathsf{C}_\lambda = Q - M(Q - \mathsf{up}_1)$ and $\mathsf{C}_\mu = M(Q - \mathsf{lo}_1) - \mathsf{lo}_1$. For $0 \le c \le m-1$ and $0 \le i \le n_c$ let

$$\mathsf{num}_c(i) = [i < n_c]\,\mathsf{Rup}[i]\,\mathsf{C}_\lambda + [i \ge 1]\,\mathsf{Rup}[i-1]\,\mathsf{C}_\mu.$$

A term whose condition fails is left out, so that $\mathsf{Rup}[-1]$ does not occur. The divisors below are $2^{14} = Q/D$, $2^{76} = Q^2/D$ and $2^{30} = Q/F$. For $c < m-1$ put

$$\mathsf{W}_c(i) = \Big\lceil \frac{\lceil \mathsf{num}_c(i)/2^{14} \rceil}{Q - \mathsf{Rup}[n_c]} \Big\rceil, \qquad \mathsf{P}_c = \Big\lceil \frac{\mathsf{Rup}[b_c] - \mathsf{Rlo}[b_{c+1}]}{2^{14}} \Big\rceil,$$

and for the last gap cell

$$\mathsf{W}_{m-1}(i) = \lceil \mathsf{num}_{m-1}(i)/2^{76} \rceil + [i = N_T]\,\lceil \mathsf{Rup}[N_T]/2^{14} \rceil, \qquad \mathsf{P}_{m-1} = \lceil \mathsf{Rup}[b_{m-1}]/2^{14} \rceil.$$

Finally $\mathsf{E}_T = \lceil \mathsf{Rup}[N_T]/2^{30} \rceil$. Here C_λ and C_μ stand for λ and μ at the scale Q, and $\mathsf{num}_c(i)$, a sum of products of two integers at the scale Q, stands for $\omega_c(i)$ at the scale Q^2. The divisor $Q - \mathsf{Rup}[n_c]$ stands for $1 - e^{-n_c\delta}$ and the difference in P_c for p_c, both at the scale Q.

Step 3: keys. The sequences V_κ are defined by recursion on $\kappa_0 + \cdots + \kappa_{m-1}$. If κ is a unit vector, with $\kappa_c = 1$ and all other entries 0, put $\mathsf{V}_\kappa = \mathsf{W}_c$. Otherwise let c be the largest index with $\kappa_c > 0$, let κ' be κ with κ_c reduced by one, and put

$$\mathsf{V}_\kappa(s) = \Big\lceil \frac{1}{D} \sum_{i+j=s} \mathsf{V}_{\kappa'}(i)\,\mathsf{W}_c(j) \Big\rceil, \qquad 0 \le s \le \sum_c \kappa_c n_c,$$

where the sum is over the pairs with $0 \le i \le \sum_c \kappa'_c n_c$ and $0 \le j \le n_c$. This mirrors the convolution that defines v_κ in Section 3, one factor at a time.

Step 4: lengths. Let $\alpha = h_2^2\, n\nu$ and $\gamma = M h_1^2$. These numbers clear the denominators of the square root in (3.2): with $b_\kappa + s = j$ and $|d\nu + t| = \tau$ that square root is

$$\sqrt{(j\delta)^2 + \Big(\frac{h^2\tau}{n\nu}\Big)^2} = \frac{\sqrt{(\alpha j)^2 + (\gamma\tau)^2}}{\alpha M},$$

with an integer under the root on the right. For integers $j, \tau \ge 0$ put

$$\mathsf{Dist}(j,\tau) = \Big\lceil \frac{\lceil \sqrt{((\alpha j)^2 + (\gamma\tau)^2) F^2}\, \rceil}{\alpha M} \Big\rceil,$$

and for $0 \le d \le n-1$

$$\mathsf{G}(j,d) = \Big\lceil \frac{1}{4\nu^2} \sum_{t=-\nu}^{\nu} \chi(t)\,\mathsf{Dist}(j, |d\nu + t|) \Big\rceil.$$

Since $\lceil \lceil y \rceil / q \rceil = \lceil y/q \rceil$ for real $y \ge 0$ and integers $q \ge 1$, the inner rounding has no effect: $\mathsf{Dist}(j,\tau) = \lceil F\sqrt{j^2 + (\gamma\tau/\alpha)^2}/M \rceil$ with $\gamma/\alpha = Mh^2/(n\nu)$. The integers of this section therefore depend on h_1 and h_2 only through h.

Step 5: the table. For every key κ and $0 \le d \le n-1$,

$$\mathsf{Tab}(\kappa, d) = \Big\lceil \frac{1}{D} \sum_s \mathsf{V}_\kappa(s)\,\mathsf{G}(b_\kappa + s, d) \Big\rceil + \kappa_{m-1}\mathsf{E}_T,$$

the sum over $0 \le s \le \sum_c \kappa_c n_c$. This mirrors (3.2) term by term.

Step 6: cell sums. For a gap tuple g,

$$\mathsf{S}_g = \sum_a \min_\pi \sum_{r=1}^{k} \mathsf{Tab}(\kappa_{\pi(r-1)\pi(r)},\, |a_{\pi(r-1)} - a_{\pi(r)}|),$$

where a runs over the n^{k+1} height tuples, π over the orders, and κ_{ij} is the key of P_iP_j in the cell (g, a). We call S_g the *cell sum* of g: it is a sum over the n^{k+1} cells with this gap tuple.

Step 7: the total. Put $\mathsf{A}_k(g) = \mathsf{S}_g$. For $j = k, k-1, \dots, 1$ and every $(g_1, \dots, g_{j-1})$ with entries in $\{0, \dots, m-1\}$ put

$$\mathsf{A}_{j-1}(g_1, \dots, g_{j-1}) = \Big\lceil \frac{1}{D} \sum_{c=0}^{m-1} \mathsf{P}_c\, \mathsf{A}_j(g_1, \dots, g_{j-1}, c) \Big\rceil.$$

The *total* is A_0, the value at the empty tuple. This mirrors the sum of $p_g\mathsf{S}_g$ over the gap tuples, taken one gap at a time from g_k back to g_1.

4.2. **The bound in integers.**

Theorem 4.1. *For all parameters as above with $M \le 2^{61}$,*

$$\beta \le \frac{h_2\, \mathsf{A}_0}{n^{k+1}\, F\, k\, h_1}.$$

Proof. We go through the steps and prove for each integer the inequality of Table 4.1. The last line of the table, inserted in (3.3), is the theorem.

Step 1. Write $x = \delta = 1/M$, and for $j \ge 1$ let

$$\vartheta_j = \sum_{i \ge 0} \frac{(-x)^i\, (j-1)!}{(j-1+i)!} = 1 - \frac{x}{j} + \frac{x^2}{j(j+1)} - \cdots.$$

Then $\vartheta_1 = e^{-x}$ and $\vartheta_j = 1 - (x/j)\vartheta_{j+1}$, so the ϑ_j are the factors of the nested form in Step 1 of Section 4.1. Since $0 < x \le 1$, the series is alternating with terms that do not increase and tend to 0, so $0 \le \vartheta_j \le 1$. We claim that

$$0 \le \mathsf{lo}_j \le Q\vartheta_j \le \mathsf{up}_j \le Q \qquad \text{for } 1 \le j \le 13.$$

This holds for $j = 13$. Suppose that it holds for $j+1$. Then $\mathsf{up}_j \le Q$ because $\mathsf{lo}_{j+1} \ge 0$, and $\mathsf{lo}_j \ge Q - \lceil Q/(Mj) \rceil \ge 0$ because $\mathsf{up}_{j+1} \le Q$. Moreover

$$\mathsf{up}_j \ge Q - \frac{\mathsf{lo}_{j+1}}{Mj} \ge Q - \frac{Q\vartheta_{j+1}}{Mj} = Q\vartheta_j, \qquad \mathsf{lo}_j \le Q - \frac{\mathsf{up}_{j+1}}{Mj} \le Q - \frac{Q\vartheta_{j+1}}{Mj} = Q\vartheta_j.$$

In particular $\mathsf{lo}_1 \le Qe^{-\delta} \le \mathsf{up}_1$, and induction on i gives $\mathsf{Rlo}[i] \le Qe^{-i\delta} \le \mathsf{Rup}[i]$: for Rup by (4.1), and for Rlo in the same way with inequalities and rounding reversed, as $\mathsf{lo}_1 \ge 0$.

Step 2 also needs that $\mathsf{Rup}[i] < Q$ for $i \ge 1$. Indeed $\mathsf{lo}_2 \ge Q - \lceil Q/(2M) \rceil \ge Q/2 \ge M$, so $\mathsf{up}_1 = Q - \lfloor \mathsf{lo}_2/M \rfloor < Q$; this is the only use of $M \le 2^{61}$. Since $\mathsf{up}_1 \le Q$, the sequence Rup is non-increasing, and so $\mathsf{Rup}[i] \le \mathsf{Rup}[1] = \mathsf{up}_1 < Q$.

Step 2. With λ, μ and $\omega_c(i)$ as in Step 2 of Section 4.1, the bounds for $e^{-\delta}$ give

$$Q\lambda \le Q - M(Q - \mathsf{up}_1) = \mathsf{C}_\lambda, \qquad Q\mu \le M(Q - \mathsf{lo}_1) - \mathsf{lo}_1 = \mathsf{C}_\mu.$$

Since λ and μ are positive (Section 3.2) and $\mathsf{Rup}[i] \ge Qe^{-i\delta}$, it follows that $\mathsf{num}_c(i) \ge Q^2\omega_c(i)$.

For $c < m-1$ the denominator of $w^{(c)}(i)$ satisfies $Q(1-e^{-n_c\delta}) \geq Q - \mathsf{Rup}[n_c] > 0$, by Step 1 and because $n_c \geq 1$. Since $Q/2^{14} = D$,

$$\mathsf{W}_c(i) \geq \frac{\mathsf{num}_c(i)}{2^{14}\,(Q - \mathsf{Rup}[n_c])} \geq \frac{Q^2\,\omega_c(i)}{2^{14}\,Q\,(1-e^{-n_c\delta})} = D\,w^{(c)}(i).$$

In the last gap cell there is no denominator. Since $Q^2/2^{76} = D$ and $\mathsf{Rup}[N_T] \geq Qe^{-T}$,

$$\mathsf{W}_{m-1}(i) \geq D\,\omega_{m-1}(i) + [i = N_T]\,De^{-T} = D\,w^{(m-1)}(i).$$

For the probabilities, Step 1 gives $\mathsf{Rup}[b_c] \geq Qe^{-t_c} > Qe^{-t_{c+1}} \geq \mathsf{Rlo}[b_{c+1}]$ for $c < m-1$, so the numerator of P_c is positive. With $Q/2^{14} = D$ and $Q/2^{30} = F$,

$$\mathsf{P}_c \geq D(e^{-t_c} - e^{-t_{c+1}}) = D\,p_c, \qquad \mathsf{P}_{m-1} \geq De^{-t_{m-1}} = D\,p_{m-1}, \qquad \mathsf{E}_T \geq Fe^{-T}.$$

Step 3. The sequence v_κ is the convolution of $v_{\kappa'}$ and $w^{(c)}$. By induction on $\kappa_0 + \cdots + \kappa_{m-1}$, with Step 2 and (4.1), $\mathsf{V}_\kappa(s) \geq D\,v_\kappa(s)$ for all s.

Step 4. By the identity in Step 4 of Section 4.1, $\mathsf{Dist}(j,\tau)$ is at least F times the square root in (3.2) with $b_\kappa + s = j$ and $|d\nu + t| = \tau$: the integer square root is at least the real one, and the quotient is rounded up. That square root depends on $d\nu + t$ only through $|d\nu + t|$, so $\mathsf{G}(j,d)$ is at least F times the inner sum in (3.2), by (4.1) with $S = 4\nu^2$ and $x_t = \chi(t)/S$.

Step 5. By Steps 3 and 4 and (4.1), the first term of $\mathsf{Tab}(\kappa,d)$ is at least F times the double sum in (3.2), and $\kappa_{m-1}\mathsf{E}_T \geq F\kappa_{m-1}e^{-T}$ by Step 2. Hence $\mathsf{Tab}(\kappa,d) \geq F\,\bar{e}(\kappa,d)$.

Steps 6 and 7. If every term of every sum is replaced by an upper bound, the minimum of the sums does not decrease. By Step 5, therefore, S_g is at least F times the sum over a in (3.3). By induction on $k-j$, using (4.1) with the scales D and 1 and the bounds $\mathsf{P}_c \geq D\,p_c$ of Step 2,

$$\mathsf{A}_j(g_1,\ldots,g_j) \geq \sum_{g_{j+1},\ldots,g_k} p_{g_{j+1}}\cdots p_{g_k}\,\mathsf{S}_g,$$

so $\mathsf{A}_0 \geq \sum_g p_g \mathsf{S}_g$. With the bound for S_g this is the last line of Table 4.1, and inserting it in (3.3) gives $\beta \leq \mathsf{A}_0/(n^{k+1}Fkh)$, which is the claim because $h = h_1/h_2$. □

Remark 4.2. No estimate of rounding errors is needed in this proof, because every rounding is in the direction that increases the total: the upper bounds are rounded up, and the lower bounds lo_j and Rlo are rounded down. The effect of the rounding and of the mesh on the result is estimated in Section 5.3.

5. The computation

5.1. **Parameters and results.** We take $M = 128$, so that $\delta = 1/128$, and $m = 4$ gap cells with $(b_1, b_2, b_3) = (48, 112, 208)$, that is

$$[0, \tfrac{3}{8}), \qquad [\tfrac{3}{8}, \tfrac{7}{8}), \qquad [\tfrac{7}{8}, \tfrac{13}{8}), \qquad [\tfrac{13}{8}, \infty),$$

together with $n = 6$ height cells, $\nu = 64$ and $T = 16$. With these parameters fixed, we ran the computation of Section 4.1 for five choices of k and h.

Theorem 5.1. *For the parameters above, the total* A_0 *of Section 4.1 has the following values. The last column is rounded up in its last digit.*

k	h	A_0	$h_2\mathsf{A}_0/(n^{k+1}Fkh_1)$
5	2	1,761,033,937,996,622	0.878820870371
6	2	12,593,568,504,391,125	0.872868757138
7	21/10	92,054,418,307,910,785	0.868075379727
8	21/10	628,708,228,549,677,266	0.864607001369
9	11/5	4,430,997,795,015,293,889	0.861715646305

Theorem 5.1 states the result of a finite computation with integers, and its proof is that computation. Section 5.2 describes the programs that carried it out and the evidence that they did so correctly. Where a bound of Theorem 5.1 is quoted below with six decimals, it is rounded up as well.

Proof of Theorem 1.1. By Theorem 4.1 and the last row of the table in Theorem 5.1, $\beta \le 5\,\mathsf{A}_0/(6^{10}\cdot 2^{32}\cdot 9\cdot 11)$ with $\mathsf{A}_0 = 4{,}430{,}997{,}795{,}015{,}293{,}889$. This is less than 0.8618. □

The *cost* of an order in a cell is the sum of the table entries of its segments, so that S_g is the sum over a of the least cost in the cell (g, a). For each of the computations of Theorem 5.1, *the rule* is that of the proof of Proposition 3.1 with $\bar{e}_{ij}(\sigma) = \mathsf{Tab}(\kappa_{ij}, |a_i - a_j|)/F$: in each cell it chooses an order of least cost. Where several orders have the least cost the choice is free, and Theorem 5.1 does not depend on it.

The values of k, m and n were limited by the time that the computation takes: there are $m^k n^{k+1}$ cells, that is $1.6\cdot 10^{13}$ for $k = 9$. The cell boundaries, among the multiples of 1/8, and the strip heights, among the multiples of 1/10, were chosen by simulating the rule for trial values; this choice affects the size of the bound and not its validity.

5.2. **The programs.** The three programs, in Python, C and Rust, are in the repository that accompanies this paper,[4] where they are called checkers. They share no code. The file `RECIPE.md` of the repository restates the definitions of Section 4.1, and the C program and the Rust program were each first written from this file alone. All three use integer arithmetic only; where values are kept in words of a fixed width, each operation that could overflow is tested, or a bound checked beforehand excludes the overflow. Each program computes from the parameters the table Tab, all m^k cell sums S_g and the total, and compares them with the values stored in the repository, in a file that is the same for all three programs. The programs therefore agree not only on the totals but on every table entry and every cell sum; for $k = 9$ these are 4284 and 262,144 numbers.

Three identities. Step 6 takes almost all of the time. Each program finds the minimum over the $(k-1)!$ orders by dynamic programming over subsets of the interior points [2, 11], and shortens the sum over the cells by three identities, which Section 7 of `RECIPE.md` derives. (i) *Reversal*: the cell with gap tuple $(g_k, \dots, g_1)$ and height tuple $(a_k, \dots, a_0)$ is the cell (g, a) read from its other end; its paths are those of (g, a) in reverse, with the same costs, so a gap tuple and its reverse have the same cell sum. (ii) *Reflection*: replacing every a_i by $n - 1 - a_i$ changes no difference $|a_i - a_j|$, so a cell and its reflection have the same least cost. (iii) *Shared paths*:

[4] https://github.com/devinokeefe/travelling-sales-person-constant-bounds

the table entry of a segment between two interior points depends neither on g_1 and g_k nor on a_0 and a_k, so for $k \geq 3$ the least cost of a path from one interior point to another through all interior points is the same in the m^2n^2 cells that differ only in these four numbers, and is computed once for all of them. Identities (i) and (ii) are two of the three symmetries by which the number of cells to be evaluated is reduced in [6, Section 4.2, after (39)]; the remaining one, a common vertical translation of all points, we do not use.

Recomputation and runs. Earlier versions of the three programs use neither (i) nor (ii), and only the C version uses a part of (iii). For $k \leq 7$ we computed every cell sum with these versions as well, and for $k = 8$ with the earlier version of the C program. The results were the same. For $k = 9$ we did not do this, so there the cell sums rest on the identities, in the routines whose results for $k \leq 8$ were confirmed in this way. The evaluation in Lean described below confirms the total for $k = 9$ by a routine that does not use (i), and whose use of (ii) and (iii) is proved correct.

All runs of the three programs printed the same result.

A check in Lean. For $k = 5, \ldots, 9$ the totals were also confirmed in the proof assistant Lean 4 [5]. The Lean development in the repository states the definitions of Section 4.1 as directly as the language allows and, beside them, an efficient version, which uses (ii) and (iii) but not (i); theorems checked by the Lean kernel say that the two give the same total for all parameters. For each of these k the equation for A_0 in Theorem 5.1 is then a theorem of Lean, whose proof evaluates the efficient version in compiled code (the tactic `native_decide`). The kernel does not check such an evaluation, so these theorems rest on the Lean compiler, interpreter and runtime as well, and on the hardware. The numbers to be proved were taken from the programs, so Lean could only confirm them or fail. The Lean definitions were transcribed from the file `RECIPE.md` and compared with it only by reading the two side by side. The proofs of Sections 2 to 4 are not formalised. The file `lean/README.md` of the repository says in full what is proved and what is assumed.

5.3. **Controls.** The controls of this section are not part of the proof. They use floating-point arithmetic or simulation, and serve to detect a mistake in Sections 3 and 4 or in the programs. Their scripts and records are in the repository. Of the three programs they use only routines of the Python program that read the parameters and the stored values and list the keys and the pairs of points. The file `controls/README.md` says what each control can and cannot show, and describes further controls.

The segments. For every key and every d we computed the conditional expected length of a segment by numerical integration, without the lemmas of Section 3. For each of the five computations of Theorem 5.1, every table entry $\mathsf{Tab}(\kappa, d)/F$ lies above the integral, by a relative margin between $4.8 \cdot 10^{-8}$ and $1.9 \cdot 10^{-4}$. This control concerns the table, that is Proposition 3.5 and Steps 1 to 5; the cell sums and the total do not enter it.

The value of the rule. We also computed, on a mesh twice as fine as that of the computation, the upper bound of Proposition 3.5 and the matching lower bound from Jensen's inequality. Summing the lower bounds, and likewise the upper bounds, along the path that the rule chooses in a cell, averaging over the cells with their probabilities and dividing by kh gives two numbers below and above the value of the rule: the ends of the *bracket* in Table 5.1.

For $k > 7$ the sums over all cells would be too costly. There the amounts by which the table entries along the path exceed the two bounds are averaged over the cells of the simulated blocks below, with standard errors below 10^{-9}, and subtracted from the bound of Theorem 5.1; the bracket is then computed from the total A_0 and would not show an error in the cell sums. For every k the bound exceeds the lower end of the bracket, and hence the value of the rule, by less than 10^{-5}; this is all that the mesh, the truncation and the rounding add to the bound.

TABLE 5.1. Controls for the computations of Theorem 5.1. The bound is rounded up. The bracket for the value of the rule is computed in floating-point arithmetic, without a bound for its rounding errors; its lower end is rounded down and its upper end up; for $k > 7$ it also has a sampling error, below 10^{-9}. The last two columns are estimates from $4 \cdot 10^8$ simulated blocks for each k, with standard errors. The samples for the different k were drawn with one seed, so they are not independent of each other.

k	h	Bound	Rule, bracket	Rule, simulated	Shortest path, simulated
5	2	0.878821	[0.878814, 0.878817]	0.878828 ± 0.000012	0.874265 ± 0.000012
6	2	0.872869	[0.872861, 0.872865]	0.872880 ± 0.000011	0.867701 ± 0.000011
7	21/10	0.868076	[0.868068, 0.868071]	0.868067 ± 0.000010	0.861794 ± 0.000010
8	21/10	0.864608	[0.864599, 0.864603]	0.864612 ± 0.000009	0.858016 ± 0.000009
9	11/5	0.861716	[0.861708, 0.861711]	0.861713 ± 0.000009	0.854004 ± 0.000009

Simulated blocks. We drew $4 \cdot 10^8$ independent blocks for each k and measured, in each block, the lengths of the path chosen by the rule and of a shortest path (Table 5.1). The simulated values of the rule lie within two standard errors of their brackets. Some lie above the bound of Theorem 5.1, as is to be expected: the bound exceeds the value of the rule by less than one standard error.

The cost of the order that the rule chooses, averaged over the same blocks and divided by Fkh, estimates the bound of Theorem 5.1 itself, from the table alone and without the cell sums. For $k = 9$ it gave 0.861722 ± 0.000008, against the bound 0.861716, and for every k it lies within two standard errors of the bound. For $k = 8$ and $k = 9$ an error in the total that changed the bound by $3 \cdot 10^{-5}$, which is more than three standard errors, would very probably show in this estimate; a smaller one might not. Theorem 1.1 leaves a margin of more than $8 \cdot 10^{-5}$ above the bound for $k = 9$.

6. REMARKS

6.1. **What limits the bound.** The shortest-path value $\mathbb{E} \min_\pi L_\pi/(kh)$, below which no rule can go, is known to us only from simulation (Tables 5.1 and 6.1). The bounds of Theorem 5.1 exceed the simulated values for the same k and h (Table 5.1) by amounts that grow with k, from 0.0046 for $k = 5$ to 0.0077 for $k = 9$. By the controls of Section 5.3 the mesh, the truncation and the rounding account for less than 10^{-5} of each difference. The rest is the price of choosing the order from the cell and not from the block itself.

Finer cells can reduce this price. For $k = 5$ and $h = 2$, the grid of [6] has 12 gap cells and 20 height cells, all equally likely. With a program of our own we simulated the rule that the second-moment bound for the segments defines on these cells, for $4 \cdot 10^6$ blocks: it had the

value 0.87467 ± 0.00012 and lost 0.000493 ± 0.000001 against the shortest path in the same blocks. (The shortest-path value reported in [6] is 0.87426, from a sample of $5 \cdot 10^6$ blocks; this agrees with the last column of Table 5.1 for $k = 5$.) On these cells, therefore, almost all of the difference between 0.8810 and the shortest-path value comes from the second-moment bound for the segments; in our computation for $k = 5$ almost all of the difference between the bound 0.8789 and the shortest-path value comes from the coarser cells.

TABLE 6.1. Simulated shortest-path values $\mathbb{E}\min_\pi L_\pi/(kh)$: the smallest bound that Theorem 2.2 can give for blocks with k gaps and strips of height h. For each k the table gives the strip height with the smallest value among the multiples of 1/10 that we tried, and that value, estimated from $4 \cdot 10^7$ blocks (the same blocks at all strip heights) with a standard error of at most 0.00004.

k	5	6	7	8	9	10
h	2	21/10	11/5	11/5	23/10	23/10
Value	0.8743	0.8671	0.8618	0.8574	0.8540	0.8509

Larger blocks help slowly. In Table 6.1 each further point lowers the shortest-path value by less than the one before, from about 0.007 between $k = 5$ and $k = 6$ to about 0.003 between $k = 9$ and $k = 10$, while the number of cells grows by the factor $mn = 24$ each time. That the gain is limited has been remarked before [19, Section 2.5], [9, Sections 3 and 5]; a figure in the software repositories of [3, 20] shows shortest-path values for $k = 3, \dots, 9$ and six strip heights.[5]

For $k = 6, \dots, 9$ the strip heights of Table 6.1 are higher by 1/10 than those of Theorem 5.1, which were chosen for our rule and not for the shortest path; the minimum over the strip heights is flat, and for $k = 7$ and $k = 9$ the next lower height gives a value larger by less than 0.0001.

6.2. **Two questions.**

(1) For $k = 4$, the partition of [20] is found by a decision tree, with one order for each part. Does such a partition, in place of a product grid, come close to the shortest-path value for k between 7 and 9 at a feasible cost? The bound of Proposition 3.5 extends to cells that are products of arbitrary intervals with endpoints on the mesh.
(2) When a point near the boundary of a strip may be visited from the adjacent strip, simulations give about 0.850 for $k = 8$ and $h = 2$ [9, Table 4]. Can the bound of [9] for such tours be evaluated rigorously for $k \geq 5$?

USE OF AI TOOLS

The author made significant use of Claude Opus 5.5, GPT-6 Pro and GPT-6.1 Sol in developing the method, writing the code and the Lean formalisation, and drafting this paper. The author reviewed the content, edited it as needed, and takes full responsibility for it.

[5] https://github.com/INFORMSJoC/2024.0538 and https://github.com/yyu56253/BHH_upper_bound, accessed 9 October 2026.

References

1. Jillian Beardwood, J. H. Halton, and J. M. Hammersley, *The shortest path through many points*, Proceedings of the Cambridge Philosophical Society **55** (1959), no. 4, 299–327.
2. Richard Bellman, *Dynamic programming treatment of the travelling salesman problem*, Journal of the ACM **9** (1962), no. 1, 61–63.
3. John Gunnar Carlsson and Julien Yu, *A new upper bound for the Euclidean TSP constant*, INFORMS Journal on Computing **38** (2026), no. 2, 341–356.
4. Damek Davis, Paata Ivanisvili, Terence Tao, and contributors, *Optimization constants in mathematics*, 2026, https://github.com/teorth/optimizationproblems, constant 12 (page `constants/12a`), accessed 9 October 2026.
5. Leonardo de Moura and Sebastian Ullrich, *The Lean 4 theorem prover and programming language*, Automated Deduction – CADE 28 (André Platzer and Geoff Sutcliffe, eds.), Lecture Notes in Computer Science, vol. 12699, Springer, Cham, 2021, pp. 625–635.
6. Zhuolun Dong and Junyu Cao, *New bounds for the Euclidean TSP constant*, Preprint, arXiv:2610.02809v1, 2 October 2026.
7. H. P. Edmundson, *Bounds on the expectation of a convex function of a random variable*, Paper P-982, The RAND Corporation, 1957.
8. L. Few, *The shortest path and the shortest road through n points*, Mathematika **2** (1955), no. 2, 141–144.
9. Julia Gaudio and Charlie K. Guan, *An improved upper bound for the Euclidean TSP constant using band crossovers*, Operations Research Letters **69** (2026), 107492, arXiv:2602.11250.
10. Julia Gaudio and Patrick Jaillet, *An improved lower bound for the traveling salesman constant*, Operations Research Letters **48** (2020), no. 1, 67–70, arXiv:1907.02390.
11. Michael Held and Richard M. Karp, *A dynamic programming approach to sequencing problems*, Journal of the Society for Industrial and Applied Mathematics **10** (1962), no. 1, 196–210.
12. C. C. Huang, W. T. Ziemba, and A. Ben-Tal, *Bounds on the expectation of a convex function of a random variable: with applications to stochastic programming*, Operations Research **25** (1977), no. 2, 315–325.
13. D. S. Johnson, L. A. McGeoch, and E. E. Rothberg, *Asymptotic experimental analysis for the Held–Karp traveling salesman bound*, Proceedings of the 7th Annual ACM–SIAM Symposium on Discrete Algorithms, 1996, pp. 341–350.
14. Peter Kall and Stein W. Wallace, *Stochastic programming*, John Wiley & Sons, Chichester, 1994, section numbers are cited from the second edition, which the authors distribute.
15. J. F. C. Kingman, *Poisson processes*, Oxford Studies in Probability, vol. 3, Oxford University Press, 1993.
16. Albert Madansky, *Bounds on the expectation of a convex function of a multivariate random variable*, The Annals of Mathematical Statistics **30** (1959), no. 3, 743–746.
17. Allon G. Percus and Olivier C. Martin, *Finite size and dimensional dependence in the Euclidean traveling salesman problem*, Physical Review Letters **76** (1996), no. 8, 1188–1191.
18. J. Michael Steele, *Probability theory and combinatorial optimization*, CBMS-NSF Regional Conference Series in Applied Mathematics, vol. 69, Society for Industrial and Applied Mathematics, Philadelphia, PA, 1997.
19. Stefan Steinerberger, *New bounds for the traveling salesman constant*, Advances in Applied Probability **47** (2015), no. 1, 27–36, arXiv:1311.6338.
20. Julien Yu and John Gunnar Carlsson, *A new upper bound of the Euclidean TSP constant*, Preprint, Optimization Online, posted 17 June 2023, updated 18 October 2023, https://optimization-online.org/?p=23315.

Independent Researcher

Email address: okeefede@tcd.ie