

**MTH 166**

**Lecture-19**

**Separation of variables solution of**  
**Partial Differential Equations**

**Topic:**

Solution of Partial Differential Equations

**Learning Outcomes:**

1. To solve first order PDE using separation of variables solution.
2. To find nature of Heat, Wave and Laplace equation.

**Find the separation of variables solution of following PDE:**

**Problem 1.**  $\frac{\partial u}{\partial x} = \frac{\partial u}{\partial y}$

**Solution.** The given equation is:  $\frac{\partial u}{\partial x} = \frac{\partial u}{\partial y}$  (1)

Let solution be:  $u(x, y) = XY$  (2) where  $X = f(x), Y = g(y)$

$$\Rightarrow \frac{\partial u}{\partial x} = X'Y \quad \text{and} \quad \frac{\partial u}{\partial y} = XY'$$

Equation (1) becomes:  $X'Y = XY'$

$$\Rightarrow \frac{X'}{X} = \frac{Y'}{Y} = k \quad (\text{Say})$$

Taking these pairs one by one

$$\Rightarrow \frac{X'}{X} = k$$

$$\Rightarrow \int \frac{X'}{X} dx = k \int dx$$

$$\Rightarrow \log X = kx + c_1$$

$$\Rightarrow X = e^{(kx+c_1)}$$

$$\text{also } \frac{Y'}{Y} = k$$

$$\Rightarrow \int \frac{Y'}{Y} dy = k \int dy$$

$$\Rightarrow \log Y = ky + c_2$$

$$\Rightarrow Y = e^{(ky+c_2)}$$

Required solution is:  $u(x, y) = XY = e^{(kx+c_1)} e^{(ky+c_2)} = e^{(c_1+c_2)} e^{k(x+y)}$

$$\Rightarrow u(x, y) = Ae^{k(x+y)} \quad \textbf{Answer.} \quad (A = e^{(c_1+c_2)})$$

**Find the separation of variables solution of following PDE:**

**Problem 2.**  $4 \frac{\partial u}{\partial x} + 3 \frac{\partial u}{\partial y} = 0$

**Solution.** The given equation is:  $4 \frac{\partial u}{\partial x} + 3 \frac{\partial u}{\partial y} = 0$  (1)

Let solution be:  $u(x, y) = XY$  (2) where  $X = f(x), Y = g(y)$

$$\Rightarrow \frac{\partial u}{\partial x} = X'Y \quad \text{and} \quad \frac{\partial u}{\partial y} = XY'$$

Equation (1) becomes:  $4X'Y + 3XY' = 0 \Rightarrow 4X'Y = -3XY'$

$$\Rightarrow 4 \frac{X'}{X} = -3 \frac{Y'}{Y} = k \quad (\text{Say})$$

Taking these pairs one by one

$$\Rightarrow 4 \frac{X'}{X} = k$$

$$\Rightarrow \int \frac{X'}{X} dx = \frac{k}{4} \int dx$$

$$\Rightarrow \log X = \frac{k}{4}x + c_1$$

$$\Rightarrow X = e^{\left(\frac{k}{4}x + c_1\right)}$$

$$\text{also } \frac{Y'}{Y} = k$$

$$\Rightarrow \int \frac{Y'}{Y} dy = -\frac{k}{3} \int dy$$

$$\Rightarrow \log Y = -\frac{k}{3}y + c_2$$

$$\Rightarrow Y = e^{\left(-\frac{k}{3}y + c_2\right)}$$

$$\text{Required solution is: } u(x, y) = XY = e^{\left(\frac{k}{4}x + c_1\right)} e^{\left(-\frac{k}{3}y + c_2\right)} = e^{(c_1 + c_2)} e^{k\left(\frac{1}{4}x - \frac{1}{3}y\right)}$$

$$\Rightarrow u(x, y) = A e^{\frac{k}{12}(3x - 4y)} \quad \textbf{Answer.}$$

$$(A = e^{(c_1 + c_2)})$$

**Find the separation of variables solution of following PDE:**

**Problem 3.**  $y \frac{\partial u}{\partial x} + x \frac{\partial u}{\partial y} = 0$

**Solution.** The given equation is:  $y \frac{\partial u}{\partial x} + x \frac{\partial u}{\partial y} = 0$  (1)

Let solution be:  $u(x, y) = XY$  (2) where  $X = f(x), Y = g(y)$

$$\Rightarrow \frac{\partial u}{\partial x} = X'Y \quad \text{and} \quad \frac{\partial u}{\partial y} = XY'$$

Equation (1) becomes:  $yX'Y + xXY' = 0 \Rightarrow yX'Y = -xXY'$

$$\Rightarrow \frac{1}{x} \frac{X'}{X} = -\frac{1}{y} \frac{Y'}{Y} = k \quad (\text{Say})$$

Taking these pairs one by one

$$\Rightarrow \frac{1}{x} \frac{X'}{X} = k$$

$$\Rightarrow \int \frac{X'}{X} dx = k \int x dx$$

$$\Rightarrow \log X = k \frac{x^2}{2} + c_1$$

$$\Rightarrow X = e^{\left(k \frac{x^2}{2} + c_1\right)}$$

$$\text{also } -\frac{1}{y} \frac{Y'}{Y} = k$$

$$\Rightarrow \int \frac{Y'}{Y} dy = -k \int y dy$$

$$\Rightarrow \log Y = -k \frac{y^2}{2} + c_2$$

$$\Rightarrow Y = e^{\left(-k \frac{y^2}{2} + c_2\right)}$$

$$\text{Required solution is: } u(x, y) = XY = e^{\left(k \frac{x^2}{2} + c_1\right)} e^{\left(-k \frac{y^2}{2} + c_2\right)} = e^{(c_1 + c_2)} e^{k\left(\frac{x^2}{2} - \frac{y^2}{2}\right)}$$

$$\Rightarrow u(x, y) = A e^{\frac{k}{2}(x^2 - y^2)} \quad \textbf{Answer.} \quad (A = e^{(c_1 + c_2)})$$

## **Classification/Nature of Partial Differential Equations:**

Let us consider a general second order homogeneous PDE:

$$A \frac{\partial^2 u}{\partial x^2} + B \frac{\partial^2 u}{\partial x \partial y} + C \frac{\partial^2 u}{\partial y^2} + D \frac{\partial u}{\partial x} + E \frac{\partial u}{\partial y} + Fu = 0 \quad \text{where } u = f(x, y)$$

or

$$Au_{xx} + Bu_{xy} + Cu_{yy} + Du_x + Eu_y + Fu = 0 \quad (1)$$

Equation (1) will be:

1. Hyperbolic iff  $(B^2 - 4AC) > 0$
2. Parabolic iff  $(B^2 - 4AC) = 0$
3. Elliptic iff  $(B^2 - 4AC) < 0$

Note: Coefficients of second order partial derivatives only decide nature of PDE.

**Classify the following PDE as Hyperbolic, Parabolic or Elliptic:**

**Problem 1.**  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$  [2D-Laplace Equation]

**Solution.** Given equ. is:  $u_{xx} + u_{yy} = 0$  (1)

Comparing with:  $Au_{xx} + Bu_{xy} + Cu_{yy} + Du_x + Eu_y + Fu = 0$

We have  $A = 1, B = 0, C = 1$

Here  $(B^2 - 4AC) = (0)^2 - 4(1)(1) = -4 < 0$

So, **Laplace** equation (1) is **Elliptic** in nature.

**Problem 2.**  $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$  [1D-Wave Equation];  $c^2$  is Diffusivity constant.

**Solution.** Given equ. is:  $c^2 u_{xx} - u_{tt} = 0$  (1)

Comparing with:  $Au_{xx} + Bu_{xt} + Cu_{tt} + Du_x + Eu_t + Fu = 0$

We have  $A = c^2, B = 0, C = -1$

Here  $(B^2 - 4AC) = (0)^2 - 4(c^2)(-1) = 4c^2 > 0$

So, **Wave** equation (1) is **Hyperbolic** in nature.

**Problem 3.**  $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$  [1D-Heat Equation];  $c^2$  is Diffusivity constant.

**Solution.** Given equ. is:  $c^2 u_{xx} - u_t = 0$  (1)

Comparing with:  $Au_{xx} + Bu_{xt} + Cu_{tt} + Du_x + Eu_t + Fu = 0$

We have  $A = c^2, B = 0, C = 0$

Here  $(B^2 - 4AC) = (0)^2 - 4(c^2)(0) = 0$

So, **Heat** equation (1) is **Parabolic** in nature.

**Problem 4.**  $(1 - y) \frac{\partial^2 u}{\partial x^2} + 2x \frac{\partial^2 u}{\partial x \partial y} + (1 + y) \frac{\partial^2 u}{\partial y^2} = 0$

**Solution.** Given equ. is:  $(1 - y)u_{xx} + 2xyu_{xy} + (1 + y)u_{yy} = 0$  (1)

Comparing with:  $Au_{xx} + Bu_{xy} + Cu_{yy} + Du_x + Eu_y + Fu = 0$

We have  $A = (1 - y), B = 2x, C = (1 + y)$

Here  $(B^2 - 4AC) = (2x)^2 - 4(1 - y)(1 + y) = 4(x^2 + y^2 - 1)$

So, equation (1) is:

Hyperbolic iff  $(x^2 + y^2 - 1) > 0$

Parabolic iff  $(x^2 + y^2 - 1) = 0$

Elliptic iff  $(x^2 + y^2 - 1) < 0$



[This Photo](#) by Unknown Author is licensed under [CC BY-NC-ND](#)