

Stat261 - Spring 2018

Assignment #5 - due Thursday, April 5, 2018 in class

The files: Stat261-Assign5-R-2018.pdf and StudentPerformanceData.pdf are required to complete this assignment. The Stat261-Assign5-R-2018.Rmd contains the code that generated Stat261-Assign5-R-2018.pdf. Section numbers in the assignment questions refer to the sections in Stat261-Assign5-R-2018.pdf.

We will use the variables **G3**, **sex**, and **Walc** in this data set. Look up the definitions of these variables in the file StudentPerformanceData.pdf.

1. Using the information (summary and graphs) in Section 1.1, comment on the variables **G3**, **sex**, and **Walc**.
2. Section 1.2 contains an analysis of the first 10 grade observations.
 - (a) Comment on the distribution of these 10 observations given Figures 3 and 4.
 - (b) What is a 95% confidence interval for the mean of the 10 observations?
 - (c) Using the 10 observations, perform a test of the hypothesis that the mean grade is 10. Include a concluding sentence which could be incorporated into a report about this data to your boss. Hint: the required computed quantities are given in this section.
3. We are interested in comparing the grades for boys and girls in Section 1.3.
 - (a) Comment on Figure 5.
 - (b) Assuming that the variances for the boys and girls are equal, perform a test of the hypothesis that the mean grade for boys is the same as the mean grade for girls. Include a concluding sentence which could be incorporated into a report about this data to your boss.
 - (c) Without assuming that the variances for the boys and girls are equal, perform a test of the hypothesis that the mean grade for boys is the same as the mean grade for girls. Include a concluding sentence which could be incorporated into a report about this data to your boss.
 - (d) Were your conclusions for the two tests above the same? Why do you suppose that is?
4. In Section 1.4, we are interested in whether there is a relationship between student grades and **Walc**, weekend alcohol consumption.
 - (a) Comment on Figure 6 and 7.
 - (b) The results from fitting a straight line model to the grades as a function of weekend alcohol consumption are given on page 10. What is the estimated straight line model for this data?
 - (c) Perform a test of the hypothesis that the slope parameter is zero. Include a concluding sentence which could be incorporated into a report about this data to your boss.

- (d) Figure 8 is a qqplot of the residuals from the straight line model fit. Comment on this plot.
- (e) Figure 10 contains the side-by-side boxplots of the grades by **Walc**, with the fitted line drawn on top. Comment on this graph.

BONUS QUESTIONS:

1. (BONUS) Suppose that Y_1, Y_2, \dots, Y_n are independent $N(\alpha, \sigma^2)$. Show that if σ is unknown, the likelihood ratio statistic for testing $H_0 : \alpha = \alpha_0$ is given by:

$$D = n \ln \left[1 + \frac{1}{n-1} T^2 \right], \text{ where}$$

$$T = \frac{\hat{\alpha} - \alpha_0}{s/\sqrt{n}}.$$

2. (BONUS) Testing equality of variances. Consider k independent normal samples of sizes n_1, n_2, \dots, n_k . Measurements from sample i have unknown variance σ_i^2 . Let $s_1^2, s_2^2, \dots, s_k^2$ be the sample variances computed from the sample data which are estimates of $\sigma_1^2, \sigma_2^2, \dots, \sigma_k^2$. Since the measurements are normally distributed, we know that.

$$(n_i - 1)s_i^2/\sigma_i^2 \sim \chi_{(n_i-1)}^2 \text{ for } i = 1, 2, \dots, k.$$

Using the above distribution, the log likelihood for σ_i is therefore:

$$\ell(\sigma_i) = -(n_i - 1) \ln \sigma_i - (n_i - 1)s_i^2/(2\sigma_i^2).$$

- (a) Find the joint log likelihood function of $\sigma_1, \sigma_2, \dots, \sigma_k$ and show that it is maximized for $\hat{\sigma}_i^2 = s_i^2, i = 1, 2, \dots, k$.
- (b) Show that if $\sigma_1 = \sigma_2 = \dots = \sigma_k = \sigma$, then the MLE of σ^2 is given by,

$$s_{pooled}^2 = \left(\sum_{i=1}^k (n_i - 1)s_i^2 \right) / \left(\sum_{i=1}^k (n_i - 1) \right).$$

- (c) Show that the likelihood ratio statistic for testing $H_0 : \sigma_1 = \sigma_2 = \dots = \sigma_k = \sigma$ is given by

$$D = \sum_{i=1}^k (n_i - 1) \ln(s_{pooled}^2/s_i^2)$$