

COUNCIL OF SCIENTIFIC & INDUSTRIAL RESEARCH**Mathematical Science****8. Numerical Analysis**

8.1 Errors

8.2 Interpolation with Equal and Unequal Intervals


8.3 Solution of Algebraic and transcendental Equations

8.4 Solution of system of linear equations

8.5 Numerical Integration

8.6 Numerical solution of Differential equations

8.7 Determination of eigenvalues by power method



teachinns
Text with Technology

8. Numerical Analysis

8.1 Errors

8.1.1 Significant figures/digits: The Significant figures/digits are the digits which are used to represent a number.

Examples (8.1):

- (i) 0, 1, 2, ..., 9 are significant figures.
- (ii) 3.15673, 4.00901 both contain six significant figures.
- (iii) 0.0008 contains only one significant figure which is 8 and all the zeros left to 8 here are used to fix the decimal positions.
- (iv) 6.0001 contain five significant figures.

8.1.2 Exact Numbers (N_E): Exact numbers are those numbers which have no approximation.

Examples (8.2): 3, $\frac{1}{7}$, e, $\sqrt{5}$, π etc

8.1.3 Approximate Numbers (N_A): Approximate numbers are those numbers which can not be represented by finite numbers of digits.

Examples (8.3):

- (i) 1.732 is the approximation of $\sqrt{3}$.
- (ii) 3.142 is the approximation of π .

8.1.4 Error (E): Error of a number is the differences of the exact value and approximate value of a number.

$$\text{Error} = \text{Exact number} - \text{Approximate number}$$

$$\text{i.e. } E = N_E - N_A$$

Examples (8.4): $\sqrt{3} - 1.732$ is the error for $\sqrt{3}$.

Errors are committed two ways: (i) rounding a number to a finite digits (Rounding-off error) (ii) due to calculation (Significant error).

8.1.5 Rounding-off Error: Rounding-off error is the error for discarding all but a pre-decided number of digits.

Rules for rounding-off a number to n-significant figures:

- (i) If the digit at (n+1)-th place is less than 5, discard all the digits after n-th place.
- (ii) If the digit at (n+1)-th place is greater than 5, add one to n-th place and discard all the digits after n-th place.
- (iii) If the digit at (n+1)-th place is exactly 5 and n-th place is even, discard all the digits after n-th place.

- (iv) If the digit at (n+1)-th place is exactly 5 and n-th place is odd, add one to n-th place and discard all the digits after n-th place.

Examples (8.4):

Correct the following numbers up to 4 significant figures.

- (i) $2.356489 \approx 2.356$ ($4 < 5$, so 6 remains same)
- (ii) $3.783763 \approx 3.784$ ($7 > 5$, so add 1 to 3)
- (iii) $5.3485345 \approx 5.348$ (8 is at 4th place even, so 8 remains same)
- (iv) $2.6735674 \approx 2.674$ (3 is at 4th place odd, so add 1 to 3)

8.1.6 Types of Errors: Significant Error (S_a), Absolute (E_a), Relative (E_r) and Relative Percentage Errors (E_p)

- (i) Significant Error (S_a) = Exact Numbers (N_E) - Approximate Numbers (N_A)
i.e. $S_a = N_E - N_A$
- (ii) Absolute Error (E_a) = | Exact Numbers (N_E) - Approximate Numbers (N_A) |
i.e. $E_a = |N_E - N_A|$
- (iii) Relative Error (E_r) = $\frac{\text{Absolute Error } (E_a)}{\text{Exact Numbers } (N_E)}$
i.e. $E_r = \frac{E_a}{N_E}$
- (iv) Relative Percentage Error (E_p) = Relative Error (E_r) \times 100%
i.e. $E_p = E_r \times 100\%$

Example (8.5):

Write the approximate representation of $\frac{1}{3}$ correct up to 4 significant figures and also find (i) Significant Error (S_a), (ii) Absolute Error (E_a), (iii) Relative Error (E_r) and (iv) Relative Percentage Error (E_p).

Solution: $\frac{1}{3} = 0.3333$

- (i) $N_E = 1.2345627$, $N_A = 1.2345584$
Significant Error (S_a) = $N_E - N_A = 0.0000043$ (loses 6 significant digits each N_E and N_A)
- (ii) Absolute Error (E_a) = $|N_E - N_A| = |\frac{1}{3} - 0.3333| = 0.000033$
- (iii) Relative Error (E_r) = $\frac{E_a}{N_E} = \frac{0.000033}{\frac{1}{3}} = 0.000099 \approx 0.0001$
- (iv) Relative Percentage Error (E_p) = $E_r \times 100 = \frac{E_a}{V_T} \times 100 = \frac{|V_T - V_A|}{V_T} \times 100$

Remark:

(i) If a number be rounded up to m decimal places the absolute error $E_a \leq \frac{1}{2} 10^{-m}$

Example (8.6)

$$V_T = 345.26132, \quad V_A = 345.261$$

$$\therefore E_a = 0.00032 \leq \frac{1}{2} \times 10^{-3} = 0.0005$$

(ii) If a number be rounded to n correct significant figures, then the relative error

$$E_r < \frac{1}{k \times 10^{n-1}}, \quad k : \text{first significant digit in the number.}$$

Example (8.7) $V_T = \frac{2}{3}, \quad V_A = 0.6667$

(a) $E_a = |V_T - V_A| = 0.000033$

(b) $E_r = \frac{E_a}{V_T} = 0.0000495 \approx 0.00005$

(c) $E_p = E_r \times 100 = 0.005\%$

$$E_p = 0.000033 < 0.00005$$

$$E_r < \frac{1}{k \times 10^{n-1}} = \frac{1}{6 \times 10^3} = 0.00166 \approx 0.0017 \quad (k = 6, n = 4)$$

8.2 Interpolation with Equal and Unequal Intervals

Let $y = f(x)$ defined in $[a, b]$. Let us consider the consecutive value of x , differing by h as $a = x_0, x_1 = x_0 + h, \dots, x_r = x_0 + r h, \dots, x_n = x_0 + h$

$$y_0 = f(x_0), \quad y_1 = f(x_1), \dots, y_n = f(x_n).$$

x_0, x_1, \dots, x_n are called nodes and y_1, y_2, \dots, y_n are called entries.

8.2.1 Forward differences: -

$$\Delta f(x_0) = f(x_0 + h) - f(x_0) = f(x_1) - f(x_0) = y_1 - y_0 = \Delta y_0$$

$$\Delta f(x_1) = f(x_1 + h) - f(x_1) = f(x_2) - f(x_1) = y_2 - y_1 = \Delta y_1$$

$$\Delta f(x_{n-1}) = f(x_{n-1} + h) - f(x_{n-1}) = f(x_n) - f(x_{n-1}) = y_n - y_{n-1} = \Delta y_{n-1}$$

$$\Delta^2 f(x_0) = \Delta^2 y_0 = y_2 - 2y_1 + y_0$$

$$\Delta^3 f(x_0) = \Delta^3 y_0 = y_3 - 3y_2 + 3y_1 - y_0$$

8.2.2 Backward differences: -

$$\nabla f(x_1) = f(x_1) - f(x_1 \cdot h) = f(x_1) - f(x_0) = y_1 - y_0 = \nabla y_1$$

$$\nabla f(x_n) = f(x_n) - f(x_{n-1}) = y_n - y_{n-1} = \nabla y_n$$

$$\diamond \quad \Delta \cdot \nabla = \Delta - \nabla$$

$$\text{Result: } \Delta^k f(x) = \sum_{i=0}^k (-1)^i \binom{k}{i} \delta [x + (k-i)h]$$

- Fundamental theorem of difference calculus:-**

If $f(x)$ be a polynomial of degree n , then the n^{th} difference of $f(x)$ is constant and $(n+1)th$ difference vanish.

8.2.3 Shift operator E:-

$$E f(x) = f(x+h) \Rightarrow E = \Delta + 1 \quad \text{and} \quad E\Delta = \Delta E$$

(i) Relation between Δ (difference operator) and $D \equiv \frac{d}{du}$ (differential operator)

By Taylor's Theorem:-

$$f(x+h) = f(x) + h f'(x) + \frac{h^2}{2!} f''(x) + \dots$$

$$\text{or, } E f(x) = f(x) + h D f(x) + \frac{h^2}{2!} D^2 f(x) + \dots$$

$$\text{or, } (\Delta + 1)f(x) = \left(1 + hD + \frac{h^2}{2!} D^2 + \dots\right) f(x)$$

$$\text{or, } \Delta + 1 = e^{hD}$$

$$\text{or, } hD = \log(1 + \Delta)$$

$$\text{or, } D = \frac{1}{h} \left[\Delta - \frac{\Delta^2}{2} + \frac{\Delta^3}{3} - \frac{\Delta^4}{4} + \dots \right]$$

(ii) Expression of any value of a function in terms of the leading term and leading difference of a difference table

By Shift operator E , we have –

$$f(x+xh) = E f(x) = (1 + \Delta)f(x)$$

$$= \left[1 + \binom{n}{1} \Delta + \binom{n}{2} \Delta^2 + \dots + \binom{n}{n-1} \Delta^{n-1} + \Delta^n \right] f(x)$$

$$= \left[f(x_0) + \binom{n}{1} \Delta f(x_0) + \binom{n}{2} \Delta^2 f(x_0) + \dots + \binom{n}{n-1} \Delta^{n-1} f(x_0) + \Delta^n f(x_0) \right] f(x)$$

$$= y_0 + \binom{n}{1} \Delta y_0 + \binom{n}{2} \Delta^2 y_0 + \dots + \binom{n}{n-1} \Delta^{n-1} y_0 + \Delta^n y_0$$

8.2.4 Factorial notation:-

$$x^{(n)} = x(x - h)(x - 2h) \dots (x - \overline{x-1} h)$$

$$(i) \Delta^n x^{(n)} = n! h^n \quad \text{and} \quad \Delta^{n+1} x^{(n)} = 0$$

Example (8.8): Find the polynomial $f(x)$, which satisfy the following data and hence find the value of $f(1.5)$.

x	1	2	3	4	5
$f(x)$	4	13	34	73	136

Difference table:

x	$\Delta f(x)$	$\Delta f(x)$	$\Delta^2 f(x)$	$\Delta^3 f(x)$	
1	4				
2	13	9			
3	34	21	12		
4	73	39	18	6	
5	136	63	24	6	(3rd degree polynomial as 3rd order difference is constant)

We know that $f(x + nh) = E^n f(x_0) = (1 + \Delta)^n f(x_0)$

Here $x = 1, h = 1, x + nh = 1 + n = x \quad \therefore x = n + 1$

$$\therefore f(x) = (1 + \Delta)^{x-1} f(1)$$

$$= \left[1 + (x-1)\Delta f(1) + \frac{(x-1)(x-2)}{2!} \Delta^2 f(1) + \frac{(x-1)(x-2)(x-3)}{3!} \Delta^3 f(1) + \dots \right]$$

$$= 1 + 9(x-1) + 12 \frac{(x-1)(x-2)}{2} + 6 \frac{(x-1)(x-2)(x-3)}{3}$$

$$= x^3 + 2x + 1$$

$$\therefore f(1.5) = (1.5)^3 + 2 \times (1.5) + 1 = 7.375$$

8.2.5 Newton's Forward Interpolation Formula:-

Let $f(x)$ is known for $(n + 1)$ distinct equispaced arguments namely

$x_0, x_1, x_2, \dots, x_{n-1}, x_n$ such that $x_r = x_0 + r h$ ($r = 0, 1, \dots, n - 1$), $h =$ setp length.

$$y = f(x) \text{ and } y_j = f(x_j), \quad j = 0, 1, \dots, n, \quad s = \frac{x - x_0}{h}$$

$$f(x) \simeq y_0 + s \Delta y_0 + s(s-1) \frac{\Delta^2 y_0}{2!} + s(s-1)(s-2) \frac{\Delta^3 y_0}{3!} + \dots + s(s-1) \dots (s - \overline{n-1}) \frac{\Delta^n y_0}{n!}$$

Which is known as Newton's Forward Interpolation Formula.

$$\text{Error:- } R_{n+1}(x) = \frac{s(s-1)(s-2) \dots (s-n)}{n+1} h^{n+1} f^{n+1}(\xi)$$

Where $\min\{x_1, x_0, x_n\} < \xi < \max\{x_1, x_0, x_n\}$

$$|R_{n+1}(x)| < 1 \text{ for } x > 1 \text{ and } 0 < s < 1$$

8.2.6 Newton's Backward Interpolation Formula: -

$$s = \frac{x - x_n}{h}$$

$$f(x) = y_n + s \Delta y_n + s(s+1) \frac{\Delta^2 y_n}{2!} + s(s+1) \dots (s+n-1) \frac{\Delta^n y_n}{n!}$$

$$\text{Error:- } R_{n+1}(x) = s(s+1) \dots (s+n) h^{n+1} \frac{f^{n+1}(\xi)}{(n+1)!}$$

Where $\min\{x_0, x, x_n\} < \xi < \max\{x, x_0, x_n\}$

8.2.7 Lagrange's Interpolation Formula: -

Let $y = f(x)$ be defined on $[a, b]$ and is only known for $a = x_0, x_1, \dots, x_n = b$, in general are not equispaced and $y_i = f(x_i)$, $i = 0, 1, 2, \dots, n$

$$f(x) \simeq \sum_{i=0}^n l_i(x) f(x_i)$$

$$\text{Where } l_i(x) = \frac{(x-x_0)(x-x_1) \dots (x-x_{i-1})(x-x_{i+1}) \dots (x-x_n)}{(x_i-x_0)(x_i-x_1) \dots (x_i-x_{i-1})(x_i-x_{i+1}) \dots (x_i-x_n)}, \quad i = 0, 1, \dots, n$$

$$\text{Error:- } R_{n+1}(x) = (x-x_0)(x-x_1) \dots (x-x_n) \frac{f^{n+1}(\xi)}{(n+1)!}, \quad (x_0 < \xi < x_n)$$

8.2.8 Divided Difference:-

Let $y = f(x)$ is known for $x_j (j = 0, 1, \dots, n)$ are not necessarily equispaced. Then the first order divided difference for $x_0, x_1, f(x_0, x_1) = \frac{f(x_1) - f(x_0)}{x_1 - x_0} = \frac{f(x_0) - f(x_1)}{x_0 - x_1} = f(x_1, x_0)$

Second order divided difference for x_0, x_1, x_2

$$f(x_0, x_1, x_2) = \frac{f(x_1, x_2) - f(x_0, x_1)}{x_2 - x_0} = \frac{f(x_0)}{(x_0 - x_1) - (x_0 - x_2)} + \frac{f(x_1)}{(x_1 - x_0) - (x_1 - x_2)} + \frac{f(x_2)}{(x_2 - x_0) - (x_2 - x_1)}$$

n th order divided difference for $x_0, x_1, x_2, \dots, x_n$

$$f(x_0, x_1, x_2, \dots, x_n) = \sum_{i=0}^n \frac{f(x_i)}{(x_i - x_0) \dots (x_i - x_{i-1})(x_i - x_{i+1}) \dots (x_i - x_n)} = \sum_{j=0}^n \frac{f(x_j)}{\prod_{i \neq j} (x_j - x_i)}$$

Some Remarks: -

(i) The n th order divided difference of a polynomial of degree n is constant.

(ii) Divided difference for equispaced arguments:-

$$f(x_0, x_1, x_2, \dots, x_n) = \frac{\Delta^n y_0}{n! h^n}$$

$$\text{Since } f(x_0, x_1) = \frac{f(x_1) - f(x_0)}{x_1 - x_0} = \frac{y_1 - y_0}{h} = \frac{\Delta y_0}{h}$$

$$f(x_0, x_1, x_2) = \frac{f(x_1, x_2) - f(x_0, x_1)}{x_2 - x_0} = \frac{\frac{\Delta y_1}{h} - \frac{\Delta y_0}{h}}{2h} = \frac{\Delta(\Delta y_0)}{2h^2} = \frac{\Delta^2 y_0}{2h^2}$$

(iii) Newton's general divided difference formula:-

Let $y = f(x)$ be known for $x_0, x_1, x_2, \dots, x_n$ not necessarily equispaced. Then the polynomial of degree n through $(x_0, y_0), \dots, (x_n, y_n)$, $y_i = f(x_i), i = 0, 1, \dots, n$ is given by –

$$f(x) \simeq f(x_0) + (x - x_0)f(x_0, x_1) + (x - x_0)(x - x_1)f(x_0, x_1, x_2) + \dots + (x - x_0) \dots (x - x_{n-1})f(x_0, x_1, \dots, x_n)$$

Error:-

$$\begin{aligned} R_{n+1}(x) &= (x - x_0)(x - x_1) \dots (x - x_n) f(x_0, x_1, x_2, \dots, x_n) \\ &= (x - x_0)(x - x_1) \dots (x - x_n) \frac{f^{n+1}(\xi)}{(n+1)!} \end{aligned}$$

8.3 Solution of Algebraic and transcendental Equations: -

Algebraic Equation: (i) If $f(x) = 0$ is a purely polynomial in x .

Transcendental Equation: If $f(x) = 0$ contains trigonometric exponential logarithmic function etc.

Assumptions:

- (i) $f(x)$ is continuous and continuously differentiable upto sufficient x_0 of times.
- (ii) $f(x) = 0$ has no multiple root, i.e., if $f(\alpha) = 0$, then in a neighbourhood of α either $f'(x) > 0$ or $f'(x) < 0$.

8.3.1 Method to find the location of roots:


(a) Method of tabulation:

x	x_0	x_1	x_2	x_3
$f(x)$	$f(x_0)$	$f(x_1)$	$f(x_2)$	$f(x_3)$
Sign	—	—	+	+

$$\Rightarrow f(\alpha) = 0 \text{ and } x_1 < \alpha < x_2$$

(b) Graphical method:

Write $f(x) = 0$ as $\phi(x) = \phi(x)$. Draw $y = \phi_1(x)$ and $y = \phi_2(x)$



$$\Rightarrow 2 < \alpha < 3$$

8.3.2 Bisection Method

We first find an interval $[a_0, b_0]$ such that the given function $f: [a_0, b_0] \rightarrow \mathbf{R}$ (set of real numbers) is (i) continuous on $[a_0, b_0]$ (ii) $f'(x)$ keeps the same sign in $[a_0, b_0]$ and (iii) $f(a_0)f(b_0) < 0$. (These three conditions ensure that the function f in $[a_0, b_0]$ has a unique root). Consider $x_0 = \frac{a_0 + b_0}{2}$.

- (i) If $f(x_0) = 0$ then x_0 is the root of f .
- (ii) If $f(x_0) \neq 0$ then either $f(a_0)f(x_0) < 0$ or $f(x_0)f(b_0) < 0$. If $f(a_0)f(x_0) < 0$,
- (iii) then the root lies in $[a_0, x_0]$ and we rename $[a_0, x_0]$ as $[a_1, b_1]$ or if $f(x_0)f(b_0) < 0$ then the root lies in $[x_0, b_0]$ and we rename $[x_0, b_0]$ as $[a_1, b_1]$

and consider a point $x_1 = \frac{a_1 + b_1}{2}$.

- (i) If $f(x_1) = 0$ then x_1 is the root of f .
- (ii) If $f(x_1) \neq 0$ then either $f(a_1)f(x_1) < 0$ or $f(x_1)f(b_1) < 0$. If $f(a_1)f(x_1) < 0$,

then the root lies in $[a_1, x_1]$ and we rename $[a_1, x_1]$ as $[a_2, b_2]$ or if $f(x_1)f(b_1) < 0$ then the root lies in $[x_1, b_1]$ and we rename $[x_1, b_1]$ as $[a_2, b_2]$

and consider a point $x_2 = \frac{a_2 + b_2}{2}$.

Continuing in this process we get a sequence $\{x_n\}$ of points in $[a_0, b_0]$. Now if

$|x_n - x_{n-1}| < \varepsilon$ (a pre-assigned error) then the root of f will be x_n in $[a_0, b_0]$.

This method is surely convergent.

Tabulation of the bisection method

Suppose $f(a_0) > 0$ and $f(b_0) < 0$

x	$a_n(+ve)$	$b_n(-ve)$	$x_{n+1} = \left(\frac{a_n + b_n}{2}\right)$	$f(x_{n+1})$
0	a_0	b_0	$x_1 \left(= \frac{a_0 + b_0}{2}\right)$	$f(x_1) > 0$ (suppose)
1	$a_1 (= x_1)$	$b_1 (= b_0)$	x_2	$f(x_2) > 0$ (suppose)
2	$a_2 (= x_2)$	$b_2 (= b_0)$	x_3	$f(x_3) > 0$ (suppose)
3	$a_3 (= a_2)$	$b_3 (= x_3)$	x_4	$f(x_3) > 0$ (suppose)

8.3.3 Method of iteration or fixed point iteration:

This method is based on the principle of finding a sequence $\{x\}$ each elements of which successively approximates to a real root α so $f(x) = 0$ in $[a, b]$. We rewrite $f(x) = 0$

as $x = \phi(x)$.

Let $x = x_0 \in [a, b]$ be the initial approximation of α , then we set its first approximation as $x_1 = \phi(x_0)$ and then the successive approximations are $x_{n+1} = \phi(x_n)$, $x = 0, 1, \dots$ (iteration formula.)

Convergence of the method of iteration:

$x = \phi(x)$ is not unique.

By MVT,

$$|\alpha - x_1| = |\phi(\alpha) - \phi(x_0)| = |\alpha - x_0| |\phi'(\xi_1)| \text{ for } x_0 < \xi_1 < \alpha$$

$$|\alpha - x_2| = |\phi(\alpha) - \phi(x_1)| = |\alpha - x_1| |\phi'(\xi_2)| \text{ for } x_1 < \xi_2 < \alpha$$

.

.

.

$$|\alpha - x_{n+1}| = |\phi(\alpha) - \phi(x_n)| = |\alpha - x_n| |\phi'(\xi_{n+1})| \text{ for } x_n < \xi_{n+1} < \alpha$$

$$\text{Thus } |\alpha - x_{n+1}| = |\alpha - x_0| |\phi'(\xi_1)| \dots |\phi'(\xi_{n+1})|$$

$$\text{Assuming } |\phi'(x)| \leq \rho \text{ in } (a \leq x \leq b), |\alpha - x_{n+1}| \leq |\alpha - x_0| \rho^{n+1}$$

$$\therefore \log_{n \rightarrow \infty} |\alpha - x_{n+1}| \leq |\alpha - x_0| \log_{n \rightarrow \infty} \rho^{n+1} \rightarrow 0 \text{ if } \rho < 1 \text{ i.e., } |\phi'(x)| < 1$$

$$\rightarrow 0 \text{ if } \rho > 1 \text{ i.e., } |\phi'(x)| > 1$$

Estimation of error:

$$|\varepsilon_{n+1}| \leq \frac{\rho}{1-\rho} |\varepsilon_n|$$

$$\text{Order} = 1$$



$$[|\varepsilon_{n+1}| \leq \rho |\alpha - x_n|]$$

8.3.4 Newton – Raphson Method: -

This is also an iterative method and it is used to find isolated roots of an equation $f(x) = 0$.

We first find an interval $[a_0, b_0]$ such that

(i) the given function $f: [a_0, b_0] \rightarrow \mathbf{R}$ (set of real numbers) satisfies the condition,

$f(a_0)f(b_0) < 0$ and $f'(x) \neq 0$ in $[a_0, b_0]$ with $f'(x)$ is not very small in $[a_0, b_0]$

(ii) the interval $[a_0, b_0]$ should be very close to the root desire root

(iii) $|f(x)f''(x)| < \{f'(x)\}^2$.

Then we find a sequence $\{x_n\}$ of points in $[a_0, b_0]$ such that $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$. If $|x_{n+1} - x_n| < \varepsilon$ (a pre-assigned error) then the root of f will be x_{n+1} in $[a_0, b_0]$.

Convergence of this method:

Comparing with the iteration method

$$\phi(x) = x - \frac{f(x)}{f'(x)} \text{ and}$$

$$|\phi'(x)| < 1 \Rightarrow |f(x)f'(x)| < \{f'(x)\}^2$$

Estimation of error:-

$$|\varepsilon_{n+1}| = |\alpha - x_{n+1}| = \frac{1}{2} \left| \frac{f''(\alpha)}{f'(\alpha)} \right| |\alpha - x_n|^2$$

Definition (Order of a method):- A method is said to be of order P . If P is the largest number for which \exists a finite number C such that $|x_{n+1} - \alpha| \leq C|x_n - \alpha|^P$ i.e.,

$$|\varepsilon_{n+1}| \leq C|\varepsilon_n|^P, n \rightarrow \alpha.$$

So, the order of Newton – Raphson method is 2.

Note:

- (i) N-R method fails if $f'(x) = 0$ or very small in a neighbourhood of the root.
- (ii) N-R method is faster than iteration method.
- (iii) The initial guess (approximation) must be taken very close to the root otherwise it may diverge.
- (iv) To find q -th root of $R > 0$. Let $x = \sqrt[q]{R} \Rightarrow x^q - R = 0$
but $f(x) = x^q - R = 0$ then $f'(x) = qx^{q-1}$, $x_{n+1} = \frac{(q-1)x_n^q + R}{qx_n^{q-1}}$ ($n = 0, 1, 2, \dots$)


Let x be an approximation of α . Then $x_1 = x_0 + h$ is correct root

$$\Rightarrow f(x_0 + h) = f(x_1) = 0 \Rightarrow f(x_0) +$$

$$hf'(x_0) + \frac{h^2}{2} f''(x_0) + \dots = 0$$

$$\Rightarrow h = -\frac{f(x_0)}{f'(x_0)} (\because h \rightarrow 0, h^n = 0, n \geq 2)$$

$$\therefore x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$



8.3.5 Regula –Falsi Method:

We first find an interval $[x_0, x_1]$ such that the given function $f: [x_0, x_1] \rightarrow \mathbf{R}$ (set of real numbers) satisfies $f(x_0)f(x_1) < 0$. We find a point $x_2 = x_1 - \frac{x_1 - x_0}{f(x_1) - f(x_0)} f(x_0)$.

- (i) If $f(x_2) = 0$ then x_2 is the root of f .
- (ii) If $f(x_2) \neq 0$ then either $f(x_0)f(x_2) < 0$ or $f(x_1)f(x_2) < 0$. If $f(x_0)f(x_2) < 0$ then the root lies in $[x_0, x_2]$ and we rename $[x_0, x_2]$ as $[x_1, x_2]$ or if $f(x_1)f(x_2) < 0$ then the root lies in $[x_1, x_2]$ and we keep the interval $[x_1, x_2]$ as same name $[x_1, x_2]$ and find a point

$$x_3 = x_2 - \frac{x_2 - x_1}{f(x_2) - f(x_1)} f(x_1)$$

- (iii) If $f(x_3) = 0$ then x_3 is the root of f .
- (iv) If $f(x_3) \neq 0$ then either $f(x_1)f(x_3) < 0$ or $f(x_2)f(x_3) < 0$. If $f(x_1)f(x_3) < 0$, then the root lies in $[x_1, x_3]$ and we rename $[x_1, x_3]$ as $[x_2, x_3]$ or if $f(x_2)f(x_3) < 0$ then the root lies in $[x_2, x_3]$ and we keep the interval $[x_2, x_3]$ as same name $[x_2, x_3]$

and find a point $x_4 = x_3 - \frac{x_3 - x_2}{f(x_3) - f(x_2)} f(x_2)$

Continuing in this process we get a sequence $\{x_n\}$ of points in $[x_0, x_1]$. Now if $|x_n - x_{n-1}| < \varepsilon$ (a pre-assigned error) then the root of f will be x_n in $[x_0, x_1]$.

Graphically,


$$\frac{AC}{AQ} = \frac{CB}{BP}$$

$$\Rightarrow AC = \frac{AQ}{BP} \cdot CB = \frac{AQ}{BP} (AB - AC)$$

$$\Rightarrow x_{n+1} - a_n = \frac{|f(a_n)|}{|f(b_n)|} (b_n - a_n - (x_{n+1} - a_n))$$

$$\Rightarrow (x_{n+1} - a_n) \left[1 + \frac{|f(a_n)|}{|f(b_n)|} \right]$$

$$= \frac{|f(a_n)|}{|f(b_n)|} (b_n - a_n)$$



$$= x_{n+1} = a_n + \frac{|f(a_n)|}{|f(a_n)| + |f(b_n)|} (b_n - a_n)$$

$$\blacksquare \quad x_{n+1} = x_n + \frac{f(x_n)(x_n - x_{n-1})}{f(x_n) - f(x_{n-1})}$$

8.4 Solution of system of linear equations: -

8.4.1 Gauss-Elimination Method

Let us consider a system of linear algebraic equation in n unknown.

$$\begin{aligned} a_{11} x_1 + a_{12} x_2 + \cdots + a_{1n} x_n &= b_1 \\ a_{21} x_1 + a_{22} x_2 + \cdots + a_{2n} x_n &= b_2 \\ a_{31} x_1 + a_{32} x_2 + \cdots + a_{3n} x_n &= b_3 \\ &\vdots \\ a_{n1} x_1 + a_{n2} x_2 + \cdots + a_{nn} x_n &= b_n \end{aligned}$$

The above system can be written as $AX = b$, where

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ a_{31} & a_{32} & \cdots & a_{3n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix}, \quad X = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{pmatrix} \text{ and } b = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_n \end{pmatrix}$$

Now making the augmented matrix $(A|b)$ in the following form by elementary row operations, we will get the solutions by back substitution.

$$(A|b) \approx \left(\begin{array}{cccc|c} a'_{11} & a'_{12} & \cdots & a'_{1n} & b'_1 \\ 0 & a'_{22} & \cdots & a'_{2n} & b'_2 \\ 0 & 0 & a'_{3n} & \cdots & a'_{3n} & b'_3 \\ & & \vdots & & \vdots & \\ 0 & 0 & 0 & \cdots & a'_{nn} & b'_n \end{array} \right)$$

$$x_n = \frac{b'_n}{a'_{nn}}$$

$$a'_{nn-2} x_{n-1} + a'_{nn-1} x_n = b'_{n-1} \text{ gives } x_{n-1} \text{ etc.}$$

8.4.2 Gauss-Jacobi Iteration Method

Let us consider a system of linear algebraic equation in n unknown.

$$\begin{aligned} a_{11} x_1 + a_{12} x_2 + \cdots + a_{1n} x_n &= b_1 \\ a_{21} x_1 + a_{22} x_2 + \cdots + a_{2n} x_n &= b_2 \\ a_{31} x_1 + a_{32} x_2 + \cdots + a_{3n} x_n &= b_3 \\ &\vdots \\ a_{n1} x_1 + a_{n2} x_2 + \cdots + a_{nn} x_n &= b_n \end{aligned}$$

Where the system is diagonally dominating i.e. $\sum_{i \neq j}^n |a_{ij}| \leq |a_{ii}|$.

The above system can be written as

$$\begin{aligned} x_1 &= \frac{1}{a_{11}} [b_1 - (a_{12} x_2 + a_{13} x_3 + \cdots + a_{1n} x_n)] \\ x_2 &= \frac{1}{a_{22}} [b_2 - (a_{21} x_1 + a_{23} x_3 + \cdots + a_{2n} x_n)] \\ x_3 &= \frac{1}{a_{33}} [b_3 - (a_{31} x_1 + a_{32} x_2 + \cdots + a_{3n} x_n)] \\ &\vdots \\ x_n &= \frac{1}{a_{nn}} [(b_n - (a_{n1} x_1 + a_{n2} x_2 + \cdots + a_{nn-1} x_{n-1}))] \end{aligned}$$



Text with Technology

This method is an iteration method with some initial guess $x_i^{(0)}$ ($i=1, 2, \dots, n$) and the $k+1$ -th (k is a natural number) iteration is given by

$$\begin{aligned} x_1^{(k+1)} &= \frac{1}{a_{11}} [b_1 - (a_{12} x_2^{(k)} + a_{13} x_3^{(k)} + \cdots + a_{1n} x_n^{(k)})] \\ x_2^{(k+1)} &= \frac{1}{a_{22}} [b_2 - (a_{21} x_1^{(k)} + a_{23} x_3^{(k)} + \cdots + a_{2n} x_n^{(k)})] \\ x_3^{(k+1)} &= \frac{1}{a_{33}} [b_3 - (a_{31} x_1^{(k)} + a_{32} x_2^{(k)} + \cdots + a_{3n} x_n^{(k)})] \\ &\vdots \\ x_n^{(k+1)} &= \frac{1}{a_{nn}} [(b_n - (a_{n1} x_1^{(k)} + a_{n2} x_2^{(k)} + \cdots + a_{nn-1} x_{n-1}^{(k)}))] \end{aligned}$$

Here the iteration depends on given error ($\epsilon > 0$). We stop the iteration if $|x_i^{(k+1)} - x_i^{(k)}| < \epsilon$ ($i=1, 2, \dots, n$) and the solution will be $x_i^{(k+1)}$ ($i=1, 2, \dots, n$).

16

8.5 Numerical Integration

8.5.1 Trapezoidal Rule:

Let f be integrable over the interval $[a, b]$. We divide the interval into n equal subintervals by the points $a = x_0, x_0 + h, x_0 + 2h, x_0 + 3h, \dots, x_0 + (n-1)h, x_0 + nh = x_n = b$ where h is the step length.


Consider $y_r = f(x_r)$ for $r = 0, 1, 2, 3, \dots, n$.

Then

$$\int_a^b f(x) dx \approx \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1})], \quad h = \frac{b-a}{n}$$

$$\text{Error} = -\frac{(b-a)^3}{12n^2} f''(\xi), \quad (a = x_0 < \xi < x_n = b)$$

$$= -\frac{n h^3}{12} f''(\xi)$$



8.5.2 Simpson's One-third Rule: ($n = 2m$)

Let f be integrable over the interval $[a, b]$. We divide the interval into n (even) equal subintervals by the points $a = x_0, x_0 + h, x_0 + 2h, x_0 + 3h, \dots, x_0 + (n-1)h, x_0 + nh = x_n = b$ where h is the step length.

Consider $y_r = f(x_r)$ for $r = 0, 1, 2, 3, \dots, n$.

Then

$$\int_a^b f(x) dx \approx \frac{h}{3} [y_0 + y_n + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$$

$$\text{Error} = -\frac{n h^5}{180} f^{iv}(\xi), \quad (a < \xi < b)$$

8.5.3 Simpson's three-eighth Rule: ($n = 3m$)

Let f be integrable over the interval $[a, b]$. We divide the interval into n (multiple of three) equal subintervals by the points $a = x_0, x_0 + h, x_0 + 2h, x_0 + 3h, \dots, x_0 + (n-1)h, x_0 + nh = x_n = b$ where h is the step length.

Consider $y_r = f(x_r)$ for $r = 0, 1, 2, 3, \dots, n$.

Then

$$\int_a^b f(x) dx \approx \frac{3h}{8} [y_0 + y_n + 2(y_3 + y_6 + y_9 + \dots) + 3(y_1 + y_2 + y_4 + y_5 + \dots)]$$

8.5.4 Weddle's Rule: ($n = 6m$)

Let f be integrable over the interval $[a, b]$. We divide the interval into n (multiple of six) equal subintervals by the points $a = x_0, x_0 + h, x_0 + 2h, x_0 + 3h, \dots, x_0 + (n-1)h, x_0 + nh = x_n = b$ where h is the step length.

Consider $y_r = f(x_r)$ for $r = 0, 1, 2, 3, \dots, n$.

$$\text{Then } \int_a^b f(x)dx = \frac{3h}{10} [(y_0 + y_n) + 5(y_1 + y_5 + y_7 + y_{11} + \dots + y_{n-5} + y_{n-1}) + (y_2 + y_4 + y_8 + y_{10} + \dots + y_{n-4} + y_{n-2}) + 6(y_3 + y_9 + y_{15} + \dots + y_{n-3}) + 2(y_6 + y_{12} + \dots + y_{n-6})]$$

8.6 Numerical solution of Differential equations**(A) Single step Method:****8.6.1 Euler's method:**

$$\frac{dy}{dx} = f(x, y) \text{ with } y(x_0) = y_0$$

$$y_{n+1} = y_n + h f(x_n, y_n), \quad n = 0, 1, 2, \dots, p \text{ where } x_r = x_0 + rh, r = 1, 2, \dots, p$$

8.6.2 Euler's Modified method (Euler-Cauchy Method):

$$y_{n+1}^{(0)} = y_n + hf(x_n, y_n)$$

$$y_{n+1}^{(1)} = y_n + \frac{h}{2} [f(x_n, y_n) + f(x_{n+1}, y_{n+1}^{(0)})]$$

·
·
·

$$y_{n+1}^r = y_n + \frac{h}{2} [f(x_n, y_n) + f(x_{n+1}, y_{n+1}^{r-1})]$$

8.6.3 Picard's Method:

$$\frac{dy}{dx} = f(x, y) \text{ with } y(x_0) = y_0$$

$$\text{Integrating } [x_0, x] \text{ we have } \int_{x_0}^x dy = \int_{x_0}^x f(t, y) dt \text{ or, } y(x) = y(x_0) + \int_{x_0}^x f(t, y) dt$$

$$y^{(n+1)}(x) = y_0 + \int_{x_0}^x f(t, y_t^n) dt \text{ where } y^n(x) = y_0$$

8.6.4 Taylor's series method: -

$$y' = \frac{dy}{dx} = f(x, y) \quad \text{with } y(x_0) = y_0$$

$$y(x) = y(x + h) = y_0 + h y_0' + \frac{h^2}{2!} y_0'' + \frac{h^3}{3!} y_0''' + \dots$$

$$y' = f(x, y)$$

$$y'' = f_x + f_y y' = f_x + f_y f$$

$$\text{Order} = \left| \frac{1}{h} f(h) \right|$$

$$f(h) = \text{error}$$

Example:

$$y_1 = y_0 + h y_0' + \frac{h^2}{2!} y_0'' + f(h)$$

Then

$$\text{Order} = \left| \frac{1}{h} t(h) \right| = O(h^2)$$

8.6.5 Runge – Kutta Method:

(a) Second order Runge – Kutta Method:

$$y' = \frac{dy}{dx} = f(x, y) \quad \text{with } y(x_0) = y_0,$$

$$k_1 = h f(x_n, y_n)$$

$$k_2 = h f(x_n + h, y_n + k_1) \quad \text{Error} = O(h^3) \text{ i.e., order} = 2$$

$$y_{n+1} = y_n + \frac{1}{2} (k_1 + k_2)$$

(b) Fourth order Runge – Kutta Method:

$$y_1' = \frac{dy}{dx} = f(x, y) \quad \text{with } y(x_0) = y_0,$$

$$k_1 = h f(x_n, y_n)$$

$$k_2 = h f\left(x_n + \frac{h}{2}, y_n + \frac{k_1}{2}\right), k_3 = h f\left(x_n + \frac{h}{2}, y_n + \frac{k_2}{2}\right)$$

$$k_4 = h f(x_n + h, y_n + k_3)$$

$$y_{n+1} = y_n + \frac{1}{t} [k_1 + 2(k_2 + k_3) + k_4]$$

(B) Multi – Step Method:

8.6.6 Mid – point Method:

$$\frac{dy}{dx} = f(x, y), \quad \text{with } y(x_0) = y_0$$

$$y_{n+1} = y_n + h y_n' + \frac{h^2}{2!} y_n'' + \frac{h^3}{3!} y_n''' + y''(\xi)$$

$$y_{n-1} = y_n - h y_n' + \frac{h^2}{2!} y_n'' - \frac{h^3}{3!} y_n''' + y''(\xi)$$

$$\Rightarrow y_{n+1} - y_{n-1} = 2h y_n' + t(h) \quad \left(\text{order} = \left| \frac{1}{h} t(h) \right| = O(h^2) \right)$$

$$\Rightarrow y_{n+1} = y_{n-1} + 2h f(x_n, y_n), \quad n = 1, 2, 3, \dots$$

8.6.7 Adams – Bash forth Method:**(i) Order – 1:**

$$y_{n+1} = y_n + h f(x_n, y_n), t(h) = \frac{h}{2} y''(\xi) \text{ [Euler's method]}$$

(ii) Order – 2:

$$y_{n+1} = y_n + \frac{h}{2} [3y'_n - y'_{n-1}], t(h) = \frac{5}{12} h^3 y'''(\xi)$$

(iii) Order – 3:

$$y_{n+1} = y_n + \frac{h}{12} [23y'_n - 16y'_{n-1} + y'_{n-2}], t(h) = \frac{3}{8} h^4 y^{(4)}(\xi)$$

8.6.8 Adams – Moulton Method:**(i) Order – 1:-**

$$y_{n+1} = y_n + h f(x_{n+1}, y_{n+1}), t(h) = -\frac{1}{2} h y''(\xi) \text{ (Backward Euler's Method)}$$

(ii) Order – 2:-

$$y_{n+1} = y_n + \frac{h}{2} [y'_{n+1} + y'_n], t(h) = \frac{h^3}{12} y''(\xi)$$

(iii) Order – 3:-

$$y_{n+1} = y_n + \frac{h}{12} [5y'_{n+1} + 8y'_n - y'_{n-1}], t(h) = -\frac{h^4}{24} y^{(4)}(\xi)$$

8.6.9 Two dimensional Newton – Raphson Method:-

$$f(x, y) = 0, g(x, y) = 0$$

$$J = \begin{bmatrix} f_x & f_y \\ g_x & g_y \end{bmatrix}$$

$$\therefore \begin{pmatrix} x_{k+1} \\ y_{k+1} \end{pmatrix} = \begin{pmatrix} x_k \\ y_k \end{pmatrix} - J^{-1}(x_k, y_k) \cdot \begin{pmatrix} f(x_k, y_k) \\ g(x_k, y_k) \end{pmatrix}.$$

Application:- Finding complex root of $f(z) = 0 \Rightarrow f(z) = u(x, y) + iv(x, y) = 0$

$$\text{Then } \begin{pmatrix} x_{k+1} \\ y_{k+1} \end{pmatrix} = \begin{pmatrix} x_k \\ y_k \end{pmatrix} - J^{-1}(x_k, y_k) \cdot \begin{pmatrix} u(x_k, y_k) \\ v(x_k, y_k) \end{pmatrix}$$

8.7 Determination of eigenvalues by power method:

Let $A = (a_{ij})_{n \times n}$ be a real symmetric matrix and $X_0 (\neq 0)$ be a real n component vector.

Let $X_1 = AX_0$, $X_2 = AX_1$, $X_3 = AX_2, \dots, Y = AX_n = AX$ ($X_n = X$)

$$m_0 = X^T X, m_1 = X^T Y, m_2 = Y^T Y.$$

Then $q = \frac{m_1}{m_0}$ is an approximate eigen values λ of A and if we set $q = \lambda + \varepsilon$ so that ε is the

error of q , then $|\varepsilon| \leq \sqrt{\frac{m_2}{m_0} - q^2}$

Example (8.8)

$$A = \begin{pmatrix} 8 & 1 & 1 \\ 1 & 5 & 2 \\ 1 & 2 & 5 \end{pmatrix}, \text{ choose } X_0 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

$$\text{Then } X_1 = \begin{pmatrix} 10 \\ 8 \\ 8 \end{pmatrix}, X_2 = \begin{pmatrix} 96 \\ 66 \\ 66 \end{pmatrix}, X_3 = \begin{pmatrix} 920 \\ 558 \\ 558 \end{pmatrix}, X_4 = \begin{pmatrix} 8316 \\ 4806 \\ 4806 \end{pmatrix}$$

Let $X = X_3, Y = X_4$ we have $m_0 = X^T X = 1432728$, $m_1 = X^T Y = 12847896$, $m_2 = Y^T Y = 115351128$

$$q = \frac{m_1}{m_0} = 8.967 \text{ and } |\varepsilon| \leq \sqrt{\frac{m_2}{m_0} - q^2} = 0.311$$

$$\Rightarrow q - \varepsilon < \lambda < q + \varepsilon \Rightarrow 8.656 < \lambda < 9.278 \Rightarrow \lambda = 9$$