



24 שעות¹

של הקלה עוצמתית בכאב ובדלקת בכדור אחד ביממה^{2*}



ARCOTIA[®]
(etoricoxib, MSD)

• **THE POWER TO MOVE YOU** •

Therapeutic Indications¹:

For the symptomatic relief of osteoarthritis (OA) (30 mg and 60 mg), rheumatoid arthritis (RA) (90 mg), ankylosing spondylitis (90 mg), and the pain and signs of inflammation associated with acute gouty arthritis (120 mg). For the short-term treatment of moderate pain associated with dental surgery (90 mg). Contraindications: • Hypersensitivity to the active substance or to any of the excipients. • Active peptic ulceration or active gastro-intestinal (GI) bleeding. • Patients who have experienced bronchospasm, acute rhinitis, nasal polyps, angioneurotic oedema, urticaria, or allergic-type reactions after taking acetylsalicylic acid or NSAIDs including COX-2 (cyclooxygenase-2) inhibitors. • Pregnancy and lactation. • Severe hepatic dysfunction (serum albumin <25 g/l or Child-Pugh score ≥10). • Estimated renal creatinine clearance <30 ml/min. • Children and adolescents under 16 years of age. • Inflammatory bowel disease. • Congestive heart failure (NYHA II-IV). • Patients with hypertension whose blood pressure is persistently elevated above 140/90mmHg and has not been adequately controlled. • Established ischemic heart disease, peripheral arterial disease, and/or cerebrovascular disease.

For special warnings, precautions for use and common/clinically significant adverse events, please refer to the Physicians information leaflet

References:

1. עלון לרופא כפי שאושר ע"י משרד הבריאות בישראל.

2. Matsumoto AK, Melian A, Mandel DR, et al; for Etoricoxib Rheumatoid Arthritis Study Group. A randomized, controlled, clinical trial of Etoricoxib in the treatment of rheumatoid arthritis. J Rheumatol. 2002;29(8): 1623-1630.




MSD

Copyright © Merck Sharp & Dohme (Israel-1996) Company Ltd., 2013.
All rights reserved. MUSC-1093234-0000

לפני מתן מרשם יש לעיין בעלון לרופא כפי שאושר ע"י משרד הבריאות
*למצבים המופיעים בהתוויה

יעילות: זמן ממוצע לרה-מדיקציה

Figure 5. Mean time to remedication in painful dental conditions.



Moore RA, Derry S et al. Single dose oral analgesics for acute postoperative pain in adults (Review) published in The Cochrane library 2011, Issue 9