



Curves and Surfaces

CS 432 Interactive Computer Graphics
Prof. David E. Breen
Department of Computer Science

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

1



Objectives

- Introduce types of curves and surfaces
 - Explicit
 - Implicit
 - Parametric
 - Strengths and weaknesses
- Discuss Modeling and Approximations
 - Conditions
 - Stability

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2



Escaping Flatland


- Until now we have worked with flat entities such as lines and flat polygons
 - Fit well with graphics hardware
 - Mathematically simple
- But the world is not composed of flat entities
 - Need curves and curved surfaces
 - May only have need at the application level
 - Implementation can render them approximately with flat primitives

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

3



Modeling with Curves



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4



What Makes a Good Representation?

- There are many ways to represent curves and surfaces
- Want a representation that is
 - Stable
 - Smooth
 - Easy to evaluate
 - Must we interpolate or can we just come close to data?
 - Do we need derivatives?


E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

5



Explicit Representation

- Most familiar form of curve in 2D
$$y=f(x)$$
- Cannot represent all curves
 - Vertical lines
 - Circles
- Extension to 3D
 - $y=f(x), z=g(x)$
 - The form $z=f(x,y)$ defines a surface



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

6



Implicit Representation

- Two dimensional curve(s)
 $g(x,y)=0$
- Much more robust
 - All lines $ax+by+c=0$
 - Circles $x^2+y^2-r^2=0$
- Three dimensions $g(x,y,z)=0$ defines a surface
 - Intersect two surface to get a curve
- In general, we cannot exactly solve for points that satisfy the equation

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

7



Algebraic Surface

$$\sum_i \sum_j \sum_k x^i y^j z^k = 0$$

- Quadric surface $2 \geq i+j+k$
- At most 10 terms
- Can solve intersection with a ray by reducing problem to solving quadratic equation


E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

8



Parametric Curves

- Separate equation for each spatial variable
 $x=x(u)$
 $y=y(u)$
 $z=z(u)$
 $\mathbf{p}(u)=[x(u), y(u), z(u)]^T$
- For $u_{\max} \geq u \geq u_{\min}$ we trace out a curve in two or three dimensions



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

9



Selecting Functions

- Usually we can select “good” functions
 - not unique for a given spatial curve
 - Approximate or interpolate known data
 - Want functions which are easy to evaluate
 - Want functions which are easy to differentiate
 - Computation of normals
 - Connecting pieces (segments)
 - Want functions which are smooth

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

10



Parametric Lines


We can normalize u to be over the interval $(0,1)$

Line connecting two points \mathbf{p}_0 and \mathbf{p}_1

$$\mathbf{p}(u)=(1-u)\mathbf{p}_0+u\mathbf{p}_1$$

Ray from \mathbf{p}_0 in the direction \mathbf{d}

$$\mathbf{p}(u)=\mathbf{p}_0+u\mathbf{d}$$



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

11



Parametric Surfaces

- Surfaces require 2 parameters

$$x=x(u,v)$$


$$y=y(u,v)$$

$$z=z(u,v)$$

$$\mathbf{p}(u,v)=[x(u,v), y(u,v), z(u,v)]^T$$


- Want same properties as curves:

- Smoothness
- Differentiability
- Ease of evaluation



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012


12



Normals


We can differentiate with respect to u and v to obtain the normal at any point p

$$\frac{\partial \mathbf{p}(u,v)}{\partial u} = \begin{bmatrix} \partial x(u,v) / \partial u \\ \partial y(u,v) / \partial u \\ \partial z(u,v) / \partial u \end{bmatrix} \quad \frac{\partial \mathbf{p}(u,v)}{\partial v} = \begin{bmatrix} \partial x(u,v) / \partial v \\ \partial y(u,v) / \partial v \\ \partial z(u,v) / \partial v \end{bmatrix}$$

$$\mathbf{n} = \frac{\partial \mathbf{p}(u,v)}{\partial u} \times \frac{\partial \mathbf{p}(u,v)}{\partial v}$$


E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

1
3



Parametric Planes

point-vector form


$$\mathbf{p}(u,v) = \mathbf{p}_0 + u\mathbf{q} + v\mathbf{r}$$

$$\mathbf{n} = \mathbf{q} \times \mathbf{r}$$

three-point form


$$\mathbf{q} = \mathbf{p}_1 - \mathbf{p}_0$$

$$\mathbf{r} = \mathbf{p}_2 - \mathbf{p}_0$$



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

1
4



Parametric Sphere

$$x(u,v) = r \cos \theta \sin \phi$$

$$y(u,v) = r \sin \theta \sin \phi$$


$$z(u,v) = r \cos \phi$$

$$360 \geq \theta \geq 0$$

$$180 \geq \phi \geq 0$$


θ constant: circles of constant longitude
 ϕ constant: circles of constant latitude

differentiate to show $\mathbf{n} = \mathbf{p}$




E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

1
5




Curve Segments

- After normalizing u , each curve is written $\mathbf{p}(u) = [x(u), y(u), z(u)]^T$, $1 \geq u \geq 0$
- In classical numerical methods, we design a single global curve
- In computer graphics and CAD, it is better to design small connected curve *segments*



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

1
6




Parametric Polynomial Curves

$$x(u) = \sum_{i=0}^N c_{xi} u^i \quad y(u) = \sum_{j=0}^M c_{yj} u^j \quad z(u) = \sum_{k=0}^L c_{zk} u^k$$

- If $N=M=L$, we need to determine $3(N+1)$ coefficients
- Equivalently we need $3(N+1)$ independent conditions
- Noting that the curves for x , y and z are independent, we can define each independently in an identical manner
 - We will use the form $\mathbf{p}(u) = \sum_{k=0}^L c_k u^k$ where \mathbf{p} can be any of x , y , z


E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

1
7



Why Polynomials

- Easy to evaluate
- Continuous and differentiable everywhere
 - Must worry about continuity at join points including continuity of derivatives



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

1
8



Cubic Parametric Polynomials

- $N=M=L=3$, gives balance between ease of evaluation and flexibility in design

$$p(u) = \sum_{k=0}^3 c_k u^k$$

- Four coefficients to determine for each of x , y and z
- Seek four independent conditions for various values of u resulting in 4 equations in 4 unknowns for each of x , y and z
 - Conditions are a mixture of continuity requirements at the join points and conditions for fitting the data

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

1
9



Cubic Polynomial Surfaces

$$p(u,v) = [x(u,v), y(u,v), z(u,v)]^T$$

where

$$p(u,v) = \sum_{i=0}^3 \sum_{j=0}^3 c_{ij} u^i v^j$$

p is any of x , y or z

Need 48 coefficients (3 independent sets of 16) to determine a surface patch

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2
0



Designing Parametric Cubic Curves

CS 432 Interactive Computer Graphics

Prof. David E. Breen

Department of Computer Science

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2
1



Objectives

- Introduce the types of curves
 - Interpolating
 - Hermite
 - Bezier
 - B-spline
- Analyze their performance

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2
2



Matrix-Vector Form

$$p(u) = \sum_{k=0}^3 c_k u^k$$

$$\text{define } \mathbf{c} = \begin{bmatrix} c_0 \\ c_1 \\ c_2 \\ c_3 \end{bmatrix} \quad \mathbf{u} = \begin{bmatrix} 1 \\ u \\ u^2 \\ u^3 \end{bmatrix}$$


$$\text{then } p(u) = \mathbf{u}^T \mathbf{c} = \mathbf{c}^T \mathbf{u}$$

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2
3



Interpolating Curve




Given four data (control) points p_0, p_1, p_2, p_3 determine cubic $p(u)$ which passes through them

Must find c_0, c_1, c_2, c_3

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2
4



Interpolation Equations

apply the interpolating conditions at $u=0, 1/3, 2/3, 1$

$$p_0 = p(0) = c_0$$

$$p_1 = p(1/3) = c_0 + (1/3)c_1 + (1/3)^2c_2 + (1/3)^3c_3$$

$$p_2 = p(2/3) = c_0 + (2/3)c_1 + (2/3)^2c_2 + (2/3)^3c_3$$


$$p_3 = p(1) = c_0 + c_1 + c_2 + c_3$$

or in matrix form with $\mathbf{p} = [p_0 \ p_1 \ p_2 \ p_3]^T$

$$\mathbf{p} = \mathbf{A}\mathbf{c} \quad \mathbf{A} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & \left(\frac{1}{3}\right) & \left(\frac{1}{3}\right)^2 & \left(\frac{1}{3}\right)^3 \\ 1 & \left(\frac{2}{3}\right) & \left(\frac{2}{3}\right)^2 & \left(\frac{2}{3}\right)^3 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2
5



Interpolation Matrix

Solving for \mathbf{c} we find the *interpolation matrix*


$$\mathbf{M}_I = \mathbf{A}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -5.5 & 9 & -4.5 & 1 \\ 9 & -22.5 & 18 & -4.5 \\ -4.5 & 13.5 & -13.5 & 4.5 \end{bmatrix}$$

$$\mathbf{c} = \mathbf{M}_I \mathbf{p}$$


Note that \mathbf{M}_I does not depend on input data and can be used for each segment in x, y , and z

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2
6



Interpolating Multiple Segments




use $\mathbf{p} = [p_0 \ p_1 \ p_2 \ p_3]^T$ use $\mathbf{p} = [p_3 \ p_4 \ p_5 \ p_6]^T$

Get continuity at join points but not continuity of derivatives

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2
7



Blending Functions

Rewriting the equation for $p(u)$

$$p(u) = \mathbf{u}^T \mathbf{c} = \mathbf{u}^T \mathbf{M}_I \mathbf{p} = \mathbf{b}(u)^T \mathbf{p}$$

where $\mathbf{b}(u) = [b_0(u) \ b_1(u) \ b_2(u) \ b_3(u)]^T$ is an array of *blending polynomials* such that

$$p(u) = b_0(u)p_0 + b_1(u)p_1 + b_2(u)p_2 + b_3(u)p_3$$

$$b_0(u) = -4.5(u-1/3)(u-2/3)(u-1)$$


$$b_1(u) = 13.5u(u-2/3)(u-1)$$

$$b_2(u) = -13.5u(u-1/3)(u-1)$$

$$b_3(u) = 4.5u(u-1/3)(u-2/3)$$


E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2
8




Blending Functions

- These functions are not smooth
- Hence the interpolation polynomial is not smooth



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

2
9




Interpolating Patch

$$p(u, v) = \sum_{i=0}^3 \sum_{j=0}^3 c_{ij} u^i v^j$$


Need 16 conditions to determine the 16 coefficients c_{ij}

Choose at $u, v = 0, 1/3, 2/3, 1$



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

3
0



Matrix Form

Define $\mathbf{v} = [1 \ v \ v^2 \ v^3]^T$

$\mathbf{C} = [c_{ij}] \quad \mathbf{P} = [p_{ij}]$

$p(u,v) = \mathbf{u}^T \mathbf{C} \mathbf{v}$


If we observe that for constant u (v), we obtain interpolating curve in v (u), we can show

$\mathbf{C} = \mathbf{M}_j \mathbf{P} \mathbf{M}_j^T$

$p(u,v) = \mathbf{u}^T \mathbf{M}_j \mathbf{P} \mathbf{M}_j^T \mathbf{v}$

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

3
1



Blending Patches


$$p(u,v) = \sum_{i=0}^3 \sum_{j=0}^3 b_i(u) b_j(v) p_{ij}$$

Each $b_i(u)b_j(v)$ is a blending patch

Shows that we can build and analyze surfaces from our knowledge of curves

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

3
2




Other Types of Curves and Surfaces


- How can we get around the limitations of the interpolating form
 - Lack of smoothness
 - Discontinuous derivatives at join points
- We have four conditions (for cubics) that we can apply to each segment
 - Use them other than for interpolation
 - Need only come close to the data

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

3
3



Hermite Form




Use two interpolating conditions and two derivative conditions per segment

Ensures continuity and first derivative continuity between segments

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

3
4



Equations

Interpolating conditions are the same at ends

$p(0) = p_0 = c_0$
 $p(1) = p_3 = c_0 + c_1 + c_2 + c_3$


Differentiating we find $p'(u) = c_1 + 2uc_2 + 3u^2c_3$

Evaluating at end points

$p'(0) = p'_0 = c_1$
 $p'(1) = p'_3 = c_1 + 2c_2 + 3c_3$

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

3
5



Matrix Form

$$\mathbf{q} = \begin{bmatrix} p_0 \\ p_3 \\ p'_0 \\ p'_3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 3 \end{bmatrix} \mathbf{c}$$

Solving, we find $\mathbf{c} = \mathbf{M}_H^{-1} \mathbf{q}$ where \mathbf{M}_H is the Hermite matrix

$$\mathbf{M}_H = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -3 & 3 & -2 & -1 \\ 2 & -2 & 1 & 1 \end{bmatrix}$$

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

3
6



Blending Polynomials

$$p(u) = b(u)^T q$$

$$b(u) = \begin{bmatrix} 2u^3 - 3u^2 + 1 \\ -2u^3 + 3u^2 \\ u^3 - 2u^2 + u \\ u^3 - u^2 \end{bmatrix}$$

Although these functions are smooth, the Hermite form is not used directly in Computer Graphics and CAD because we usually have control points but not derivatives

However, the Hermite form is the basis of the Bezier form

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

3
7



Parametric and Geometric Continuity

- We can require the derivatives of x , y , and z to each be continuous at join points (*parametric continuity*)
- Alternately, we can only require that the tangents of the resulting curve be continuous (*geometry continuity*)
- The latter gives more flexibility as we have need satisfy only two conditions rather than three at each join point


E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

3
8



Parametric Continuity

- **Continuity** (recall from the calculus):
 - Two curves are C^i continuous at a point p iff the i -th derivatives of the curves are equal at p




39

Pics/Math courtesy of Dave Mount @ UMD-CP



Example

- Here the p and q have the same tangents at the ends of the segment but different derivatives
- Generate different Hermite curves
- This technique is used in drawing applications



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4
0



Higher Dimensional Approximations

- The techniques for both interpolating and Hermite curves can be used with higher dimensional parametric polynomials
- For interpolating form, the resulting matrix becomes increasingly more ill-conditioned and the resulting curves less smooth and more prone to numerical errors
- In both cases, there is more work in rendering the resulting polynomial curves and surfaces

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4
1



Bezier and Spline Curves and Surfaces

CS 432 Interactive Computer Graphics
Prof. David E. Breen
Department of Computer Science

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4
2



Objectives

- Introduce the Bezier curves and surfaces
- Derive the required matrices
- Introduce the B-spline and compare it to the standard cubic Bezier

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4
3



Bezier's Idea


- In graphics and CAD, we do not usually have derivative data
- Bezier suggested using the same 4 data points as with the cubic interpolating curve to approximate the derivatives in the Hermite form

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4
4



Approximating Derivatives



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4
5



Equations

Interpolating conditions are the same

$$p(0) = p_0 = c_0$$

$$p(1) = p_3 = c_0 + c_1 + c_2 + c_3$$

Approximating derivative conditions

$$p'(0) = 3(p_1 - p_0) = c_0$$

$$p'(1) = 3(p_3 - p_2) = c_1 + 2c_2 + 3c_3$$

Solve four linear equations for $c = M_B p$

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4
6



Bezier Matrix

$$M_B = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -3 & 3 & 0 & 0 \\ 3 & -6 & 3 & 0 \\ -1 & 3 & -3 & 1 \end{bmatrix}$$

$$p(u) = u^T M_B p = b(u)^T p$$

blending functions


E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4
7



Blending Functions

$$b(u) = \begin{bmatrix} (1-u)^3 \\ 3u(1-u)^2 \\ 3u^2(1-u) \\ u^3 \end{bmatrix}$$



Note that all zeros are at 0 and 1 which forces the functions to be smooth over (0,1)

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4
8



Cubic Bezier Curve

- Multiplying it all out gives

$$p(u) = (1-u)^3 p_0 + 3u(1-u)^2 p_1 + 3u^2(1-u) p_2 + u^3 p_3$$

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

4
9



Bernstein Polynomials

- The blending functions are a special case of the Bernstein polynomials

$$b_{kd}(u) = \frac{d!}{k!(d-k)!} u^k (1-u)^{d-k}$$

- These polynomials give the blending polynomials for any degree Bezier form
 - All zeros at 0 and 1
 - For any degree they all sum to 1
 - They are all between 0 and 1 inside (0,1)


E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

5
0



Convex Hull Property

- The properties of the Bernstein polynomials ensure that all Bezier curves lie in the convex hull of their control points
- Hence, even though we do not interpolate all the data, we cannot be too far away



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

5
1



General Form of Bezier Curve

$$\vec{p}(u) = \sum_{i=0}^k \vec{p}_{i+1} \binom{k}{i} (1-u)^{k-i} u^i$$

52



Analysis

- Although the Bezier form is much better than the interpolating form, its derivatives are not continuous at join points
- Can we do better?
 - Go to higher order Bezier
 - More work
 - Derivative continuity still only approximate
 - Supported by OpenGL
 - Apply different conditions
 - Tricky without letting order increase

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

5
4




B-Splines

- Basis splines: use the data at $\mathbf{p} = [p_{i-2} \ p_{i-1} \ p_i \ p_{i+1}]^T$ to define curve only between p_{i-1} and p_i
- Allows us to apply more continuity conditions to each segment
- For cubics, we can have continuity of function, first and second derivatives at join points
- Cost is 3 times as much work for curves
 - Add one new point each time rather than three
- For surfaces, we do 9 times as much work


E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

5
6




Cubic B-spline

$$p(u) = u^T M_s p = b(u)^T p$$

$$M_s = \begin{bmatrix} 1 & 4 & 1 & 0 \\ -3 & 0 & 3 & 0 \\ 3 & -6 & 3 & 0 \\ -1 & 3 & -3 & 1 \end{bmatrix}$$



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012


5
6



Blending Functions

$$b(u) = \frac{1}{6} \begin{bmatrix} (1-u)^3 \\ 4-6u^2+3u^3 \\ 1+3u+3u^2-3u^3 \\ u^3 \end{bmatrix}$$






convex hull property

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

5
7



Splines and Basis


- If we examine the cubic B-spline from the perspective of each control (data) point, each interior point contributes (through the blending functions) to four segments
- We can rewrite $p(u)$ in terms of the data points as

$$p(u) = \sum B_i(u) p_i$$

defining the basis functions $\{B_i(u)\}$

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012


5
9



Basis Functions


In terms of the blending polynomials

$$B_i(u) = \begin{cases} 0 & u < i-2 \\ b_0(u+2) & i-2 \leq u < i-1 \\ b_1(u+1) & i-1 \leq u < i \\ b_2(u) & i \leq u < i+1 \\ b_3(u-1) & i+1 \leq u < i+2 \\ 0 & u \geq i+2 \end{cases}$$



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

6
0




Generalizing Splines

- We can extend to splines of any degree
- Data and conditions to not have to given at equally spaced values (the *knots*)
 - Nonuniform and uniform splines
 - Can have repeated knots
 - Can force spline to interpolate points
- Cox-deBoor recursion gives method of evaluation

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

6
1



NURBS

- Nonuniform Rational B-Spline curves and surfaces add a fourth variable w to x, y, z
 - Can interpret as weight to give more importance to some control data
 - Can also interpret as moving to homogeneous coordinate
- Requires a perspective division
 - NURBS act correctly for perspective viewing
- Quadrics are a special case of NURBS


E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

6
2



Issues with 3D “mesh” formats

- Easy to acquire
- Easy to render
- Harder to model with
- Error prone
 - split faces, holes, gaps, etc




63



BRep Data Structures

- Winged-Edge Data Structure (Weiler)
- Vertex
 - n edges
- Edge
 - 2 vertices
 - 2 faces
- Face
 - m edges



64



BRep Data Structure


- Vertex structure
 - X,Y,Z point
 - Pointers to n coincident edges
- Edge structure
 - 2 pointers to end-point vertices
 - 2 pointers to adjacent faces
 - Pointer to next edge
 - Pointer to previous edge
- Face structure
 - Pointers to m edges

65



Biparametric Surfaces

- Biparametric surfaces
 - A generalization of parametric curves
 - 2 parameters: s, t (or u, v)
 - Two parametric functions



66



Bicubic Surfaces

- Recall the 2D curve: $Q(s) = G \cdot M \cdot S$
 - G : Geometry Matrix
 - M : Basis Matrix
 - S : Polynomial Terms $[s^3 \ s^2 \ s \ 1]$
- For 3D, we allow the points in G to vary in 3D along t as well:


$$Q(s,t) = \begin{bmatrix} G_1(t) & G_2(t) & G_3(t) & G_4(t) \end{bmatrix} \cdot M \cdot S$$

67



Observations About Bicubic Surfaces

- For a fixed t_1 , $Q(s, t_1)$ is a curve
- Gradually incrementing t_1 to t_2 , we get a new curve
- The combination of these curves is a surface
- $G_i(t)$ are 3D curves



68



Bicubic Surfaces

- Each $G_i(t)$ is $G_i(t) = \mathbf{G}_i \cdot \mathbf{M} \cdot \mathbf{T}$, where

$$\mathbf{G}_i = \begin{bmatrix} g_{i1} & g_{i2} & g_{i3} & g_{i4} \end{bmatrix}$$

- Transposing $G_i(t)$, we get

$$G_i(t) = \mathbf{T}^T \cdot \mathbf{M}^T \cdot \mathbf{G}_i^T$$

$$= \mathbf{T}^T \cdot \mathbf{M}^T \cdot \begin{bmatrix} g_{i1} & g_{i2} & g_{i3} & g_{i4} \end{bmatrix}^T$$

69



Bicubic Surfaces

- Substituting $G_i(t)$ into $Q(s) = \mathbf{G} \cdot \mathbf{M} \cdot \mathbf{S}$ we get $Q(s, t)$
- The g_{11} , etc. are the *control points* for the Bicubic surface patch:

$$Q(s, t) = \mathbf{T}^T \cdot \mathbf{M}^T \cdot \begin{bmatrix} g_{11} & g_{21} & g_{31} & g_{41} \\ g_{12} & g_{22} & g_{32} & g_{42} \\ g_{13} & g_{23} & g_{33} & g_{43} \\ g_{14} & g_{24} & g_{34} & g_{44} \end{bmatrix} \cdot \mathbf{M} \cdot \mathbf{S}$$

70



Bicubic Surfaces

- Writing out $Q(s, t) = \mathbf{T}^T \cdot \mathbf{M}^T \cdot \mathbf{G} \cdot \mathbf{M} \cdot \mathbf{S}$ $0 \leq s, t \leq 1$ gives

$$x(s, t) = \mathbf{T}^T \cdot \mathbf{M}^T \cdot G_x \cdot \mathbf{M} \cdot \mathbf{S}$$

$$y(s, t) = \mathbf{T}^T \cdot \mathbf{M}^T \cdot G_y \cdot \mathbf{M} \cdot \mathbf{S}$$


$$z(s, t) = \mathbf{T}^T \cdot \mathbf{M}^T \cdot G_z \cdot \mathbf{M} \cdot \mathbf{S}$$

71



Bézier Patches

- Bézier Surfaces (similar definition)



$$x(s, t) = \mathbf{T}^T \cdot \mathbf{M}_B^T \cdot G_{B_x} \cdot \mathbf{M}_B \cdot \mathbf{S}$$

$$y(s, t) = \mathbf{T}^T \cdot \mathbf{M}_B^T \cdot G_{B_y} \cdot \mathbf{M}_B \cdot \mathbf{S}$$

$$z(s, t) = \mathbf{T}^T \cdot \mathbf{M}_B^T \cdot G_{B_z} \cdot \mathbf{M}_B \cdot \mathbf{S}$$


73



Bezier Patches

Using same data array $\mathbf{P}=[p_{ij}]$ as with interpolating form

$$p(u, v) = \sum_{i=0}^3 \sum_{j=0}^3 b_i(u) b_j(v) p_{ij} = \mathbf{u}^T \mathbf{M}_B \mathbf{P} \mathbf{M}_B^T \mathbf{v}$$




E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

74




Bézier Surfaces

- C^0 and G^0 continuity can be achieved between two patches by setting the 4 boundary control points to be equal
- G^1 continuity achieved when cross-wise CPs are co-linear




75




Bézier Surfaces: Example Utah Teapot

- Utah Teapot modeled with 306 3D control points that define 32 Bézier patches with G^1 continuity



76




Rendering Curves and Surfaces

CS 432 Interactive Computer Graphics
 Prof. David E. Breen
 Department of Computer Science

E. Angel and D Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

7
7




Objectives

- Introduce methods to draw curves
 - Approximate with lines
 - Finite Differences
- Derive the recursive method for evaluation of Bézier curves and surfaces
- Learn how to convert all polynomial data to data for Bézier polynomials

E. Angel and D Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

7
8



Every Curve is a Bézier Curve


- We can render a given polynomial using the recursive method if we find control points for its representation as a Bézier curve
- Suppose that $p(u)$ is given as an interpolating curve with control points q

$$p(u) = u^T M_I q$$
- There exist Bézier control points p such that

$$p(u) = u^T M_B p$$
- Equating and solving, we find $p = M_B^{-1} M_I q$

E. Angel and D Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

7
9



Matrices

Interpolating to Bézier $M_B^{-1} M_I =$


$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ -\frac{5}{6} & 3 & -\frac{3}{2} & \frac{1}{3} \\ \frac{1}{3} & -\frac{3}{2} & 3 & -\frac{5}{6} \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

B-Spline to Bézier $M_B^{-1} M_S =$

$$\begin{bmatrix} 1 & 4 & 1 & 0 \\ 0 & 4 & 2 & 0 \\ 0 & 2 & 4 & 0 \\ 0 & 1 & 4 & 1 \end{bmatrix}$$


E. Angel and D Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

8
0



Example

These three curves were all generated from the same original data using Bézier recursion by converting all control point data to Bézier control points



E. Angel and D Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

8
1



Evaluating Polynomials

- Simplest method to render a polynomial curve is to evaluate the polynomial at many points and form an approximating polyline
- For surfaces we can form an approximating mesh of triangles or quadrilaterals
- Use Horner's method to evaluate polynomials

$$p(u) = c_0 + u(c_1 + u(c_2 + uc_3))$$

- 3 multiplications/evaluation for cubic

E. Angel and D. Shreiner. Interactive Computer Graphics 6E © Addison-Wesley 2012

8



The de Casteljau Algorithm

Basic case, with two points:


- Plotting a curve via *repeated linear interpolation*

- Given $\langle p_0, p_1, \dots \rangle$

a sequence of control points

- Simple case: Mapping a parameter u to the line $\overline{p_0, p_1}$

$$p(u) = (1-u)p_0 + up_1 \quad \text{for } 0 \leq u \leq 1.$$



86
Pics/Math courtesy of Dave Mount @ UMD-CP



The de Casteljau Algorithm


- The complete solution from the algorithm for three iterations:

$$p_{01}(u) = (1-u)p_0 + up_1$$

$$p_{11}(u) = (1-u)p_1 + up_2$$

$$p(u) = (1-u)p_{01}(u) + up_{11}(u)$$

p_0
 p_1 p_{01}
 p_2 p_{11} $p(u)$
Final Value




87
Pics/Math courtesy of Dave Mount @ UMD-CP



The de Casteljau Algorithm

- The solution after four iterations:



88
Pics/Math courtesy of Dave Mount @ UMD-CP



The de Casteljau Algorithm

- Input: $p_0, p_1, p_2, \dots, p_n \in R^3, t \in R$

- Iteratively set:

$$p_{ir}(t) = (1-t)p_{i(r-1)}(t) + tp_{(i+1)(r-1)}(t) \quad \begin{cases} r = 1, \dots, n \\ i = 0, \dots, n-r \end{cases}$$


$$\text{and } p_{i0}(t) = p_i$$

Then $p_{0n}(t)$ is the point with parameter value t on the Bézier curve defined by the p_i 's


89




De Casteljau: Arc Segment Animation



90
Animated by Max Peysakhov @ Drexel University




De Casteljau: Cubic Curve Animation




91

Animated by Max Peysakhov @ Drexel University




Subdivision

- Common in many areas of graphics, CAD, CAGD, vision
- Basic idea
 - primitives defined by control polygons
 - set of control points is not unique
 - more than one way to compute a curve
 - subdivision refines representation of an object by introducing more control points
- Allows for local modification
- Subdivide to pixel resolution



92


Pics/Math courtesy of G. Farin @ ASU



Bézier Curve Subdivision

- Subdivision allows display of curves at different/adaptive levels of resolution
- Rendering systems (OpenGL, ActiveX, etc) only display polygons or lines
- Subdivision generates the lines/facets that approximate the curve/surface
 - output of subdivision sent to renderer

93




deCasteljau Recursion

- We can use the convex hull property of Bezier curves to obtain an efficient recursive method that does not require any function evaluations
 - Uses only the values at the control points
- Based on the idea that “any polynomial and any part of a polynomial is a Bezier polynomial for properly chosen control data”


94

E. Angel and D.Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012



Splitting a Cubic Bezier


p_0, p_1, p_2, p_3 determine a cubic Bezier polynomial and its convex hull



Consider left half $l(u)$ and right half $r(u)$


95

E. Angel and D.Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012




$l(u)$ and $r(u)$

Since $l(u)$ and $r(u)$ are Bezier curves, we should be able to find two sets of control points $\{l_0, l_1, l_2, l_3\}$ and $\{r_0, r_1, r_2, r_3\}$ that determine them



96


E. Angel and D.Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012




Convex Hulls

$\{l_0, l_1, l_2, l_3\}$ and $\{r_0, r_1, r_2, r_3\}$ each have a convex hull that is closer to $p(u)$ than the convex hull of $\{p_0, p_1, p_2, p_3\}$. This is known as the *variation diminishing property*.

The polyline from l_0 to $l_3 (= r_0)$ to r_3 is an approximation to $p(u)$. Repeating recursively we get better approximations.




9
7




Efficient Form

$$\begin{aligned}
 l_0 &= p_0 \\
 r_3 &= p_3 \\
 l_1 &= \frac{1}{2}(p_0 + p_1) \\
 r_1 &= \frac{1}{2}(p_2 + p_3) \\
 l_2 &= \frac{1}{2}(l_1 + \frac{1}{2}(p_1 + p_2)) \\
 r_2 &= \frac{1}{2}(r_1 + \frac{1}{2}(p_2 + p_3)) \\
 l_3 &= r_0 = \frac{1}{2}(l_2 + r_1)
 \end{aligned}$$



Requires only shifts and adds!

9
9




Drawing Parametric Curves

Two basic ways:


- **Iterative evaluation** of $x(t)$, $y(t)$, $z(t)$ for incrementally spaced values of t
 - can't easily control segment lengths and error
- **Recursive Subdivision** via de Casteljau, that stops when control points get sufficiently close to the curve
 - i.e. when the curve is nearly a straight line

100




Drawing Parametric Curves via Recursive Subdivision

- Idea: stop subdivision when segment is flat enough to be drawn w/ straight line
- **Curve Flatness Test:**
 - based on the convex hull
 - if d_2 and d_3 are both less than some ϵ , then the curve is declared flat




9
9




FYI: Computing the Distance from a Point to a Line

- Line is defined with two points
- Basic idea:
 - Project point P onto the line
 - Find the location of the projection



$$d(P, L) = \frac{(y_0 - y_1)x + (x_1 - x_0)y + (x_0y_1 - x_1y_0)}{\sqrt{(x_1 - x_0)^2 + (y_1 - y_0)^2}}$$

102




Drawing Parametric Curves via Recursive Subdivision

The Algorithm:


- DrawCurveRecSub(curve, e)
 - If straight(curve, e) then DrawLine(curve)
 - Else
 - SubdivideCurve(curve, LeftCurve, RightCurve)
 - DrawCurveRecSub(LeftCurve, e)
 - DrawCurveRecSub(RightCurve, e)

103




Biparametric Patch

- (u,v) pair maps to a 3D point on patch



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

1
0
4




Bezier Patches

Using same data array $P=[\vec{p}_{ij}]$ as with interpolating form


$$\vec{p}(u,v) = \sum_{i=0}^3 \sum_{j=0}^3 b_i(u)b_j(v)\vec{p}_{ij} = u^T \mathbf{M}_b \mathbf{P} \mathbf{M}_b^T v$$

Patch lies in convex hull



E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012

1
0
5




Bézier Patches

- Expanding the summation

$$\vec{p}(u,v) = \sum_{i=0}^3 \sum_{j=0}^3 b_i(u)b_j(v)\vec{p}_{ij} =$$


$$\begin{aligned}
 &b_0(u)b_0(v)\vec{p}_{00} + \\
 &b_0(u)b_1(v)\vec{p}_{01} + \\
 &b_0(u)b_2(v)\vec{p}_{02} + \\
 &b_0(u)b_3(v)\vec{p}_{03} + \\
 &b_1(u)b_0(v)\vec{p}_{10} + \\
 &\text{etc.}
 \end{aligned}$$

106



Bezier Blending Functions

$$b(u) = \begin{bmatrix} (1-u)^3 \\ 3u(1-u)^2 \\ 3u^2(1-u) \\ u^3 \end{bmatrix}$$



Note that all zeros are at 0 and 1 which forces the functions to be smooth over (0,1)

E. Angel and D. Shreiner: Interactive Computer Graphics 6E © Addison-Wesley 2012


1
0
7




Faceting



108



Faceting



109