


CGL-LF: $t \in [6, 18]$; 6 time blocks


Local projections of the velocity gradient

$$G_{ij} = \partial_j u_i, \quad P_{ij} = \delta_{ij} - b_i b_j$$




Field-strength-changing strain and compression


$$I_B = S_{\parallel} - \nabla \cdot \mathbf{u} = (D \ln |\mathbf{B}| / Dt)_{\text{ideal}}$$



$$u_{\parallel} = \mathbf{u} \cdot \hat{\mathbf{b}}, \quad \mathbf{u}_{\perp} = \mathbf{u} - u_{\parallel} \hat{\mathbf{b}}$$



Strain, induction proxy, and curvature contribution



$\hat{\mathbf{b}} = \mathbf{B}/|\mathbf{B}|$; $\hat{\mathbf{b}} \cdot \nabla u_{\parallel} - S_{\parallel} = \mathbf{u} \cdot [(\hat{\mathbf{b}} \cdot \nabla) \hat{\mathbf{b}}]$ in the continuum (discrete product-rule residual is retained).

Slope guides are illustrative, not fitted: perpendicular derivatives give $k_{\perp}^2 E_u \propto k_{\perp}^{1/3}$; critical balance gives $k_{\perp}^2 E_u \propto k_{\perp}^{-1/3}$ for $E_u \propto k_{\perp}^{-5/3}$.

Local projections need not follow these slopes. Spectra sum all parallel wavenumbers; the cutoff is not resolved-scale evidence.