

# Bio Final Review

Daniel Filipski and Christopher Milan

May 29, 2018

## 1 General Information

For this, see the google doc.

## 2 Genetics

### 2.1 Sex-linked genes

These are genes located on the sex chromosomes. They will show different phenotype frequencies based on gender.

- (1) Gene A is on the X chromosome. A is the wild type, and  $\alpha$  is the diseased type.  
 $X^A X^\alpha \times X^A Y$ :

$$\begin{array}{c|c} X^A Y & X^\alpha Y \\ \hline X^A X^A & X^\alpha X^A \end{array}$$

### 2.2 Pedigrees

- = Unaffected Female
- = Affected Female
- = Unaffected Male
- = Affected Male

Connecting lines on pedigrees work just as they do on family trees. Relatively simple logic can be used to determine the genotypes of each member of the pedigree; however, some can be more difficult than others. My general method is to use the “method of staring” in the words of Mr. Letarte.

### 2.3 Genetic Disorders – Sickle Cell, Cystic Fibrosis, Huntington’s

#### Sickle Cell

- Red blood cells contain hemoglobin, which bind  $O_2$
- Hemoglobin is made up of two  $\alpha$ -globin and two  $\beta$ -globin polypeptides
- Mutation in  $\beta$ -globin makes it slightly less soluble
- When  $O_2$  is low, hemoglobin without  $O_2$  will start to clump and form long fibers that will change the shape of the red blood cell, which will then get stuck in capillaries
- If one is a heterozygote of this disease, they have an advantage against milleria
- Sickle Cell disease is recessive, because its effects are not great enough with only some of the  $\beta$ -globin broken.

#### Cystic Fibrosis

- In frame three base pair deletion in gene for CFTR
- CFTR is missing one amino acid (phenylalanine), which causes it to misfold and be destroyed
- CFTR is a channel in the epithelial cell membranes for  $Cl^-$

- Without CFTR, there is too much extracellular  $\text{Cl}^-$ , which makes the fluid outside the cell thicker
- Mucus clogs lungs and serves as a growth substance for *Pseudomonas aeruginosa*
- The allele for Cystic Fibrosis is recessive, as with one of the two CFTR, cells still have enough paths for  $\text{Cl}^-$

### **Huntington's Disease**

- Mutation is dominant, but the disease does not present itself until late 30's or early 40's
- Huntingtin gene expressed in nerve cells. Its developmental role in adults is unclear
- CAG (codes for glutamine)
- Wild type 6-35 repeats
- Diseased 36+ repeats
- Diseased protein forms aggregates in neurons, which lead to cell death.

## **2.4 Nondisjunction**

Nondisjunction Event – Failure to separate chromosomes

This is more common in meiosis I. Trisomy 21 causes down syndrome.

## **2.5 Recombinant DNA – Restriction Enzymes, Ligase, Electrophoresis, GFP, PCR, Selectable Markers, Screens, Plasmids, Transformations**

Recombinant DNA – Combination of two or more pieces of DNA to create an artificial construct.

Building Pieces of DNA:

1. Synthesize from scratch
2. Cut and Paste
  - Ligase is not specific and will join any two pieces of DNA.
  - Restriction enzymes originate from bacteria, where they served as a type of immune system.
  - Restriction System: methylase adds  $\text{CH}_3$
  - Restriction Enzyme cuts DNA if not regulated.

Eco RI:

- Sticky ends of *GAATTC* (a DNA reverse palindrome)
- Eco RI cuts between the G and the A

Plasmid:

- Mini chromosome in bacteria
- Must have an ori, a selectable marker
- Plasmids can be shared between cells. They also can be picked up from the environment, when they are there for whatever reason.
- An example of plasmids which can be shared between cells is antibiotic resistance.

DNA Sequencing:

- Denature DNA into single strands
- Add primer for only one strand
- Provide DNA polymerase and the four nucleotides, with a small fraction of the nucleotides modified so that DNA polymerase cannot extend from them (remove 3' hydroxyl)

## 2.6 Selective Breeding – Hybridization vs Inbreeding

Selective breeding – only allowing parents with certain characteristics to breed.

Hybridization:

- Crossing two organisms (typically plants) to get the best traits from both in a hybrid
- Can be different species.

Inbreeding:

- Continued breeding of individuals with similar characteristics
- Dramatically decreases genetic variation
- Increases prominence of some traits

## 3 Evolution

### 3.1 Darwin's Observations and Background Knowledge

<++>

### 3.2 Evolutionary Theory – Adaptations, Fitness, Natural Selection, etc.

<++>

### 3.3 Evidence for Evolution

<++>

### 3.4 Population Genetics – Hardy-Weinberg Equation and Conditions that disrupt equilibrium, selection on polygenic traits

<++>

### 3.5 Speciation

<++>

### 3.6 Molecular Clocks

<++>

### 3.7 Binomial Nomenclature and Linnaean Classification

<++>

### 3.8 Cladograms

<++>

### 3.9 Domains and Kingdoms

<++>

### 3.10 Fossil Record

<++>

### **3.11 Speciation and Extinction Events**

<++>

### **3.12 Origins of Life including Endosymbiont Theory**

<++>