



THE UNIVERSITY OF QUEENSLAND  
A U S T R A L I A

# TORIC VARIETIES

DECLAN FLETCHER

BACHELOR OF MATHEMATICS (HONOURS)

SUPERVISOR: MASOUD KAMGARPOUR  
CO-SUPERVISOR: YANG ZHANG

THE UNIVERSITY OF QUEENSLAND  
SCHOOL OF MATHEMATICS AND PHYSICS

# CONTENTS

1. Introduction	1
1.1. Algebraic geometry	1
1.2. Toric varieties	1
1.3. Geometric invariant theory	1
1.4. Contents of this thesis	1
2. Algebraic subsets of affine space	2
2.1. Affine space and algebraic sets	2
2.2. The map $\mathbf{V}$	3
2.3. The Zariski topology	3
2.4. The map $\mathbf{I}$	4
2.5. The Nullstellensatz	6
2.6. Coordinate rings and regular functions	8
2.7. Products of algebraic sets	9
2.8. Polynomial maps between algebraic subsets	9
2.9. Open subsets of algebraic sets	12
2.10. Regular functions	13
3. Affine varieties	15
3.1. Sheaves and their morphisms	15
3.2. Affine varieties	16
3.3. Tangent spaces	17
4. Convex geometry	22
4.1. Convex cones	22
4.2. Polyhedral cones	26
4.3. The semigroup of a cone	28
5. Affine toric varieties	29
5.1. The semigroup algebra $S_\sigma$	29
5.2. Affine toric varieties	31
5.3. Points of $U_\sigma$	31
5.4. The torus action	31
5.5. Singularities of $U_\sigma$	31
6. Affine GIT quotients as affine toric varieties	32
6.1. Algebraic groups	32
6.2. Reductive groups	32
6.3. The affine GIT quotient	32
6.4. Algebraic tori	35
6.5. The invariant ring for a torus acting on affine space	35
6.6. $\mathbb{A}^n//T$ as a toric variety	37
7. Abstract varieties	41
7.1. Motivation: projective varieties	41
7.2. Abstract varieties	42
7.3. The glueing construction	43
8. Abstract toric varieties	45
8.1. Fans	45
8.2. Abstract toric varieties	46

8.3. The dense torus	46
8.4. The orbit-cone correspondence	46
8.5. Classification of toric surfaces	46
8.6. Toric morphisms	46
References	47

## 1. INTRODUCTION

- 1.1. Algebraic geometry.
- 1.2. Toric varieties.
- 1.3. Geometric invariant theory.
- 1.4. Contents of this thesis.

## 2. ALGEBRAIC SUBSETS OF AFFINE SPACE

In this chapter, we introduce affine varieties, which are the central object of study in this project. Algebraic geometry establishes a connection between spaces defined by zero sets of polynomials (geometric objects), and ideals in a polynomial ring (algebraic objects). We detail this connection, and explain how algebraic properties of rings and ideals inform properties of the corresponding geometric spaces.

**2.1. Affine space and algebraic sets.** Let  $k$  be a field. Affine  $n$ -space over  $k$ , denoted  $\mathbb{A}_k^n$  or  $\mathbb{A}^n$ , is the set

$$\mathbb{A}^n := \{(a_1, \dots, a_n) : a_i \in k\}.$$

Elements of  $\mathbb{A}^n$  are called points, and if  $P = (a_1, \dots, a_n) \in \mathbb{A}^n$  is a point, then the  $a_i$  are called the coordinates of  $P$ .

Let  $A := k[X_1, \dots, X_n]$ . We interpret a polynomial  $f \in A$  as a function on  $\mathbb{A}^n$  by evaluating  $f$  at the coordinates of a point  $P = (a_1, \dots, a_n)$ , i.e.,  $f(P) := f(a_1, \dots, a_n)$ . This allows us to talk about the zeros of the polynomial, which is the set

$$\mathbf{V}(f) := \{P \in \mathbb{A}^n : f(P) = 0\} \subseteq \mathbb{A}^n.$$

More generally, if  $T \subseteq A$  is a set of polynomials, define

$$\mathbf{V}(T) := \{P \in \mathbb{A}^n : f(P) = 0 \text{ for all } f \in T\}.$$

A set  $V \subseteq \mathbb{A}^n$  is called *algebraic* if  $V = \mathbf{V}(T)$  for some  $T \subseteq A$ .

Observe that if  $\langle T \rangle \subseteq A$  is the ideal generated by  $T$ , then  $\mathbf{V}(T) = \mathbf{V}(\langle T \rangle)$ . Moreover, Hilbert's famous Basis theorem tells us all ideals of  $A$  are finitely generated [Rei95, §3.6]. Therefore, if  $f_1, \dots, f_r$  generate  $\langle T \rangle$ , then

$$\mathbf{V}(T) = \mathbf{V}(\langle T \rangle) = \mathbf{V}(\{f_1, \dots, f_r\}).$$

We conclude that any algebraic set is the set of zeros of a *finite* number of polynomials.

**Example 1.** We list some examples of algebraic sets:

- (1)  $\mathbb{A}^n$  and  $\emptyset$  are algebraic, since  $\mathbb{A}^n = \mathbf{V}(0)$  and  $\emptyset = \mathbf{V}(1)$ . Here, by 0 and 1 we mean the constant polynomials in  $A$ .
- (2) Any line in  $\mathbb{A}^2$  has the form  $\mathbf{V}(aX + bY - c)$  for some  $a, b, c \in k$ , so lines are algebraic.
- (3) The parabola  $\mathbf{V}(Y - X^2)$  is an algebraic set.
- (4) The hyperbola  $\mathbf{V}(XY - 1)$  is an algebraic set.
- (5) The twisted cubic  $C = \{(t, t^2, t^3) \in \mathbb{A}^3 : t \in k\}$  is an algebraic set. We see this by noting  $C = \mathbf{V}(\{X^2 - Y, X^3 - Z\})$ .
- (6) The curve  $\mathbf{V}(Y^2 - X^3)$  is algebraic, and it is an example of a so-called *cuspidal cubic*.

**2.2. The map  $\mathbf{V}$ .** Our discussion above tells us that to study zero sets of polynomials, it suffices to study zero sets of ideals in  $A$ . The map

$$\mathbf{V} : \{\text{ideals } I \subseteq A\} \rightarrow \{\text{algebraic subsets } V \subseteq \mathbb{A}^n\}, \quad I \mapsto \mathbf{V}(I),$$

is our first link between algebra and geometry. The following result describes the behaviour of  $\mathbf{V}$ :

**Proposition 2.** (1) If  $I \subseteq J$  are ideals, then  $\mathbf{V}(I) \supseteq \mathbf{V}(J)$ .

(2) If  $I_1, I_2$  are ideals, then  $\mathbf{V}(I_1) \cup \mathbf{V}(I_2) = \mathbf{V}(I_1 I_2)$ .

(3) If  $\{I_\alpha\}_{\alpha \in \mathcal{A}}$  is an arbitrary collection of ideals, then  $\bigcap_{\alpha \in \mathcal{A}} \mathbf{V}(I_\alpha) = \mathbf{V}(\sum_{\alpha \in \mathcal{A}} I_\alpha)$ .<sup>1</sup>

*Proof.* (1) If  $P \in \mathbf{V}(J)$ , then we have  $f(P) = 0$  for all  $f \in I$ , and  $P \in \mathbf{V}(I)$ .

(2) Assume without loss of generality that  $P \in \mathbf{V}(I_1)$ . Then for all  $f \in I_1$  and  $g \in I_2$ , we have  $(fg)(P) = 0$ , implying all polynomials in  $I_1 I_2$  vanish at  $P$ . Conversely, if  $P \in \mathbf{V}(I_1 I_2)$  but  $P \notin \mathbf{V}(I_2)$ , there is  $g \in I_2$  with  $g(P) \neq 0$ . But for any  $f \in I_1$ , there holds  $(fg)(P) = 0$ , so  $f(P) = 0$ .

(3) Suppose  $P \in \bigcap_{\alpha \in \mathcal{A}} \mathbf{V}(I_\alpha)$ . Then for all  $\alpha \in \mathcal{A}$  and all  $f_\alpha \in I_\alpha$ , we have  $f_\alpha(P) = 0$ , implying every element of  $\sum_{\alpha} I_\alpha$  vanishes at  $P$ . Conversely, for each  $\alpha$ , part (1) tells us  $\mathbf{V}(I_\alpha) \supseteq \mathbf{V}(\sum_{\alpha} I_\alpha)$  and so  $\bigcap_{\alpha \in \mathcal{A}} \mathbf{V}(I_\alpha) \supseteq \mathbf{V}(\sum_{\alpha \in \mathcal{A}} I_\alpha)$ .  $\square$

**2.3. The Zariski topology.** Proposition 2 tells us arbitrary intersections and finite unions of algebraic sets are algebraic. In Example 1, we saw  $\emptyset$  and  $\mathbb{A}^n$  are algebraic. Together, these imply algebraic subsets of  $\mathbb{A}^n$  form the closed sets for a topology on  $\mathbb{A}^n$ ; this topology is called the *Zariski topology*.

**Example 3** (The Zariski topology on  $\mathbb{A}^1$ ). *Any nonconstant polynomial in one variable has finitely many roots. Then for any ideal  $I \subseteq A$ ,  $\mathbf{V}(I)$  is either finite or all of  $\mathbb{A}^1$ . In other words, any closed set is either finite or  $\mathbb{A}^1$ , so the Zariski topology on  $\mathbb{A}^1$  is the finite complement topology. When  $k$  is an infinite field, this topology is not Hausdorff; any two nonempty open sets have finite complements, so they must necessarily intersect.*

Example 3 shows us that the Zariski topology is a very coarse topology, in the sense that open sets are large. Nonetheless, the Zariski topology plays an important role in studying algebraic sets.

**2.4. The map  $\mathbf{I}$ .** The map  $\mathbf{V}$  gave us a map from ideals to algebraic subsets; this is our first link between algebra and geometry. There is another map  $\mathbf{I}$ , taking subsets of  $\mathbb{A}^n$  to ideals, defined as

$$\mathbf{I} : \{\text{subsets } V \subseteq \mathbb{A}^n\} \rightarrow \{\text{ideals } I \subseteq A\}, \quad V \mapsto \mathbf{I}(V) := \{f \in A : f(P) = 0 \text{ for all } P \in V\}.$$

In other words,  $\mathbf{I}(V)$  is the ideal of functions vanishing on  $V$ ;  $\mathbf{I}(V)$  is called the ideal of  $V \subseteq \mathbb{A}^n$ . The following result describes the behaviour of the map  $\mathbf{I}$ ;

**Proposition 4.** (1) If  $V \subseteq U \subseteq \mathbb{A}^n$ , then  $\mathbf{I}(V) \supseteq \mathbf{I}(U)$ .

(2) If  $V \subseteq \mathbb{A}^n$ , then  $V \subseteq \mathbf{V}(\mathbf{I}(V))$ , with equality if and only if  $V$  is algebraic.

(3) If  $I \subseteq A$ , then  $I \subseteq \mathbf{I}(\mathbf{V}(I))$ .

*Proof.* (1) If  $f \in \mathbf{I}(U)$ , then we have  $f(P) = 0$  for all  $P \in U$ , so  $f \in \mathbf{I}(V)$ .

(2) If  $P \in V$ , then  $f(P) = 0$  for all  $f \in \mathbf{I}(V)$  and so  $P \in \mathbf{V}(\mathbf{I}(V))$ . If  $V = \mathbf{V}(\mathbf{I}(V))$ , then  $V$  is algebraic by definition. Conversely, if  $V = \mathbf{V}(I)$  is algebraic, then the ideal of functions vanishing on  $V$  will contain  $I$ . Then  $\mathbf{V}(\mathbf{I}(V)) \subseteq \mathbf{V}(I) = V$  and  $V = \mathbf{V}(\mathbf{I}(V))$ .

(3) If  $f \in I$ , then for  $P \in \mathbf{V}(I)$ , we have  $f(P) = 0$ , and so  $f \in \mathbf{I}(\mathbf{V}(I))$ .  $\square$

Proposition 4 begs a question: do  $\mathbf{V}$  and  $\mathbf{I}$  give a bijection between algebraic sets and ideals? Unfortunately, the inclusion  $I \subseteq \mathbf{I}(\mathbf{V}(I))$  may be strict, so  $\mathbf{V}$  and  $\mathbf{I}$  are not always inverses of each other. We give two examples of when this is the case:

<sup>1</sup>Here  $\sum_{\alpha \in \mathcal{A}} I_\alpha = \{\sum_{\alpha \in C} r_\alpha : C \text{ is a finite subset of } \mathcal{A}, r_\alpha \in I_\alpha\}$  is the usual sum of ideals, which is defined even if  $\mathcal{A}$  is infinite.

**Example 5.** (1) Consider  $I = (X^2) \subseteq k[X]$ . Then  $\mathbf{V}(I) = \{0\}$  but  $\mathbf{I}(\mathbf{V}(I)) = (X) \supsetneq I$ .  
(2) Consider  $I = (X^2 + 1)$  as an ideal in  $\mathbb{R}[X]$ . Then since  $X^2 + 1$  never vanishes on  $\mathbb{A}_{\mathbb{R}}^1$ ,  $\mathbf{V}(I) = \emptyset$ , and it holds vacuously that  $\mathbf{I}(\mathbf{V}(I)) = \mathbb{R}[X] \supsetneq I$ .

Example 5 indicates two reasons why  $I \subseteq \mathbf{I}(\mathbf{V}(I))$  may be a strict inclusion: problems can occur when the equations defining an algebraic subset have “unwanted multiplicities,” or when  $k$  is not algebraically closed. In §2.5, we resolve these problems and make the maps  $\mathbf{V}$  and  $\mathbf{I}$  into bijections which are inverses of each other.

For the remainder of this section, we study the basic topological property of irreducibility, and explain how the map  $\mathbf{I}$  gives an algebraic characterisation of this property. We say a nonempty subset  $Y$  of a topological space  $X$  is *reducible* if  $Y = Y_1 \cup Y_2$ , where  $Y_1$  and  $Y_2$  are proper closed subsets of  $Y$  [Har77, Chapter I]. Otherwise, we say  $Y$  is *irreducible*. Then in the context of the Zariski topology, an algebraic set  $V \subseteq \mathbb{A}^n$  is irreducible if it is not a union of proper algebraic subsets.

**Proposition 6.** *Let  $V \subseteq \mathbb{A}^n$  be algebraic. Then  $V$  is irreducible if and only if  $\mathbf{I}(V)$  is prime.*

*Proof.* Suppose  $\mathbf{I}(V)$  is not prime. Then there exist  $f_1, f_2 \notin \mathbf{I}(V)$  such that  $f_1 f_2 \in \mathbf{I}(V)$ . Let  $I_i := (\mathbf{I}(V), f_i)$  for  $i = 1, 2$ . To see  $V$  is reducible, we show that  $V = \mathbf{V}(I_1) \cup \mathbf{V}(I_2)$  and each  $\mathbf{V}(I_i)$  is a strict subset of  $V$ . Since  $I_i \supsetneq \mathbf{I}(V)$ , we have  $\mathbf{V}(I_i) \subsetneq \mathbf{V}(\mathbf{I}(V)) = V$ , with strict inclusion because there is  $P \in V$  with  $f_i(P) \neq 0$ . Then we see  $\mathbf{V}(I_1) \cup \mathbf{V}(I_2) \subseteq V$ . On the other hand, if  $P \in V$ , then  $g(P) = 0$  for all  $g \in \mathbf{I}(V)$ , and also  $(f_1 f_2)(P) = 0$ . Thus,  $f_1(P) = 0$  or  $f_2(P) = 0$ , and  $P \in \mathbf{V}(I_1) \cup \mathbf{V}(I_2)$ .

Conversely, let  $V = V_1 \cup V_2$  be reducible. Since  $V_1, V_2 \neq V$ ,  $\mathbf{I}(V_i) \supsetneq \mathbf{I}(V)$ , and there exists  $f_i \in \mathbf{I}(V_i) \setminus \mathbf{I}(V)$  for  $i = 1, 2$ . But  $(f_1 f_2)(P) = 0$  for all  $P \in V$ , since if  $P \in V_j$ , then  $f_j(P) = 0$ . Thus,  $f_1 f_2 \in \mathbf{I}(V)$  and  $\mathbf{I}(V)$  is not prime.  $\square$

**Example 7.** (1) Let  $k$  be an infinite field. Proposition 6 implies that  $\mathbb{A}^n$  is irreducible, since  $\mathbf{I}(\mathbb{A}^n) = \{0\}$  is a prime ideal. We can also use Example 3 to see  $\mathbb{A}^1$  is irreducible without appealing to Proposition 6. Any proper closed subset of  $\mathbb{A}^1$  is finite, so  $\mathbb{A}^1$  cannot be a union two of proper closed subsets.

(2) Let  $k$  be finite. Since points are closed, a set is irreducible if and only if it is a singleton. In particular,  $\mathbb{A}^n$  is not irreducible in this case.

(3) An example of a reducible algebraic set is  $V = \mathbf{V}(XY) = \mathbf{V}(X) \cup \mathbf{V}(Y)$ , the union of the  $X$ - and  $Y$ -axes. Algebraically, we can see the reducibility of  $V$  since  $\mathbf{I}(V) = (XY)$  is not prime ( $X$  and  $Y$  do not lie in  $(XY)$ , but  $XY$  lies in  $(XY)$ ).

**2.5. The Nullstellensatz.** Our goal in this section is to upgrade the maps  $\mathbf{V}$  and  $\mathbf{I}$  to a bijection between algebraic sets and a particular class of ideals. This is achieved by Hilbert’s Nullstellensatz (Theorem 10). To state and prove the theorem, we need the following definition:

**Definition 8.** *Let  $I$  be an ideal of  $A$ . The radical of  $I$ , denoted  $\sqrt{I}$ , is defined as*

$$\sqrt{I} := \{f \in A : f^n \in I \text{ for some } n \in \mathbb{Z}_{>0}\}.$$

*We say an ideal is radical if  $I = \sqrt{I}$ .*

Observe that  $I \subseteq \sqrt{I}$  for any ideal  $I$ . We claim that prime ideals are radical. If  $I$  is prime and  $f \in \sqrt{I}$ , then  $f^n \in I$  for some  $n \in \mathbb{Z}_{>0}$ , which implies  $f \in I$  since  $I$  is prime.

To prove Theorem 10, we use the following fact from algebra without proof:

**Theorem 9** ([Rei88, §3.8]). *Let  $k$  be an infinite field, and  $B = k[a_1, \dots, a_n]$  a finitely generated  $k$ -algebra. If  $B$  is a field, then  $B$  is algebraic over  $k$ .*

**Theorem 10** (Hilbert's Nullstellensatz [Rei88, §3.10]). *Let  $k$  be an algebraically closed field.*

- (1) *Every maximal ideal of  $A = k[X_1, \dots, X_n]$  is of the form  $\mathfrak{m}_P = (X_1 - a_1, \dots, X_n - a_n)$  for some  $P = (a_1, \dots, a_n) \in \mathbb{A}^n$ .*
- (2) *If  $I$  is a proper ideal of  $A$ , then  $\mathbf{V}(I) \neq \emptyset$ .*
- (3) *For any ideal  $I$ ,  $\mathbf{I}(\mathbf{V}(I)) = \sqrt{I}$ .*

*Proof.* (1) Let  $\mathfrak{m} \subseteq A$  be a maximal ideal. Denote  $K := k[X_1, \dots, X_n]/\mathfrak{m}$ , and let  $\varphi$  be the composition of the natural inclusion and quotient maps

$$\varphi: k \xhookrightarrow{\iota} k[X_1, \dots, X_n] \xrightarrow{\pi} K.$$

Since  $K$  is a field and finitely generated by  $\pi(X_1), \dots, \pi(X_n)$  as a  $k$ -algebra, Theorem 9 implies  $K$  is algebraic over  $k$ . Then  $K/k$  is an algebraic field extension and  $\varphi$  is the inclusion of  $k$  into  $K$ ; since  $k$  is algebraically closed,  $\varphi$  is an isomorphism. For each  $i$ , let  $a_i = (\varphi^{-1} \circ \pi)(X_i)$ , and set  $P = (a_1, \dots, a_n)$ . Then  $\pi(X_i - a_i) = 0$  and  $\mathfrak{m}_P = (X_1 - a_1, \dots, X_n - a_n) \subseteq \ker \pi = \mathfrak{m}$ . But the map  $k[X_1, \dots, X_n] \rightarrow k$  defined by evaluation at  $P$  induces the isomorphism  $k[X_1, \dots, X_n]/\mathfrak{m}_P \cong k$ . Therefore  $\mathfrak{m}_P$  is maximal and  $\mathfrak{m}_P = \mathfrak{m}$ .

(2) Proper ideals are contained in some maximal ideal, so  $I \subseteq \mathfrak{m}_P$  for some  $P \in \mathbb{A}^n$ . Then  $\mathbf{V}(I) \supseteq \mathbf{V}(\mathfrak{m}_P) = \{P\}$  and  $\mathbf{V}(I) \neq \emptyset$ .

(3) Let  $I$  be any ideal in  $A = k[X_1, \dots, X_n]$  and let  $f \in A$  be arbitrary. We introduce a new variable  $Y$  and define the new ideal

$$\tilde{I} := (I, fY - 1) \subseteq k[X_1, \dots, X_n, Y].$$

Intuitively,  $\mathbf{V}(\tilde{I}) \subseteq \mathbb{A}^{n+1}$  is the set of points  $P \in \mathbf{V}(I)$  with  $f(P) \neq 0$ . Specifically, if  $Q = (a_1, \dots, a_n, b) \in \mathbf{V}(\tilde{I})$ , then  $g(a_1, \dots, a_n) = 0$  for all  $g \in \mathbf{V}(I)$  and  $f(a_1, \dots, a_n) \cdot b = 1$  (i.e.,  $f(a_1, \dots, a_n) \neq 0$ ). Now assume that  $f \in \mathbf{I}(\mathbf{V}(I))$  so that  $f(P) = 0$  for all  $P \in \mathbf{V}(I)$ ; our previous discussion implies  $\mathbf{V}(\tilde{I}) = \emptyset$ . Then  $\tilde{I} = A$  by part (2). In particular,  $1 \in \tilde{I}$ , so there exist  $f_i \in I$  and  $g_0, g_i \in k[X_1, \dots, X_n, Y]$  such that

$$1 = \sum g_i f_i + g_0(fY - 1)$$

as a polynomial in  $k[X_1, \dots, X_n, Y]$ . Evaluating the above expression at  $Y = \frac{1}{f}$  yields

$$1 = \sum g_i(X_1, \dots, X_n, 1/f) f_i(X_1, \dots, X_n).$$

Each term in the sum is a rational function where the denominator is a power of  $f$ . Thus there is some  $N \in \mathbb{Z}_{>0}$  such that

$$f^N = \sum f^N g_i(X_1, \dots, X_n, 1/f) f_i(X_1, \dots, X_n)$$

lies in  $k[X_1, \dots, X_n]$ , and in particular, lies in  $I$ . So  $f \in \sqrt{I}$ , proving  $\sqrt{I} \supseteq \mathbf{I}(\mathbf{V}(I))$ . If  $f \in \sqrt{I}$ ,  $f^N \in I \subseteq \mathbf{I}(\mathbf{V}(I))$  for some  $N \in \mathbb{Z}_{>0}$ . But then for any  $P \in \mathbf{V}(I)$ , we must have  $f(P) = 0$ , so  $f \in \mathbf{I}(\mathbf{V}(I))$ .  $\square$

**Corollary 11.** *The maps*

$$\{\text{ideals } I \subseteq A\} \xrightleftharpoons[\mathbf{I}]{\mathbf{V}} \{\text{subsets } V \subseteq \mathbb{A}^n\}$$



induce bijections:

$$\begin{aligned} \{\text{radical ideals}\} &\longleftrightarrow \{\text{algebraic subsets}\}, \\ \{\text{prime ideals}\} &\longleftrightarrow \{\text{irreducible subsets}\}, \\ \{\text{maximal ideals}\} &\longleftrightarrow \{\text{points}\}. \end{aligned}$$

**2.6. Coordinate rings and regular functions.** Let  $V \subseteq \mathbb{A}^n$  be an algebraic set. The *coordinate ring* of  $V$  is defined as

$$k[V] := k[X_1, \dots, X_n]/\mathbf{I}(V).$$

This is a finitely generated  $k$ -algebra. In view of Proposition 6, the ring  $k[V]$  is an integral domain if and only if  $V$  is irreducible.

The coordinate ring is also a *reduced*  $k$ -algebra, meaning it has no non-zero nilpotent elements. Since  $\mathbf{I}(V)$  is radical, this follows from the following general fact:

**Proposition 12.** *Let  $I$  be an ideal in a ring  $R$ . Then  $R/I$  is reduced if and only if  $I$  is radical.*

*Proof.* The ring  $R/I$  is reduced if and only if for all  $n \in \mathbb{Z}_{>0}$ ,  $f^n + I = I$  implies  $f + I = I$ . As a statement about elements instead of cosets, this says that  $f^n \in I$  implies  $f \in I$ , which is equivalent to  $I = \sqrt{I}$ .  $\square$

We say a function  $\varphi: V \rightarrow k$  is *regular* if there exists  $f \in k[X_1, \dots, X_n]$  such that  $\varphi = f|_V$ . Two polynomials  $f, g \in k[X_1, \dots, X_n]$  define the same regular function on  $V$  if and only if  $(f - g)(P) = 0$  for all  $P \in V$ , equivalently, if  $f + \mathbf{I}(V) = g + \mathbf{I}(V)$ . Thus, we identify the ring of regular functions on  $V$  with  $k[V]$ .

Let  $\pi: k[X_1, \dots, X_n] \twoheadrightarrow k[V]$  be the quotient map. The correspondence theorem from ring theory tells us that there is a bijection

$$(1) \quad \left\{ \begin{array}{c} \text{ideals of} \\ k[V] = k[X_1, \dots, X_n]/\mathbf{I}(V) \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{ideals of } k[X_1, \dots, X_n] \\ \text{containing } \mathbf{I}(V) \end{array} \right\}.$$

In particular, any ideal of  $k[V]$  is of the form  $J/\mathbf{I}(V)$ , where  $J$  is an ideal of  $k[X_1, \dots, X_n]$  containing  $\mathbf{I}(V)$ . If  $J/\mathbf{I}(V)$  is an ideal in  $k[V]$ , define

$$\mathbf{V}(J/\mathbf{I}(V)) := \{P \in V : f(P) = 0 \text{ for all } f \in J/\mathbf{I}(V)\}.$$

If we think of elements of  $J$  and  $J/\mathbf{I}(V)$  as functions on  $V$ , they are equal as sets (the quotient  $J/\mathbf{I}(V)$  identifies elements of  $J$  if they define the same function). It then follows that

$$\mathbf{V}(J/\mathbf{I}(V)) = \mathbf{V}(J).$$

The following result extends Corollary 11 to a correspondence between ideals in  $k[V]$  and subsets of  $V$ .

**Corollary 13.** *There are bijections:*

$$\begin{aligned} \{\text{radical ideals in } k[V]\} &\longleftrightarrow \{\text{algebraic subsets of } V\}, \\ \{\text{prime ideals in } k[V]\} &\longleftrightarrow \{\text{irreducible subsets of } V\}, \\ \{\text{maximal ideals in } k[V]\} &\longleftrightarrow \{\text{points of } V\}. \end{aligned}$$

*Proof.* The crux of the proof is that whether an ideal is radical, prime or maximal is preserved by the bijection in Equation 1. Algebraic subsets  $W$  contained in  $V$  are in bijection with radical ideals  $\mathbf{I}(W)$  containing  $\mathbf{I}(V)$ . We have that

$$\frac{k[X_1, \dots, X_n]}{\mathbf{I}(W)} \cong \frac{k[X_1, \dots, X_n]/\mathbf{I}(V)}{\mathbf{I}(W)/\mathbf{I}(V)},$$

so in view of Proposition 12,  $\mathbf{I}(W)$  is radical if and only if  $\mathbf{I}(W)/\mathbf{I}(V)$  is. This establishes the first bijection, and the other two are analogous.  $\square$

## 2.7. Products of algebraic sets.

**2.8. Polynomial maps between algebraic subsets.** Throughout this section, let  $V \subseteq \mathbb{A}^n$  and  $W \subseteq \mathbb{A}^m$  be algebraic sets. We write  $X_1, \dots, X_n$  for the coordinates on  $\mathbb{A}^n$  and  $Y_1, \dots, Y_m$  for the coordinates on  $\mathbb{A}^m$ .

**Definition 14.** We say a map  $\varphi : V \rightarrow W$  is polynomial if there exist  $m$  polynomials  $\varphi_1, \dots, \varphi_m \in k[X_1, \dots, X_n]$  such that

$$\varphi(P) = (\varphi_1(P), \dots, \varphi_m(P))$$

for all  $P \in V$ .

We claim a map  $\varphi : V \rightarrow W$  is polynomial if and only if  $Y_j \circ \varphi \in k[V]$  for all  $j$ . If  $\varphi$  is polynomial given by the components  $\varphi_1, \dots, \varphi_m$ ,  $Y_j \circ \varphi = \varphi_j$  is regular. Conversely, if  $\tilde{\varphi}_j := Y_j \circ \varphi \in k[V]$  and  $\varphi_j \in k[X_1, \dots, X_n]$  such that  $\varphi_j \equiv \tilde{\varphi}_j \pmod{\mathbf{I}(V)}$ ,  $\varphi = (\varphi_1, \dots, \varphi_m)$  and  $\varphi$  is polynomial.

We also claim that the composition of polynomial maps is polynomial. Let  $U \subseteq \mathbb{A}^l$  be algebraic, and let  $\varphi : V \rightarrow W$  and  $\psi : W \rightarrow U$  be polynomial maps. If  $\varphi_1, \dots, \varphi_m$  and  $\psi_1, \dots, \psi_l$  are the components of  $\varphi$  and  $\psi$ , respectively, the components of  $\psi \circ \varphi : V \rightarrow U$  are

$$\psi_1(\varphi_1, \dots, \varphi_m), \dots, \psi_l(\varphi_1, \dots, \varphi_m) \in k[X_1, \dots, X_n].$$

We say a polynomial map  $\varphi : V \rightarrow W$  is an *isomorphism* of algebraic sets if there exists a polynomial map  $\psi : W \rightarrow V$  such that  $\psi \circ \varphi = \text{id}_V$  and  $\varphi \circ \psi = \text{id}_W$ .

**Theorem 15.** Let  $V \subseteq \mathbb{A}^n$ ,  $W \subseteq \mathbb{A}^m$  and  $U \subseteq \mathbb{A}^l$  be algebraic sets.

- (1) A polynomial map  $\varphi : V \rightarrow W$  induces a  $k$ -algebra homomorphism  $\varphi^* : k[W] \rightarrow k[V]$ ,  $f \mapsto \varphi^* f := f \circ \varphi$ .
- (2) Any  $k$ -algebra homomorphism  $\Phi : k[W] \rightarrow k[V]$  is of the form  $\Phi = \varphi^*$  for a unique polynomial map  $\varphi : V \rightarrow W$ .
- (3) If  $\varphi : V \rightarrow W$  and  $\psi : W \rightarrow U$  are polynomial maps, then  $(g \circ f)^* = f^* \circ g^*$ .

**Remark 16.** Together, part (1) and (2) says that the map  $\varphi \mapsto \varphi^*$  induces a bijection

$$\left\{ \begin{array}{c} \text{polynomial maps} \\ V \rightarrow W \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} k\text{-algebra homomorphisms} \\ k[W] \rightarrow k[V] \end{array} \right\}.$$

The map  $\varphi^*$  is called the *pullback* of  $\varphi$ .

*Proof.* (1) Since the composition of polynomial maps is polynomial,  $\varphi^* f = f \circ \varphi \in k[V]$  for all  $f \in k[W]$ . For  $f, g \in k[W]$ , we have

$$\varphi^*(f + g) = (f + g) \circ \varphi = f \circ \varphi + g \circ \varphi = \varphi^* f + \varphi^* g, \text{ and}$$

$$\varphi^*(fg) = (fg) \circ \varphi = (f \circ \varphi)(g \circ \varphi) = (\varphi^*f)(\varphi^*g),$$

so  $\varphi^*$  is a  $k$ -algebra homomorphism.

(2) We first show there exists a polynomial map  $\varphi : V \rightarrow W$  with  $\Phi = \varphi^*$ . If  $g \in k[Y_1, \dots, Y_m]$ , we write  $\bar{g}$  for the coset of  $g$  in  $k[W]$ , e.g.,  $\bar{Y}_j = Y_j + \mathbf{I}(W)$ . Let  $\varphi_i := \Phi(\bar{Y}_i) \in k[V]$  for  $i = 1, \dots, m$ , and define the polynomial map  $\varphi : V \rightarrow \mathbb{A}^m$  by

$$\varphi(P) = (\varphi_1(P), \dots, \varphi_m(P)).$$

We need to show  $\varphi(V) \subseteq W$  and  $\varphi^* = \Phi$ . Since  $\Phi$  is a homomorphism, we have for any  $\bar{g} \in k[W]$  that

$$\Phi(\bar{g}) = \Phi(g(\bar{Y}_1, \dots, \bar{Y}_m)) = g(\Phi(\bar{Y}_1), \dots, \Phi(\bar{Y}_m)) = g(\varphi_1, \dots, \varphi_m).$$

Then for any  $v \in V$ ,

$$\Phi(\bar{g})(v) = g(\varphi_1(v), \dots, \varphi_m(v)) = g(\varphi(v)).$$

When  $g \in \mathbf{I}(W)$ ,  $\bar{g} = 0$  and the above equation implies that

$$g(\varphi(v)) = 0$$

for all  $v \in V$ , so  $\varphi(V) \subseteq W$ . To see  $\varphi^* = \Phi$ , note that  $\varphi^*(\bar{Y}_i) = \bar{Y}_i \circ \varphi = \varphi_i = \Phi(\bar{Y}_i)$ . To show the uniqueness of  $\varphi$ , we prove the map  $\varphi \mapsto \varphi^*$  is injective. If  $\varphi, \phi : V \rightarrow W$  are polynomial maps with components  $\varphi_i$  and  $\phi_i$ , respectively, and  $\varphi^* = \phi^*$ , then for each  $i$ ,

$$\varphi_i = \varphi^*(\bar{Y}_i) = \phi^*(\bar{Y}_i) = \phi_i.$$

Therefore,  $\varphi$  and  $\phi$  have the same components and  $\varphi = \phi$ .

(3) Note  $\psi \circ \varphi : V \rightarrow U$ . For any  $f \in k[U]$ , we have

$$(\psi \circ \varphi)^*f = f \circ (\psi \circ \varphi) = (f \circ \psi) \circ \varphi = \varphi^*(\psi^*f),$$

so  $(\psi \circ \varphi)^* = \varphi^* \circ \psi^*$ . □

**Corollary 17.** *A polynomial map  $\varphi : V \rightarrow W$  is an isomorphism of algebraic sets if and only if  $\varphi^* : k[W] \rightarrow k[V]$  is an isomorphism of  $k$ -algebras.*

**Example 18.** *We give examples of polynomial maps between algebraic sets and their pullbacks.*

- (1) Let  $C = \mathbf{V}(\{X^2 - Y, X^3 - Z\})$  be the twisted cubic. Consider the map  $\varphi : \mathbb{A}^1 \rightarrow C$  defined by  $t \mapsto (t, t^2, t^3)$ . Note that  $X \in k[C] = k[X, Y, Z]/(X^2 - Y, X^3 - Z)$  generates  $k[C]$ . We write  $k[T]$  for the coordinate ring of  $\mathbb{A}^1$ . Then the pullback  $\varphi^* : k[C] \rightarrow k[T]$  is given by

$$X \mapsto X \circ \varphi = T.$$

Then  $\varphi^*$  is a  $k$ -algebra isomorphism, and  $C$  and  $\mathbb{A}^1$  are isomorphic as algebraic sets.

- (2) Let  $V = \mathbf{V}(Y^2 - X^3) \subseteq \mathbb{A}^2$ . Consider  $\varphi : \mathbb{A}^1 \rightarrow V$  given by  $t \mapsto (t^2, t^3)$ . Note  $k[V] = k[X, Y]/(Y^2 - X^3)$  is generated by  $X, Y \in k[V]$ . The pullback is  $\varphi^* : k[V] \rightarrow k[T]$ , given by

$$X \mapsto T^2, \quad Y \mapsto T^3.$$

Then  $\varphi^*(k[C]) = k[T^2, T^3] \neq k[T]$ .

- (3) Consider  $\varphi : \mathbb{A}^2 \rightarrow \mathbb{A}^2$ , given by  $(x, y) \mapsto (xy, y)$ . The image is

$$\varphi(\mathbb{A}^2) = \{(x, y) \in \mathbb{A}^2 : x = y = 0 \text{ or } y \neq 0\}.$$

This set is not algebraic, so the image of a polynomial map is not necessarily an algebraic set.

**2.9. Open subsets of algebraic sets.** Let  $V \subseteq \mathbb{A}^n$  be an algebraic set and  $f \in k[V]$  a polynomial function on  $V$ . Then the set

$$V_f := \{P \in V : f(P) \neq 0\}$$

is an open subset of  $V$ , since its complement (the zero set of  $f$ ) is algebraic. It turns out that the set  $\{V_f : f \in k[V]\}$  is a basis for the Zariski topology on  $V$ :

**Proposition 19** ([Mil13, Proposition 2.37]). *The set  $\{V_f : f \in k[V]\}$  is a basis for the Zariski topology on  $V$ . Specifically, every open set is a finite union of the form  $\bigcup V_f$ .*

*Proof.* Every open set  $U \subseteq V$  is the complement of  $\mathbf{V}(J)$  for some ideal  $J$  of  $k[V]$ . If  $J$  is generated by  $f_1, \dots, f_m$ , then  $U = \bigcup V_{f_i}$ .  $\square$

In light of the proposition, sets of the form  $V_f$  are called principal open subsets of  $V$ .

We can think of a principal open subset  $V_f$  as an algebraic set in its own right. Specifically, suppose  $\mathbf{I}(V)$  is generated by  $f_1, \dots, f_m \in k[X_1, \dots, X_n]$ . Let  $g \in k[X_1, \dots, X_n]$  be a coset representative for  $f$  in  $k[V] = k[X_1, \dots, X_n]/\mathbf{I}(V)$ . Writing  $k[X_1, \dots, X_n, Y]$  for the coordinate ring of  $k^n \times k$ , we define a new algebraic set  $W \subseteq k^n \times k$  by

$$W := \mathbf{V}(f_1, \dots, f_m, gY - 1).$$

A point  $(x_1, \dots, x_n, y) \in W$  satisfies  $f_i(x_1, \dots, x_n) = 0$  for all  $i$ , and  $g(x_1, \dots, x_n) = \frac{1}{y} \neq 0$ . It follows that the projection  $k^n \times k \rightarrow k^n$  identifies  $V_f$  with the algebraic set  $W$ .

It is natural to ask what the coordinate ring of  $V_f$  is; we will explain what it is when  $V$  is irreducible. In this case,  $k[V]$  is an integral domain. We write  $k(V)$  for the fraction field  $k(V) := \text{Frac}(k[V])$ , and call  $k(V)$  the field of rational functions on  $V$ . For any  $f \in k[V]$ , consider the subring

$$k[V]_f := \left\{ \frac{g}{f^r} : g \in k[V], r \geq 0 \right\} \subseteq k(V).$$

The ring  $k[V]_f$  is called the localisation of  $k[V]$  at  $f$ .<sup>2</sup> One can prove

$$k[V]_f \cong (k[V])[Y]/(fY - 1),$$

so that  $k[V]_f \cong k[X_1, \dots, X_n]/(f_1, \dots, f_m, gY - 1)$ , and therefore  $k[V]_f$  is the coordinate ring of  $V_f$  [Mil13, Lemma 1.13].

An important example of an algebraic set which arises as a principal open subset of  $\mathbb{A}^n$  is the algebraic torus,

$$(k^\times)^n := \{(x_1, \dots, x_n) \in k^n : x_i \neq 0\}.$$

Observe that  $(k^\times)^n = \mathbb{A}^n \setminus \mathbf{V}(X_1 \cdots X_n)$ . Then, the coordinate ring of  $(k^\times)^n$  is

$$k[X_1, \dots, X_n]_{X_1 \cdots X_n} = k[X_1, X_1^{-1}, \dots, X_n, X_n^{-1}],$$

the ring of Laurent polynomials.

---

<sup>2</sup>We emphasise that  $V$  is irreducible. This assumption is not necessary but it simplifies the discussion.

**2.10. Regular functions.** Let  $V$  be an algebraic subset of  $\mathbb{A}^n$ . When studying some class of objects, we need an appropriate idea of the structure-preserving maps between them. One such kind of a map for algebraic sets are the polynomial maps, which we saw in §2.8. A function  $V \rightarrow k$  being polynomial is a global property, as the function must be defined by a single polynomial for all points of  $V$ . This global property is often too restrictive, so we must develop a local structure-preserving property. For example, in real analysis, differentiability is a local property: a function is differentiable if it is differentiable at each point, and this can be checked on an arbitrary neighbourhood of a point. In this section, we define the notion of a regular function on  $V$ , which is a local property. This gives us a less restrictive concept of a structure-preserving map for algebraic sets.

**Definition 20.** Let  $U$  be an open subset of  $V$ . A function  $f : U \rightarrow k$  is called *regular at*  $P \in U$  if there exist  $g, h \in k[V]$  with  $h(P) \neq 0$  such that  $f = \frac{g}{h}$  on some neighbourhood of  $P$ . A function  $f : U \rightarrow k$  is called *regular* if it is regular at every  $P \in U$ .

**Example 21.** Consider  $V = \mathbf{V}(X_1X_4 - X_2X_3) \subseteq \mathbb{A}^4$  and the open subset

$$U = V \setminus \mathbf{V}(X_2X_4) = \{(x_1, \dots, x_4) \in V : x_2 \neq 0 \text{ or } x_4 \neq 0\}.$$

Define the function

$$\varphi : U \rightarrow k, \quad (x_1, \dots, x_4) \mapsto \begin{cases} \frac{x_1}{x_2} & \text{if } x_2 \neq 0, \\ \frac{x_3}{x_4} & \text{if } x_4 \neq 0. \end{cases}$$

It is well-defined since  $\frac{x_1}{x_2} = \frac{x_3}{x_4}$  if  $x_2 \neq 0$  and  $x_4 \neq 0$ , and regular since it is locally given by quotients of polynomials.

We write  $\mathcal{O}_V(U)$  for the ring of regular functions  $U \rightarrow k$ . The assignment of an open set  $U \mapsto \mathcal{O}_V(U)$  satisfies the following properties:

- Proposition 22.** (1)  $\mathcal{O}_V(U)$  is a  $k$ -subalgebra of all  $k$ -valued functions on  $U$ , i.e.,  $\mathcal{O}_V(U)$  contains all the constant functions and is closed under addition and multiplication.  
(2) If  $f \in \mathcal{O}_V(U)$  and  $U'$  is an open subset of  $U$ , then  $f|_{U'} \in \mathcal{O}_V(U')$ .  
(3) If  $\{U_i\}$  is an open cover of  $U$  and  $f_i \in \mathcal{O}_V(U_i)$  satisfy  $f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$  for all  $i$  and  $j$ , then there is a unique  $f \in \mathcal{O}_V(U)$  such that  $f|_{U_i} = f_i$  for all  $i$ .

*Proof.* (1) It is clear that a constant function is regular. Let  $f_1, f_2$  be regular on  $U$  and fix a point  $P \in U$ . Then, there exists a neighbourhood  $W$  of  $P$  and  $g_1, g_2, h_1, h_2 \in k[V]$  such that  $f_i = \frac{g_i}{h_i}$  on  $W$ . Therefore,  $f_1 + f_2 = \frac{g_1h_2 + g_2h_1}{h_1h_2}$  and  $f_1f_2 = \frac{g_1g_2}{h_1h_2}$  on  $W$ .  
(2) For any  $P \in U'$ ,  $f$  is regular at  $P \in U$ , so  $f|_{U'}$  is regular.  
(3) The function  $f$  is uniquely determined by the requirement  $f = f_i$  on  $U_i$ . To see  $f \in \mathcal{O}_V(U)$ , note that if  $P \in U$ , then  $P \in U_i$  for some  $i$ , and  $f$  is regular at  $P$  since it is locally equal to  $f_i$ . □

### 3. AFFINE VARIETIES

**3.1. Sheaves and their morphisms.** Proposition 22 alludes to an important object appearing in many areas of mathematics, called a sheaf. We will use the notion of a sheaf of  $k$ -algebras to define affine varieties, opting to avoid the most general definition of sheaves appearing in the literature (c.f. [Har77, Chapter II, §1]).

**Definition 23** ([Mil13, Chapter 3, a.]). *Let  $V$  be a topological space and  $k$  a field. Suppose that for every open subset  $U \subseteq V$ , we have a set  $\mathcal{O}_V(U)$  of functions  $U \rightarrow k$ . We say the assignment  $U \mapsto \mathcal{O}_V(U)$  is a sheaf of  $k$ -algebras if the following hold for every open subset  $U \subseteq V$ :*

- (1)  $\mathcal{O}_V(U)$  is a  $k$ -subalgebra of all  $k$ -valued functions on  $U$ , i.e.,  $\mathcal{O}_V(U)$  contains the constant functions and is closed under addition and multiplication;
- (2) if  $f \in \mathcal{O}_V(U)$  and  $U' \subseteq U$  is an open subset, then  $f|_{U'} \in \mathcal{O}_V(U')$ ; and,
- (3) if  $\{U_i\}$  is an open cover of  $U$  and  $f_i \in \mathcal{O}_V(U_i)$  satisfy  $f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$  for all  $i$  and  $j$ , then there exists a unique  $f \in \mathcal{O}_V(U)$  such that  $f|_{U_i} = f_i$  for each  $i$ .

A pair  $(V, \mathcal{O}_V)$  consisting of a topological space  $V$  and a sheaf of  $k$ -algebras on  $V$  is called a  $k$ -ringed space, or a ringed space when  $k$  is understood. The  $k$ -algebra  $\mathcal{O}_V(U)$  is sometimes denoted  $\Gamma(U, \mathcal{O}_V)$ , and its elements are called the sections of  $\mathcal{O}_V$  over  $U$ .

Proposition 22 says that for an algebraic set  $V$ , the assignment of an open subset  $U \subseteq V$  to its ring of regular functions  $\mathcal{O}_V(U)$  is a sheaf of  $k$ -algebras. This sheaf is called the structure sheaf of  $V$ . As we will see, structure sheaves are one of the main tools we use to define and study varieties. To further illustrate the definition of a sheaf, we give another example as well as a counter-example:

**Example 24.** (1) *Let  $V$  be any topological space. For any open set  $U$ , let  $\mathcal{O}_V(U)$  be the set of continuous functions  $U \rightarrow \mathbb{R}$ . It is clear  $\mathcal{O}_V$  satisfies condition (1) in Definition 23, and conditions (2) and (3) hold since continuity is a local property. Thus  $\mathcal{O}_V$  is a sheaf of  $\mathbb{R}$ -algebras.*

- (2) *Consider  $V = \mathbb{R}$  with the standard topology, and let  $\mathcal{O}_V(U)$  be the set of bounded functions  $U \rightarrow \mathbb{R}$ . This does not define a sheaf since condition (3) of Definition 23 does not hold; let  $\{U_i\}$  be the open cover of  $\mathbb{R}$  given by  $U_i := (-i, i)$  and observe that  $f_i(x) := x$  lies in  $\mathcal{O}_V(U_i)$  for each  $i$ ; however,  $f(x) = x$  does not lie in  $\mathcal{O}_V(\mathbb{R})$ . This fails to be a sheaf because boundedness is not determined locally.*

So far, we have studied algebraic subsets embedded in an ambient affine space. Roughly speaking, an *affine variety* is an algebraic subset of  $\mathbb{A}^n$ , defined without choosing a specific embedding of our algebraic set into affine space; this is analogous to how a smooth manifold is defined intrinsically as a topological space, without needing to be embedded in some Euclidean space. The following definition uses ringed spaces to make this precise:

**Definition 25.** *An affine variety over  $k$  is a  $k$ -ringed space isomorphic to one of the form  $(V, \mathcal{O}_V)$  for some algebraic set  $V \subseteq \mathbb{A}^n$ .*

We have seen that any algebraic set  $V$  gives a reduced finitely generated  $k$ -algebra,  $k[V]$ . Conversely, we can ask, given a reduced finitely generated  $k$ -algebra  $A$ , whether there is an associated affine variety. Choose generators  $a_1, \dots, a_n$  for  $A$ , so that  $A = k[a_1, \dots, a_n]$ . The  $k$ -algebra homomorphism  $\varphi : k[X_1, \dots, X_n] \rightarrow A$  given by  $X_i \mapsto a_i$  induces an isomorphism

$k[X_1, \dots, X_n]/\ker \phi \cong A$ . Then  $\ker \phi$  is radical, so that  $\mathbf{V}(\ker \phi)$  is an algebraic subset of  $\mathbb{A}^n$  with coordinate ring  $A$ . In view of Corollary 17, isomorphic  $k$ -algebras correspond to isomorphic algebraic sets. To construct an affine variety that

The answer is yes—this variety is the *maximal spectrum*,  $\text{Spec}(A)$ , of  $A$ . The maximal spectrum  $\text{Spec}(A)$  is a topological space with a certain sheaf that defines an affine variety. This construction is technical so we omit it in this document, but the details can be found in [Mil13, Chapter 3].

For the remainder of the document, we will use  $\text{Spec}(A)$  to denote the affine variety which has  $A$  as its coordinate ring. The algebra  $A$  will usually be defined in terms of generators and relations so that  $A \cong k[X_1, \dots, X_n]/I$  for some radical ideal  $I$ ; then we can think of  $\text{Spec}(A)$  as  $\mathbf{V}(I) \subseteq \mathbb{A}^n$

### 3.2. Affine varieties.

**Definition 26.** A morphism of  $k$ -ringed spaces  $(V, \mathcal{O}_V) \rightarrow (W, \mathcal{O}_W)$  is a continuous map  $\varphi : V \rightarrow W$  such that for all open subsets  $U \subseteq W$ , if  $f \in \mathcal{O}_W(U)$ , then  $f \circ \varphi \in \mathcal{O}_V(\varphi^{-1}(U))$ .

For every pair of open subsets  $U \subseteq V$  and  $U' \subseteq W$  satisfying  $\varphi(U) \subseteq U'$ , we have a homomorphism of  $k$ -algebras,  $\varphi^* : \mathcal{O}_W(U') \rightarrow \mathcal{O}_V(U)$ , given by  $f \mapsto \varphi^* f := f \circ \varphi$ .

**3.3. Tangent spaces.** In differential geometry, a key tool used to study a surface in  $\mathbb{R}^3$  is the tangent space. In algebraic geometry, there is also a notion of the tangent space, but this is defined without the use of calculus. The dimensions of the tangent spaces of an affine variety give one way to define the dimension of the variety. With dimension defined, we can also consider the non-singular and singular points on a variety; the singular points are the points where the tangent space does not provide a well-defined linear approximation to the variety.


We first define the tangent space for a hypersurface in  $\mathbb{A}^n$ , i.e., an algebraic set of the form  $V = \mathbf{V}(f) \subseteq \mathbb{A}^n$  for some non-constant irreducible  $f \in k[X_1, \dots, X_n]$ . Let  $P$  be a point  $V$ . In what follows, let  $\frac{\partial f}{\partial X_i}$  denote the formal partial derivative of the polynomial  $f$ . For example,  $\frac{\partial}{\partial X_i} X_i^n = nX_i^{n-1}$ , and the partial derivative of an arbitrary polynomial can be computed using the Leibniz rule and linearity; calculus is not needed to define this formal partial derivative, so the definition makes sense for any field  $k$ . Define the first-order part of  $f$  at  $P$  by

$$(2) \quad f_P^{(1)} := \sum_{i=1}^n \frac{\partial f}{\partial X_i}(P)(X_i - a_i).$$

Then the tangent space of  $V$  at  $P$  is

$$T_P V := \mathbf{V}(f_P^{(1)}).$$

This is an affine subspace of  $\mathbb{A}^n$  (i.e., a translation of a vector subspace) which passes through  $P$  and is tangent to  $f$ , matching our usual intuition for what a tangent space is. The following figure gives an example to illustrate:



$\mathbf{V}(f)$  and  $\mathbf{V}(f_P^{(1)})$  for  $f = y^2 - x^3 + x - 1$  and  $P = (-1, 1)$ .

Now let  $V \subseteq \mathbb{A}^n$  be a general algebraic set. For any  $f \in \mathbf{I}(V)$ , the first-order part  $f_P^{(1)}$  is defined as in equation 2. The tangent space of  $V$  at  $P$  is

$$T_P V := \bigcap_{f \in \mathbf{I}(V)} \mathbf{V}(f_P^{(1)}).$$

Then  $T_P V$  is the intersection of all affine subspaces tangent at  $P$  to some  $f$  in the ideal  $\mathbf{I}(V)$  defining  $V$ . Suppose  $f_1, \dots, f_m$  generate  $\mathbf{I}(V)$ . Then for any  $g = \sum_{i=1}^m h_i f_i \in \mathbf{I}(V)$  and  $P \in V$ ,

$$\begin{aligned} g_P^{(1)} &= \sum_{i=1}^n \frac{\partial g}{\partial X_i}(P)(X_i - a_i) \\ &= \sum_{i=1}^n \sum_{j=1}^m \left( \frac{\partial h_j}{\partial X_i}(P) f_j(P) + h_j(P) \frac{\partial f_j}{\partial X_i}(P) \right) (X_i - a_i) \\ &= \sum_{j=1}^m h_j(P) \sum_{i=1}^n \frac{\partial f_j}{\partial X_i}(P)(X_i - a_i), \end{aligned}$$

so  $g_P^{(1)}$  is a linear combination of the polynomials  $f_{j,P}^{(1)}$ . This implies that

$$\bigcap_{i=1}^m \mathbf{V}(f_i^{(1)}) \subseteq \mathbf{V}(g_P^{(1)}),$$

and the definition of  $T_P V$  simplifies to

$$T_P V = \bigcap_{i=1}^m \mathbf{V}(f_{i,P}^{(1)}).$$

Let us consider how  $T_P V$  is a vector space over  $k$ . The zero vector is the point  $P$ . If  $Q_1, Q_2 \in \mathbb{A}^n$  lying in  $T_P V$  can be written  $Q_i = \tilde{Q}_i + P$ , then

$$Q_1 + Q_2 = (\tilde{Q}_1 + \tilde{Q}_2) + P, \quad \lambda \cdot Q_i = \lambda \tilde{Q}_i + P, \quad \lambda \in k.$$



This is the usual vector space structure on  $k^n$  if the origin is translated to  $P$ .

We can now define the dimension of the algebraic set  $V$  by

$$\dim V := \min\{\dim T_P V : P \in V\}.$$

A point  $P \in V$  is called non-singular if  $\dim T_P V = \dim V$  and singular if  $\dim T_P V < \dim V$ . We denote the set of non-singular and singular points by  $V_{\text{non-sing}}$  and  $V_{\text{sing}}$ , respectively; we say  $V$  is non-singular if it is non-singular at each  $P \in V$ . For an integer  $r$  such that  $0 \leq r \leq n$ , define the subset

$$S(r) := \{P \in V : \dim T_P V \geq r\} \subseteq V.$$

If  $d := \dim V$ , we have that  $V = S(d)$  and  $V_{\text{sing}} = S(d+1)$ . We claim that the set of non-singular points  $V_{\text{non-sing}}$  is an open subset of  $V$ , because

$$V_{\text{non-sing}} = V \setminus V_{\text{sing}} = V \setminus S(d+1),$$

and  $S(d+1)$  is closed. That  $S(d+1)$  is closed follows from the following proposition:

**Proposition 27** ([Rei88, §6.5]). *For any integer  $r$  such that  $0 \leq r \leq n$ , the subset  $S(r)$  is closed.*

*Proof.* Suppose Thus,  $T_P V$  is the set of  $(x_1, \dots, x_n) \in k^n$  solving

$$\sum_{i=1}^n \frac{\partial f_j}{\partial X_i}(x_i - a_i) = 0, \quad \text{for all } j = 1, \dots, m.$$

Then  $T_P V$  can be identified with the kernel of the matrix  $J := \left( \frac{\partial f_i}{\partial X_j}(P) \right)_{i=1, \dots, m, j=1, \dots, n}$ .

Checking  $P \in S(r)$  is equivalent to checking  $\text{rank } J \leq n - r$ . In turn, this is equivalent to ensuring every  $(n - r + 1) \times (n - r + 1)$  minor of  $J$  vanishes. The entries of  $J$  are polynomials in  $P = (a_1, \dots, a_n)$ . Then each minor of  $J$  is also a polynomial and so  $S(r) \subseteq V$  is an algebraic set.  $\square$

**Example 28.** *Let us give some examples of how to compute tangent spaces and dimensions of algebraic sets.*

- (1)  $\mathbb{A}^n$
- (2)  $\mathbf{V}(f)$  for  $f$  non-constant and irreducible
- (3) Singular and non-singular points of  $y^2 = x^3 - 6x^2 - 9x$

Although the description we have given for  $T_P V$  in terms of first-order parts of polynomials is geometrically intuitive, we now prove a theorem which gives an intrinsic description of  $T_P V$  in terms of ideals in the coordinate ring of  $V$ .

For the setup, let  $V$  be an algebraic subset of  $\mathbb{A}^n$  and suppose  $P \in V$ . Changing coordinates if necessary, we may assume without loss of generality that  $P = (0, \dots, 0)$ . Now  $T_P V$  is a vector subspace of  $k^n$ . Write  $\mathfrak{m}_P \subseteq k[V]$  for the maximal ideal of regular functions vanishing at  $P$ . Denote by  $M_P$  the ideal  $(X_1, \dots, X_n) \subseteq k[X_1, \dots, X_n]$ . We have  $\mathfrak{m}_P = M_P / \mathbf{I}(V)$ .

**Theorem 29.** *There is a natural isomorphism of vector spaces*

$$(T_P V)^* \cong \mathfrak{m}_P / \mathfrak{m}_P^2.$$

*Proof.* We first prove the special case  $V = \mathbb{A}^n$ , where we must show  $M_P/M_P^2 \cong (k^n)^*$ . Note that  $X_1, \dots, X_n$  form a basis for  $(k^n)^*$ . As  $P = (0, \dots, 0)$ ,

$$f_P^{(1)} = \sum_{i=1}^n \frac{\partial f}{\partial X_i}(P) X_i$$

is a linear form on  $k^n$ , and we can define a map  $d : M_P \rightarrow (k^n)^*$  by  $f \mapsto f_P^{(1)}$ . Then  $d$  is surjective since the  $X_i \in M_P$  map to a basis for  $(k^n)^*$ . A general  $f \in M_P$  can be written

$$f = \sum_{i=1}^n c_i X_i + \text{higher order terms},$$

for some  $c_i \in k$ . Since  $P = (0, \dots, 0)$ ,

$$f_P^{(1)} = \sum_{i=1}^n \frac{\partial f}{\partial X_i}(P) X_i = \sum_{i=1}^n c_i X_i.$$

Therefore  $f \in \ker d$ , i.e.,  $f_P^{(1)} = 0$ , if and only if each  $c_i = 0$ . This is equivalent to  $f$  lying in  $M_P^2$ , as  $M_P^2$  is generated by the monomials  $X_i X_j$ . Thus  $d$  induces the isomorphism  $M_P/M_P^2 \cong (k^n)^*$ .

For the general case, first observe that we have a surjective map  $(k^n)^* \rightarrow (T_P V)^*$  which takes a linear form to its restriction to  $T_P V$ . Composing  $d$  with this restriction map gives the map

$$D : M_P \rightarrow (k^n)^* \rightarrow (T_P V)^*,$$

which is surjective since each factor is. To complete the proof, we show the following pair of isomorphisms:

$$(T_P V)^* \cong M_P/(M_P^2 + \mathbf{I}(V)) \cong \mathfrak{m}_P/\mathfrak{m}_P^2.$$

The first isomorphism follows from the fact that  $\ker D = M_P^2 + \mathbf{I}(V)$ , which we prove by giving a chain of equivalent conditions for  $f$  to lie in  $\ker D$ . By definition of  $D$ ,  $f \in \ker D$  means  $f_P^{(1)}|_{T_P V} = 0$ . If  $g_i \in \mathbf{I}(V)$  denote generators for  $\mathbf{I}(V)$ , we know  $T_P V = \bigcap_i \mathbf{V}(g_{i,P}^{(1)})$ . Then  $f_P^{(1)}|_{T_P V} = 0$  if and only if

$$f_P^{(1)} = \sum_i a_i g_{i,P}^{(1)}$$

for some  $a_i \in k$ . The above equality is equivalent to  $f$  and  $\sum_i a_i g_i$  only differing by quadratic terms, i.e.,

$$f - \sum_i a_i g_i \in M_P^2,$$

which is the same as  $f \in M_P^2 + \mathbf{I}(V)$ . To see why  $M_P/(M_P^2 + \mathbf{I}(V)) \cong \mathfrak{m}_P/\mathfrak{m}_P^2$ , define the map  $\varphi : M_P \rightarrow \mathfrak{m}_P/\mathfrak{m}_P^2$  by  $h \mapsto (h + \mathbf{I}(V)) + \mathfrak{m}_P^2$ . This is well-defined since  $\mathfrak{m}_P = M_P/\mathbf{I}(V)$ . Then

$$\ker \varphi = \{h \in M_P : h + \mathbf{I}(V) \in \mathfrak{m}_P^2\} = \{h \in M_P : h + \mathbf{I}(V) \in M_P^2/\mathbf{I}(V)\} = M_P^2 + \mathbf{I}(V),$$

so  $\varphi$  induces the desired isomorphism.  $\square$

## 4. CONVEX GEOMETRY

Affine toric varieties are a class of algebraic varieties which are determined by a *cone* in a vector space. The interplay between the variety and its cone leads to a rich theory combining the algebraic geometry of the variety and the convex geometry of the cone. Moreover, computations with the cone can often be done explicitly, which means toric varieties are useful to study as examples.

The most succinct definition of a toric variety  $X$  is the following:  $X$  is a normal variety with an algebraic torus  $T$  as a dense open subset, and  $T$  acts on  $X$  by an action which extends the natural action of  $T$  on itself. However, this description doesn't indicate the relationship with convex geometry. In this chapter, we will define and study toric varieties; we will also see how the GIT quotient of an affine space by a torus has the structure of an affine toric variety. We start by discussing the prerequisite convex geometry.

**4.1. Convex cones.** Let  $N$  be a lattice, i.e., a free abelian group of finite rank  $n$ . Let  $M = \text{Hom}(N, \mathbb{Z})$  denote the dual lattice, with dual pairing  $\langle \cdot, \cdot \rangle : M \times N \rightarrow \mathbb{Z}$ . We can consider  $N$  and  $M$  as subsets of the  $n$ -dimensional vector spaces  $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$  and  $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$ , respectively. Note that  $N_{\mathbb{R}}$  and  $M_{\mathbb{R}}$  are dual vector spaces, and we retain the notation  $\langle \cdot, \cdot \rangle$  for the dual pairing  $M_{\mathbb{R}} \times N_{\mathbb{R}} \rightarrow \mathbb{R}$ .

A subset  $\sigma \subseteq N_{\mathbb{R}}$  is a *cone* if it is closed under nonnegative scalar multiplication, i.e.,  $\lambda x \in \sigma$  for all  $x \in \sigma$  and all  $\lambda \in \mathbb{R}_{\geq 0}$ . A set  $\sigma \subseteq N_{\mathbb{R}}$  is *convex* if for any two points in  $\sigma$ , the line segment joining them is contained in  $\sigma$ , i.e.,  $x, y \in \sigma$  implies  $\lambda x + (1 - \lambda)y \in \sigma$  for all  $\lambda \in [0, 1]$ . Since cones are closed under positive scalar multiplication by assumption, a cone  $C$  is convex if and only if it is closed under addition.

**Example 30.** *The union of rays*

$$\sigma_1 = \{(x, x/\sqrt{3}) \in \mathbb{R}^2 : x \in \mathbb{R}_{\geq 0}\} \cup \{(x, \sqrt{3}x) \in \mathbb{R}^2 : x \in \mathbb{R}_{\geq 0}\}$$




is cone but not convex. We can “fill in”  $\sigma_1$  to get the convex cone

$$\sigma_2 = \{(x, y) \in \mathbb{R}^2 : x \in \mathbb{R}_{\geq 0}, x/\sqrt{3} \leq y \leq \sqrt{3}x\}.$$

The set

$$\sigma_3 = \{(x, r) \in \mathbb{R}^2 \times \mathbb{R} : \|x\| \leq r\}$$

is a convex cone in  $\mathbb{R}^3$ .



**Definition 31.** Let  $\sigma$  be a cone. The dual cone  $\sigma^\vee$  is

$$\sigma^\vee = \{u \in M_{\mathbb{R}} : \langle u, v \rangle \geq 0 \text{ for all } v \in \sigma\}.$$

**Example 32.** In the following examples, we let  $N = \mathbb{Z}^n \subseteq \mathbb{R}^n \cong N_{\mathbb{R}}$ . Let  $e_1, \dots, e_n$  be the standard basis for  $N$  and  $e_1^*, \dots, e_n^*$  the dual basis for  $M$ .

- (1) Let  $\sigma := \text{span}_{\mathbb{R}_{\geq 0}}\{e_1, \dots, e_n\}$ . Observe that a functional  $\sum_{i=1}^n a_i e_i^*$  is in the dual cone if and only if  $a_i \geq 0$  for all  $i$ . Then  $\sigma^\vee = \text{span}_{\mathbb{R}_{\geq 0}}\{e_1^*, \dots, e_n^*\}$ .
- (2) Let  $n = 2$  and let  $\sigma = \text{span}_{\mathbb{R}_{\geq 0}}\{e_1, -e_1 + 2e_2\}$ . If  $u \in M_{\mathbb{R}}$ , checking  $\langle u, v \rangle \geq 0$  for all  $v \in \sigma$  is equivalent to checking for the generators  $v = e_1$  and  $v = -e_1 + 2e_2$ . So then  $u = ae_1^* + be_2^* \in M_{\mathbb{R}}$  is in  $\sigma^\vee$  exactly when the following two inequalities hold:

$$\langle u, e_1 \rangle = a \geq 0, \quad \langle u, -e_1 + 2e_2 \rangle = -a + 2b \geq 0.$$

Below we have a sketch of  $\sigma$  and  $\sigma^\vee$ :



Then we can see that  $\sigma^\vee = \text{span}_{\mathbb{R}_{\geq 0}}\{2e_1 + e_2, e_2\}$ .

The following theorem relates a cone to its dual and has many important consequences (which we will see in §4.2):

**Theorem 33** ([Ful93, §1.2]). Let  $\sigma$  be a closed convex cone in  $N_{\mathbb{R}}$ . If  $v_0 \notin \sigma$ , then there exists  $u_0 \in \sigma^\vee$  such that  $\langle u_0, v_0 \rangle < 0$ .

Theorem 33 is a foundational result for the theory of toric varieties. References such as [Ful93], [CLS11], and [Oda88] omit a proof, but we present one for completeness. We begin with a lemma from analysis.

**Lemma 34.** Let  $A$  and  $B$  be disjoint closed subsets of a finite-dimensional normed vector space  $(V, \|\cdot\|)$ , and assume  $A$  is compact. Then there exist  $a_0 \in A$  and  $b_0 \in B$  which minimise the distance  $\|a - b\|$  over all  $a \in A$  and  $b \in B$ .

*Proof.* Take arbitrary  $a \in A$  and  $b \in B$  and set  $r_1 = \|a - b\| > 0$ . Since  $A$  is compact, there exists  $r_2 > 0$  such that  $A \subseteq \overline{B_{r_2}(a)}$ . Let  $S = B \cap \overline{B_{r_1+r_2}(a)}$ , which is nonempty since  $b \in S$ . Since the distance function is continuous and  $A \times S$  is compact, there exist  $(a_0, b_0) \in A \times S$  minimising the distance  $\|a_0 - b_0\|$  for all pairs of points in  $A \times S$ . We claim that this is in fact the minimum for all pairs of points in  $A \times B$ ; suppose to the contrary that there is  $(a', b') \in A \times B$  with  $\|a' - b'\| < \|a_0 - b_0\|$ . In particular, since  $\|a_0 - b_0\| \leq r_1$ ,  $\|a' - b'\| < r_1$  and so  $\|a - b'\| \leq \|a - a'\| + \|a' - b'\| < r_2 + r_1$ . This implies  $b' \in S$ , contradicting that  $\|a_0 - b_0\|$  attained the minimum distance for pairs in  $A \times S$ .  $\square$

**Theorem 35** (Hyperplane Separation Theorem [BV04, §2.5.1]). Let  $A$  and  $B$  be nonempty disjoint closed convex sets in a Euclidean vector space  $(V, (\cdot, \cdot))$ , and assume  $A$  is compact. Then there exists  $v \in V \setminus \{0\}$  and  $\lambda \in \mathbb{R}$  such that for all  $a \in A$ ,  $(v, a) \leq \lambda$ , and for all

$b \in B$ ,  $(v, b) \geq \lambda$ . In other words, there is an affine hyperplane such that  $A$  is contained in the negative halfspace and  $B$  is contained in the positive halfspace.

*Proof.* Lemma 34 yields  $a_0 \in A$  and  $b_0 \in B$  minimising the distance between points in  $A$  and points in  $B$ . Define

$$v := b_0 - a_0, \quad \lambda := \frac{\|b_0\|^2 - \|a_0\|^2}{2} = \frac{1}{2}(b_0 - a_0, b_0 + a_0).$$

The set of  $x \in V$  satisfying  $(v, x) = \lambda$  is the hyperplane orthogonal to the line segment joining  $a_0$  and  $b_0$ , and passing through its midpoint. Let us prove that  $(v, b) \geq \lambda$  for all  $b \in B$  (a similar argument shows  $(v, a) \leq \lambda$  for all  $a \in A$ ). Assume for a contradiction that there exists  $u \in B$  with  $(v, u) < \lambda$ . Then,

$$(v, u) - \frac{1}{2}(v, b_0 + a_0) < 0 \implies (v, u - b_0 + \frac{1}{2}(b_0 - a_0)) < 0 \implies (v, u - b_0) + \frac{1}{2}\|v\|^2 < 0.$$

Thus  $(v, u - b_0) < 0$ . Using the Leibniz rule for differentiating inner products, we see

$$\left. \frac{d}{dt} \|v + t(u - b_0)\|^2 \right|_{t=0} = 2(v + t(u - b_0), u - b_0)|_{t=0} = 2(v, u - b_0) < 0,$$

so for small  $t > 0$ , we have

$$\|b_0 + t(u - b_0) - a_0\| = \|v + t(u - b_0)\| < \|v\| = \|b_0 - a_0\|.$$

But  $B$  is convex and  $b_0, u \in B$ , so  $(b_0 + t(u - b_0)) \in B$  when  $t$  is sufficiently small. This contradicts the minimality of  $\|b_0 - a_0\|$ .  $\square$

*Proof of Theorem 33* ([BV04, Example 2.20]). Fix a basis  $e_1, \dots, e_n$  for  $N_{\mathbb{R}}$  and endow  $N_{\mathbb{R}}$  with the Euclidean inner product which makes the basis orthonormal (i.e.,  $(e_i, e_j) = \delta_{ij}$ ). Since  $\sigma$  is closed and  $v_0 \notin \sigma$ , there exists  $\varepsilon > 0$  such that  $\overline{B_\varepsilon(v_0)}$  does not intersect  $\sigma$ . By Theorem 35, there exist  $v \in N_{\mathbb{R}}$  and  $\lambda \in \mathbb{R}$  such that

$$(v, x) \geq \lambda \quad \text{and} \quad (v, y) \leq \lambda$$

for all  $x \in \sigma$  and all  $y \in \overline{B_\varepsilon(v_0)}$ . Since  $x = 0 \in \sigma$ ,  $\lambda \leq 0$ . In fact, we claim that  $(v, x) \geq 0$  for all  $x \in \sigma$  and  $(v, v_0) < 0$ , which completes the proof as then  $u_0 := (x \mapsto (v, x))$  is a linear form in  $\sigma^\vee$  with  $\langle u_0, v_0 \rangle < 0$ . Suppose there exists  $x \in \sigma$  with  $(v, x) = \lambda_0 < 0$ . Then for any  $s \in \mathbb{R}_{\geq 0}$ ,  $sx \in \sigma$  and  $(v, sx) = s\lambda_0$ , contradicting that  $\{(v, x) : x \in \sigma\}$  is bounded from below. Assume for a contradiction  $(v, v_0) = 0$ . We have  $y = v_0 + \frac{\varepsilon v}{\|v\|} \in \overline{B_\varepsilon(v_0)}$  and

$$0 \geq \lambda \geq (v, y) = (v, v_0 + \frac{\varepsilon v}{\|v\|}) = \varepsilon \|v\| > 0,$$

a contradiction.  $\square$

**Corollary 36.** *Let  $\sigma$  be a closed cone in  $N_{\mathbb{R}}$ . Then  $(\sigma^\vee)^\vee = \sigma$ .*

*Proof.* If  $v \in \sigma$ ,  $\langle u, v \rangle \geq 0$  for all  $u \in \sigma^\vee$ , so  $v \in (\sigma^\vee)^\vee$ . If  $v \notin \sigma$ , Theorem 33 implies there exists  $u \in \sigma^\vee$  with  $\langle u, v \rangle < 0$  and hence  $v \notin (\sigma^\vee)^\vee$ .  $\square$

**4.2. Polyhedral cones.** In the study of toric varieties, we are interested in the following class of closed convex cones:

**Definition 37.** A subset  $\sigma \subseteq N_{\mathbb{R}}$  is called a *convex polyhedral cone* if

$$\sigma = \text{span}_{\mathbb{R}_{\geq 0}}\{v_1, \dots, v_s\}$$

for a (finite) set of generators  $v_1, \dots, v_s \in N_{\mathbb{R}}$ .

It follows from the definition that convex polyhedral cones are indeed convex cones. Every example of a cone we have seen so far has been polyhedral, except for  $\sigma_3 = \{(x, r) \in \mathbb{R}^2 \times \mathbb{R} : \|x\| \leq r\}$  in Example 30. As given, a more apt name for this definition would be “finitely-generated cone”; the name polyhedral is justified by the fact that a cone satisfies Definition 37 if and only if it is a finite intersection of closed halfspaces with 0 on their boundary (i.e., is polyhedral)—see [DLHK13, §1.3] for a proof.

Fulton [Ful93, §1.2] states and proves many important consequences of Theorem 33 for convex polyhedral cones. We explain a few of these now.

Any  $u \in M_{\mathbb{R}}$  defines a hyperplane  $u^{\perp}$  and corresponding nonnegative halfspace  $u^{\vee}$ . These are defined as

$$u^{\perp} := \{v \in N_{\mathbb{R}} : \langle u, v \rangle = 0\}, \quad u^{\vee} := \{v \in N_{\mathbb{R}} : \langle u, v \rangle \geq 0\}.$$

A face  $\tau$  of  $\sigma$  is the intersection of  $\sigma$  with a hyperplane which is nonnegative on  $\sigma$ :

$$\tau = \sigma \cap u^{\perp} = \{v \in \sigma : \langle u, v \rangle = 0\}, \text{ for some } u \in \sigma^{\vee}.$$

A facet is a codimension one face. Any proper face is the intersection of all facets containing it. When  $\sigma$  spans  $N_{\mathbb{R}}$  and  $\tau$  is a facet of  $\sigma$ , there exists  $u \in \sigma^{\vee}$ , which is unique up to multiplication by a positive scalar, such that

$$\tau = \sigma \cap u^{\perp}.$$

We denote such a vector by  $u_{\tau}$ , which gives the equation for the hyperplane spanned by  $\tau$ . The face  $\sigma \cap u^{\perp}$  is generated by the vectors  $v_i$  in the generating set for  $\sigma$  such that  $\langle u, v_i \rangle = 0$ . If  $\sigma$  spans  $N_{\mathbb{R}}$ , the dual cone is generated by  $\{u_{\tau} : \tau \text{ is a facet}\}$ . This implies the dual of a convex polyhedral cone is also a convex polyhedral cone (this result is true without assuming  $\sigma$  spans  $N_{\mathbb{R}}$ ). Example 38 below gives an example of computing facets and dual cone generators.

**Example 38.** Let  $N = \mathbb{Z}^3$  and let  $\sigma = \text{span}_{\mathbb{R}_{\geq 0}}\{e_1, e_2, e_1 + e_3, e_2 + e_3\} \subseteq N_{\mathbb{R}} \cong \mathbb{R}^3$ . Observe that

$$u_{\tau_1} = e_1^*, \quad u_{\tau_2} = e_2^*, \quad u_{\tau_3} = e_3^*, \quad u_{\tau_4} = e_1^* + e_2^* - e_3^*$$

are all nonnegative on  $\sigma$  and hence lie in  $\sigma^{\vee}$ . For each  $u_{\tau_i}$ , there is the corresponding facet  $\tau_i = \sigma \cap u_{\tau_i}^{\perp}$ . These are:


$$\begin{aligned} \tau_1 &= \text{span}_{\mathbb{R}_{\geq 0}}\{e_2, e_2 + e_3\}, & \tau_2 &= \text{span}_{\mathbb{R}_{\geq 0}}\{e_1, e_1 + e_3\}, \\ \tau_3 &= \text{span}_{\mathbb{R}_{\geq 0}}\{e_1, e_2\}, & \tau_4 &= \text{span}_{\mathbb{R}_{\geq 0}}\{e_1 + e_3, e_2 + e_3\}. \end{aligned}$$

The other faces in  $\sigma$  are the rays

$$\begin{aligned} \tau_1 \cap \tau_3 &= \text{span}_{\mathbb{R}_{\geq 0}}\{e_2\}, & \tau_1 \cap \tau_4 &= \text{span}_{\mathbb{R}_{\geq 0}}\{e_2 + e_3\}, \\ \tau_2 \cap \tau_3 &= \text{span}_{\mathbb{R}_{\geq 0}}\{e_1\}, & \tau_2 \cap \tau_4 &= \text{span}_{\mathbb{R}_{\geq 0}}\{e_1 + e_3\}, \end{aligned}$$

and the origin  $\bigcap_{i=1}^4 \tau_i = \{0\}$ . The dual cone is generated by  $\{u_{\tau_i}\}_{i=1}^4$ , so

$$\sigma^\vee = \text{span}_{\mathbb{R}_{\geq 0}} \{e_1^*, e_2^*, e_3^*, e_1^* + e_2^* - e_3^*\}.$$



A diagram of  $\sigma$  [CLS11, Figure 2]

**4.3. The semigroup of a cone.** Recall that a semigroup is a set with an associative binary operation. For each cone  $\sigma$ , the points of the dual lattice which are contained in  $\sigma^\vee$  form a semigroup,

$$S_\sigma := \sigma^\vee \cap M.^3$$

A convex polyhedral cone  $\sigma \subseteq N_{\mathbb{R}}$  is called *rational* if it can be generated by elements in the lattice  $N$ . When  $\sigma$  is rational,  $\sigma^\vee$  is also rational [Ful93, §1.2]. Gordan's lemma tells us that the semigroup of a rational cone is finitely generated:

**Theorem 39** (Gordan's lemma [Ful93, §1.2]). *If  $\sigma$  is a rational convex polyhedral cone, then  $S_\sigma$  is a finitely generated semigroup.*

*Proof.* Let  $u_1, \dots, u_s \in \sigma^\vee \cap M$  generate  $\sigma^\vee$  as a cone. Define

$$K = \left\{ \sum t_i u_i : 0 \leq t_i \leq 1 \right\}.$$

Since  $K$  is compact and  $M$  is discrete, the intersection  $K \cap M$  is finite. We claim  $K \cap M$  generates  $S_\sigma$ . Suppose that  $u \in S_\sigma$ . Then  $u = \sum r_i u_i$  for some  $r_i \in \mathbb{R}_{\geq 0}$  since  $\{u_i\}$  generates  $\sigma^\vee$ . Write each  $r_i$  as  $m_i + t_i$  for  $m_i \in \mathbb{Z}_{\geq 0}$  and  $0 \leq t_i \leq 1$ , so  $u = \sum m_i u_i + \sum t_i u_i$ . Clearly  $\sum t_i u_i \in K$ . Also,  $\sum t_i u_i = u - \sum m_i u_i \in M$  since  $u$  and  $\sum m_i u_i$  lie in  $M$  and  $M$  is a group. Then  $\sum t_i u_i \in K \cap M$ , and since  $u_1, \dots, u_s \in K \cap M$ , we have that  $u = \sum m_i u_i + \sum t_i u_i \in \text{span}_{\mathbb{Z}_{\geq 0}} K \cap M$ .  $\square$

<sup>3</sup>We have  $0 \in S_\sigma$  for any convex polyhedral cone  $\sigma$ . Then  $S_\sigma$  is always a semigroup with identity, i.e., a monoid. In the toric varieties literature, it is conventional to call  $S_\sigma$  a semigroup, even though calling it a monoid would be more precise; we follow the convention and refer to  $S_\sigma$  as a semigroup.

## 5. AFFINE TORIC VARIETIES

Affine toric varieties are a class of algebraic varieties which are determined by a *cone* in a vector space. The interplay between the variety and its cone leads to a rich theory combining the algebraic geometry of the variety and the convex geometry of the cone. Moreover, computations with the cone can often be done explicitly, which means toric varieties are useful to study as examples.

The most succinct definition of a toric variety  $X$  is the following:  $X$  is a normal variety with an algebraic torus  $T$  as a dense open subset, and  $T$  acts on  $X$  by an action which extends the natural action of  $T$  on itself. However, this description doesn't indicate the relationship with convex geometry. In this chapter, we will define and study toric varieties; we will also see how the GIT quotient of an affine space by a torus has the structure of an affine toric variety. We start by discussing the prerequisite convex geometry.

**5.1. The semigroup algebra  $S_\sigma$ .** Given a cone  $\sigma$ , we have seen how to construct a semigroup  $S_\sigma = \sigma^\vee \cap M$ . There is a corresponding semigroup algebra  $\mathbb{C}[S_\sigma]$ , which is a finitely generated commutative  $\mathbb{C}$ -algebra.

**Definition 40.** *Let  $\sigma$  be a rational cone in  $N$ . We define the affine toric variety corresponding to  $\sigma$  to be*

$$U_\sigma := \text{Spec}(\mathbb{C}[S_\sigma]).$$

Let us explain the details of the construction. Given the semigroup  $S_\sigma = \sigma^\vee \cap M$ , the semigroup algebra  $\mathbb{C}[S_\sigma]$  is defined as having a basis of formal symbols

$$\{\chi^u : u \in S_\sigma\},$$

with multiplication determined by addition in  $S_\sigma$ ,

$$\chi^u \cdot \chi^{u'} = \chi^{u+u'}.$$

Note the multiplication is commutative since  $S_\sigma$  is. We have the unit  $\chi^0 \in \mathbb{C}[S_\sigma]$  corresponding to  $0 \in S_\sigma$ . Since  $S_\sigma$  is finitely generated,  $\mathbb{C}[S_\sigma]$  is a finitely generated  $\mathbb{C}$ -algebra.

Consider the semigroup algebra corresponding to the full dual lattice,  $\mathbb{C}[M]$ . If  $e_1, \dots, e_n$  is a basis for  $N$  and  $e_1^*, \dots, e_n^*$  is the dual basis for  $M$ , denote

$$X_i := \chi^{e_i^*} \in \mathbb{C}[M].$$

As a semigroup,  $M$  is generated by  $\pm e_1^*, \dots, \pm e_n^*$ , so

$$\mathbb{C}[M] = \mathbb{C}[X_1^\pm, \dots, X_n^\pm],$$

and  $\mathbb{C}[M]$  can be identified with the ring of Laurent polynomials. For any cone  $\sigma$ ,  $S_\sigma$  is contained in  $M$ , so  $\mathbb{C}[S_\sigma] \subseteq \mathbb{C}[M]$ . Then any semigroup algebra  $\mathbb{C}[S_\sigma]$  can be thought of as a subalgebra of the Laurent polynomials.

As  $\mathbb{C}[S_\sigma]$  is a finitely generated and nilpotent free  $\mathbb{C}$ -algebra,  $U_\sigma = \text{Spec}(\mathbb{C}[S_\sigma])$  is a complex affine variety. For example, if  $\sigma = \{0\}$ , then  $S_{\{0\}} = M$ , and we have

$$U_{\{0\}} = \text{Spec}(\mathbb{C}[X_1^\pm, \dots, X_n^\pm]) = (\mathbb{C}^\times)^n.$$

**Example 41.** *Let us see more examples arising from the cones we have seen previously.*




- (1) If  $\sigma = \text{span}_{\mathbb{R}_{\geq 0}}\{e_1, \dots, e_n\}$ , then  $\sigma^\vee = \text{span}_{\mathbb{R}_{\geq 0}}\{e_1^*, \dots, e_n^*\}$ . The vectors  $e_1^*, \dots, e_n^*$  generate  $S_\sigma$  as a semigroup, so

$$\mathbb{C}[S_\sigma] = \mathbb{C}[\chi^{e_1^*}, \dots, \chi^{e_n^*}] = \mathbb{C}[X_1, \dots, X_n],$$

and

$$U_\sigma = \text{Spec}(\mathbb{C}[X_1, \dots, X_n]) = \mathbb{C}^n.$$

- (2) Recall that for  $\sigma = \text{span}_{\mathbb{R}_{\geq 0}}\{e_1, -e_1 + 2e_2\}$ , we have  $\sigma^\vee = \text{span}_{\mathbb{R}_{\geq 0}}\{2e_1^* + e_2^*, e_2^*\}$ .



Observe that while  $\{2e_1^* + e_2^*, e_2^*\}$  generates  $\sigma^\vee$  as a cone, it does not generate  $S_\sigma$  as a semigroup. For example,  $e_1^* + e_2^* \in S_\sigma$ , but  $e_1^* + e_2^* \notin \text{span}_{\mathbb{Z}_{\geq 0}}\{2e_1^* + e_2^*, e_2^*\}$ . However,  $\{2e_1^* + e_2^*, e_2^*, e_1^* + e_2^*\}$  does generate  $S_\sigma$ . Then

$$\mathbb{C}[S_\sigma] = \mathbb{C}[\chi^{e_2^*}, \chi^{2e_1^* + e_2^*}, \chi^{e_1^* + e_2^*}] = \mathbb{C}[X_2, X_1^2 X_2, X_1 X_2] \cong \mathbb{C}[X, Y, Z]/(XY - Z^2),$$

and

$$U_\sigma \cong \text{Spec}(\mathbb{C}[X, Y, Z]/(XY - Z^2)) = \mathbf{V}(XY - Z^2).$$

- (3) For  $\sigma = \text{span}_{\mathbb{R}_{\geq 0}}\{e_1, e_2, e_1 + e_3, e_2 + e_3\}$ , we saw  $\sigma^\vee = \text{span}_{\mathbb{R}_{\geq 0}}\{e_1^*, e_2^*, e_3^*, e_1^* + e_2^* - e_3^*\}$ . We have

$$\begin{aligned} \mathbb{C}[S_\sigma] &= \mathbb{C}[\chi^{e_1^*}, \chi^{e_2^*}, \chi^{e_3^*}, \chi^{e_1^* + e_2^* - e_3^*}] = \mathbb{C}[X_1, X_2, X_3, X_1 X_2 X_3^{-1}] \\ &\cong \mathbb{C}[X, Y, Z, W]/(XY - ZW), \end{aligned}$$

and so

$$U_\sigma \cong \text{Spec}(\mathbb{C}[X, Y, Z, W]/(XY - ZW)) = \mathbf{V}(XY - ZW).$$

In all the examples of cones that we have seen so far, the origin is a face of the cone. The following proposition gives equivalent conditions for the origin to be face:

**Proposition 42** ([Ful93, §1.2, Proposition 3]). *For a convex polyhedral cone  $\sigma$ , the following conditions are equivalent:*

- (1)  $\sigma \cap (-\sigma) = \{0\}$ ;
- (2)  $\sigma$  contains no nonzero linear subspace;
- (3) there is  $u \in \sigma^\vee$  with  $\sigma \cap u^\perp = \{0\}$ ;
- (4)  $\sigma^\vee$  spans  $M_{\mathbb{R}}$ .

A cone is called *strongly convex* if it satisfies the conditions of Proposition 42. We are usually interested in studying toric varieties arising from strongly convex cones. This assumption implies that when  $N$  is a rank  $n$  lattice, the dimension of the variety  $U_\sigma$  is also  $n$ . Specifically, when  $\sigma$  is strongly convex,  $U_\sigma$  has an  $n$ -dimensional torus as an open subset [CLS11, Theorem 1.2.18]. Then the affine toric variety  $U_\sigma$  defined in terms of the cone  $\sigma$  satisfies the definition of a toric variety from the introduction to this chapter, i.e.,  $U_\sigma$  has a torus as a dense open subset.

5.2. **Affine toric varieties.** Note: defined by monomial equations

5.3. **Points of  $U_\sigma$ .**

5.4. **The torus action.** Discuss distinguished points

5.5. **Singularities of  $U_\sigma$ .**

## 6. AFFINE GIT QUOTIENTS AS AFFINE TORIC VARIETIES

In this chapter, we introduce the affine GIT quotient and give examples. One example we give is  $\mathfrak{g} // T$  for  $G = \mathrm{GL}_2(\mathbb{C})$ . We then compute a basis of  $\mathbb{C}[\mathfrak{g}]^T$  for general  $G$ . Finally, we find generators for the invariant ring  $\mathbb{C}[\mathfrak{g}]^T$  when  $G = \mathrm{GL}_3(\mathbb{C})$ . Throughout this chapter, many technical details may be missing; our focus is on understanding specific examples which can guide our investigation of  $\mathfrak{g} // T$  for general  $G$ .

### 6.1. Algebraic groups.

### 6.2. Reductive groups.

**6.3. The affine GIT quotient.** Let  $G$  be an affine algebraic group over  $\mathbb{C}$  and  $X = \mathrm{Spec}(A)$  a complex affine variety with coordinate ring  $A$ . Suppose that  $G$  acts on  $X$ . Recall that the  $G$ -invariant functions on  $X$  are

$$A^G := \{f \in A : f(g \cdot P) = f(P) \text{ for all } g \in G \text{ and } P \in X\}.$$

**Definition 43.** *The GIT quotient is defined as*

$$X // G := \mathrm{Spec}(A^G).$$

Note that points in  $X // G$  are not necessarily in bijection with the orbits  $X/G$ , but the GIT quotient defines a quotient variety even when  $X/G$  does not have the structure of an affine variety. Let us see some examples of GIT quotients:

**Example 44.** (1) *Consider the group  $\mathbb{C}^\times$  acting on the affine space  $\mathbb{C}^2$  by*

$$t \cdot (x, y) = (tx, t^{-1}y), \quad t \in \mathbb{C}^\times, (x, y) \in \mathbb{C}^2.$$

*Suppose a polynomial  $p$  has coefficients  $a_{ij} \in \mathbb{C}$  such that*

$$p(X, Y) = \sum_{i,j} a_{ij} X^i Y^j.$$

*Then  $p$  is  $\mathbb{C}^\times$  invariant if and only if for all  $t \in \mathbb{C}^\times$ ,*

$$\sum_{i,j} a_{ij} X^i Y^j = \sum_{i,j} a_{ij} t^{i-j} X^i Y^j.$$

*This is equivalent to having  $a_{ij} \neq 0$  if and only if  $i = j$ . Thus, we have that*

$$\mathbb{C}[X, Y]^{\mathbb{C}^\times} = \mathbb{C}[XY], \quad \text{and} \quad \mathbb{C}^2 // \mathbb{C}^\times = \mathrm{Spec}(\mathbb{C}[XY]) \cong \mathbb{C}.$$

(2) *Let  $G$  be  $\mathrm{GL}_2(\mathbb{C})$  and  $T$  the maximal torus of invertible diagonal matrices in  $G$ . The action of  $G$  on its Lie algebra  $\mathfrak{g} = \mathfrak{gl}_2(\mathbb{C})$  by conjugation induces an action of  $T$  on  $\mathfrak{g}$ . Specifically, if  $\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \in T$  and  $\begin{pmatrix} x & y \\ z & w \end{pmatrix} \in \mathfrak{g}$ ,*

$$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \cdot \begin{pmatrix} x & y \\ z & w \end{pmatrix} = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \begin{pmatrix} x & y \\ z & w \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}^{-1} = \begin{pmatrix} x & ab^{-1}y \\ a^{-1}bz & w \end{pmatrix}.$$

*Let  $X \in \mathbb{C}[\mathfrak{g}]$  be the coordinate function defined by  $X \left( \begin{pmatrix} x & y \\ z & w \end{pmatrix} \right) := x$ , and define  $Y, W, Z \in \mathbb{C}[\mathfrak{g}]$  analogously. Then  $\mathbb{C}[\mathfrak{g}] = \mathbb{C}[X, Y, Z, W]$ . Since  $T$  acts trivially on the diagonal entries of an element in  $\mathfrak{g}$ ,  $X, W \in \mathbb{C}[\mathfrak{g}]^T$ . The action on the off-diagonal*

entries is analogous to the action in part (1); by the same argument that we used in part (1), a polynomial  $p$  is invariant if and only if for each monomial in  $p$ , the exponent of  $Y$  is equal to the exponent of  $Z$ . Therefore,

$$\mathbb{C}[\mathfrak{g}]^T = \mathbb{C}[X, W, YZ], \quad \text{and} \quad \mathfrak{g} // T = \text{Spec}(\mathbb{C}[X, W, YZ]) \cong \mathbb{C}^3.$$

(3) Let  $S_2 = \langle \sigma \mid \sigma^2 = 1 \rangle$  act on  $\mathbb{C}^2$  by

$$\sigma \cdot (x, y) = (y, x).$$

Polynomials which are invariant under the induced action on  $\mathbb{C}[X, Y]$  are called symmetric polynomials. We compute  $\mathbb{C}[X, Y]^{S_2}$ , i.e., the ring of symmetric polynomials in two variables; this computation is a special case of the fundamental theorem of symmetric polynomials, which characterises  $\mathbb{C}[X_1, \dots, X_n]^{S_n}$  when  $S_n$  acts on  $\mathbb{C}^n$  by permuting coordinates (see [Lan02, Chapter IV, §6] for a proof of the general theorem).

We claim that  $\mathbb{C}[X, Y]^{S_2} = \mathbb{C}[XY, X + Y]$ . It is clear that  $XY$  and  $X + Y$  are symmetric, so we just need to show any symmetric polynomial lies in  $\mathbb{C}[XY, X + Y]$ . A polynomial can be written uniquely as a sum of homogeneous polynomials, and the polynomial is symmetric if and only if each homogeneous part is. In turn, each homogeneous part is symmetric if and only if it is a  $\mathbb{C}$ -linear combination of terms of the form  $X^i Y^j + X^j Y^i$  for some  $i, j \in \mathbb{Z}_{\geq 0}$ . If  $i = j$ , then clearly  $X^i Y^j + X^j Y^i = 2(XY)^i \in \mathbb{C}[XY, X + Y]$ . Otherwise, we can assume without loss of generality  $i < j$ . Then  $X^i Y^j + X^j Y^i = (XY)^i (Y^{j-i} + X^{j-i})$ , and it suffices to show  $X^n + Y^n \in \mathbb{C}[XY, X + Y]$  for all  $n \geq 1$  to prove our claim. We proceed by induction on  $n$ ; clearly  $X + Y$  and  $X^2 + Y^2 = (X + Y)^2 - 2XY$  lie in  $\mathbb{C}[XY, X + Y]$ , so the  $n = 1$  and  $n = 2$  cases hold. Then for  $n \geq 3$ ,

$$X^n + Y^n = (X^{n-1} + Y^{n-1})(X + Y) - XY(X^{n-2} + Y^{n-2}) \in \mathbb{C}[XY, X + Y],$$

which completes the induction. It can be shown that  $XY$  and  $X + Y$  are algebraically independent [Lan02, Chapter IV, §6]. We then have that

$$\mathbb{C}^2 // S_2 = \text{Spec}(\mathbb{C}[XY, X + Y]) \cong \mathbb{C}^2.$$

(4) This example is studied in [Kam21]. Consider the group  $G = \text{GL}_2(\mathbb{C})$  acting on its Lie algebra  $\mathfrak{g} = \mathfrak{gl}_2(\mathbb{C})$  by conjugation. We use the same notation as part (2) for coordinate functions so that  $\mathbb{C}[\mathfrak{g}] = \mathbb{C}[X, Y, Z, W]$ . A polynomial is invariant if and only if it is constant on the orbits of the action. Then  $f \in \mathbb{C}[\mathfrak{g}]^G$  is determined by its values on the orbit representatives which, by Jordan normal form, we can take to be

$$\begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}, \begin{pmatrix} \lambda & 0 \\ 0 & \mu \end{pmatrix}, \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}, \quad \lambda, \mu \in \mathbb{C}.$$

We claim that  $f\left(\begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}\right) = f\left(\begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}\right)$  for  $f \in \mathbb{C}[\mathfrak{g}]^G$ . Indeed, since  $f$  is continuous and invariant,

$$\begin{aligned} f\left(\begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}\right) &= f\left(\lim_{t \rightarrow 0} \begin{pmatrix} \lambda & t \\ 0 & \lambda \end{pmatrix}\right) = \lim_{t \rightarrow 0} f\left(\begin{pmatrix} t^{-1} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \lambda & t \\ 0 & \lambda \end{pmatrix} \begin{pmatrix} t^{-1} & 0 \\ 0 & 1 \end{pmatrix}^{-1}\right) \\ &= \lim_{t \rightarrow 0} f\left(\begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}\right) = f\left(\begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}\right). \end{aligned}$$

Then  $f \in \mathbb{C}[\mathfrak{g}]^G$  is in fact determined by its restriction  $f|_{\mathfrak{h}}$ , where  $\mathfrak{h}$  is the Cartan subalgebra of diagonal matrices in  $\mathfrak{g}$ . Note that since  $\text{diag}(\lambda, \mu)$  and  $\text{diag}(\mu, \lambda)$  are in the same orbit,  $f|_{\mathfrak{h}}$  is a symmetric polynomial in the variables  $X$  and  $W$ . We then have an inclusion

$$\mathbb{C}[\mathfrak{g}]^G \hookrightarrow \mathbb{C}[X, W]^{S_2} = \mathbb{C}[XW, X + W], \quad f \mapsto f|_{\mathfrak{h}}.$$

We know from linear algebra that  $\text{tr}, \det \in \mathbb{C}[\mathfrak{g}]^G$ . Noting  $\det|_{\mathfrak{h}} = XW$  and  $\text{tr}|_{\mathfrak{h}} = X + W$ , we see the inclusion  $\mathbb{C}[\mathfrak{g}]^G \hookrightarrow \mathbb{C}[X, W]^{S_2}$  is surjective. Thus we have an isomorphism  $\mathbb{C}[\mathfrak{g}]^G \cong \mathbb{C}[X, W]^{S_2}$ . Therefore,

$$\mathfrak{g} // G = \text{Spec}(\mathbb{C}[\mathfrak{g}]^G) \cong \text{Spec}(\mathbb{C}[X, W]^{S_2}) \cong \mathbb{C}^2.$$

#### 6.4. Algebraic tori.

**6.5. The invariant ring for a torus acting on affine space.** The goal of the next two subsections is to prove that if a torus  $T$  acts linearly on an affine space  $\mathbb{A}^n$ , then the affine GIT quotient  $\mathbb{A}^n // T$  has the structure of a toric variety. With this goal in mind, in this section we will compute the ring of invariants for such an action; in particular, we will show that there is a semigroup  $\mathcal{M}$  so that the ring of invariants is isomorphic to the semigroup algebra  $\mathbb{C}[\mathcal{M}]$ . Then in the next section, we will find a lattice  $N$  and cone  $\sigma$ , such that  $S_\sigma \cong \mathcal{M}$ , which will be sufficient to show that  $\mathbb{A}^n // T$  is an affine toric variety.

Let  $T$  be an algebraic torus acting linearly on  $\mathbb{A}^n$ , i.e., suppose there is a set of characters  $S = \{\chi_1, \dots, \chi_n\} \subseteq X^*(T)$  such that

$$t \cdot (z_1, \dots, z_n) = (\chi_1(t)z_1, \dots, \chi_n(t)z_n)$$

for all  $t \in T$  and  $z = (z_1, \dots, z_n) \in \mathbb{A}^n$ . For notational convenience, we will index the coordinates of  $\mathbb{A}^n$  by the character that the torus acts by for that coordinate, instead of a number  $i = 1, \dots, n$ . Thus, a point  $z \in \mathbb{A}^n$  will be written  $z = (z_\chi)_{\chi \in S}$ , such that

$$(3) \quad t \cdot z = (\chi(t)z_\chi)_{\chi \in S}.$$

We define the polynomial  $Z_\chi$  for  $\chi \in S$  by

$$Z_\chi(z) = z_\chi, \quad \text{for } z = (z_\chi)_{\chi \in S}.$$

Then the coordinate ring of  $\mathbb{A}^n$  is  $\mathbb{C}[Z_\chi : \chi \in S]$ . Let  $(\mathbb{Z}_{\geq 0})^S = \text{Fun}(S, \mathbb{Z}_{\geq 0})$  denote the ring of functions  $S \rightarrow \mathbb{Z}_{\geq 0}$ , and for  $\eta \in (\mathbb{Z}_{\geq 0})^S$ , write  $\eta = (\eta_\chi)_{\chi \in S}$ , where  $\eta_\chi = \eta(\chi)$ . An element  $\eta \in (\mathbb{Z}_{\geq 0})^S$  defines a monomial in  $\mathbb{C}[Z_\chi : \chi \in S]$ , namely

$$Z^\eta := \prod_{\chi \in S} Z_\chi^{\eta_\chi}.$$

A polynomial  $p \in \mathbb{C}[Z_\chi : \chi \in S]$  can be written

$$p = \sum_{\eta} p_\eta Z^\eta,$$

where the sum is over all  $\eta \in (\mathbb{Z}_{\geq 0})^S$  and all but finitely many of the coefficients  $p_\eta \in \mathbb{C}$  are zero.

Let us now describe the invariant ring  $\mathbb{C}[Z_\chi : \chi \in S]^T$ . Recall that since  $T$  acts on  $\mathbb{A}^n$ , there is an induced action of  $T$  on the coordinate ring; if  $t \in T$  and  $p \in \mathbb{C}[Z_\chi : \chi \in S]$ , then

$(t \cdot p)(z) := p(t^{-1} \cdot z)$ . Using the definition of the action in equation 3, we see that  $T$  acts on a monomial  $Z^\eta$  by

$$t \cdot Z^\eta = \prod_{\chi \in S} (\chi(t^{-1}) Z_\chi)^{\eta_\chi} = \left( \sum_{\chi \in S} -\eta_\chi \chi(t) \right) Z^\eta.$$

It follows that  $p \in \mathbb{C}[Z_\chi : \chi \in S]$  is invariant for the action of  $T$  if and only if

$$\sum_{\eta} p_{\eta} Z^{\eta} = \sum_{\eta} p_{\eta} \left( \sum_{\chi \in S} -\eta_{\chi} \chi(t) \right) Z^{\eta}$$

for all  $t \in T$ . Since  $Z^\eta$  and  $Z^\mu$  are linearly independent for distinct  $\eta$  and  $\mu$ , the above equality holds if and only if

$$Z^\eta = \left( \sum_{\chi \in S} -\eta_\chi \chi(t) \right) Z^\eta$$

for all  $\eta$  such that  $p_\eta \neq 0$ . Equivalently,  $p$  is invariant for the action of  $T$  if and only if

$$\sum_{\chi \in S} \eta_\chi \chi = 0$$

for all  $\eta$  such that  $p_\eta \neq 0$ . The following lemma summarises our discussion:

**Lemma 45.** *We have that*

$$\mathbb{C}[Z_\chi : \chi \in S]^T = \mathbb{C} \left[ Z^\eta : \eta \in (\mathbb{Z}_{\geq 0})^S \text{ and } \sum_{\chi \in S} \eta_\chi \chi = 0 \right].$$

**Corollary 46.** *Let  $\mathcal{M}$  be the semigroup*

$$\mathcal{M} := \left\{ \eta \in (\mathbb{Z}_{\geq 0})^S : \sum_{\chi \in S} \eta_\chi \chi = 0 \right\}.$$

*Denoting the semigroup algebra of  $\mathcal{M}$  by  $\mathbb{C}[\mathcal{M}]$ , we have that*

$$\mathbb{C}[Z_\chi : \chi \in S]^T \cong \mathbb{C}[\mathcal{M}].$$

**6.6.  $\mathbb{A}^n // T$  as a toric variety.** We now want to find a lattice  $N$  and a cone  $\sigma$  such that  $\mathbb{A}^n // T$  is isomorphic to the affine toric variety  $U_\sigma$ . Since Corollary 46 expresses the invariant ring  $\mathbb{C}[Z_\chi : \chi \in S]^T$  as the semigroup algebra  $\mathbb{C}[\mathcal{M}]$ , it suffices to find  $N$  and  $\sigma$  such that  $S_\sigma$  is isomorphic to  $\mathcal{M}$ . We then have that

$$\mathbb{A}^n // T \cong \text{Spec}(\mathbb{C}[Z_\chi : \chi \in S]^T) \cong \text{Spec}(\mathbb{C}[S_\sigma]) = U_\sigma,$$

showing  $\mathbb{A}^n // T$  has the structure of an affine toric variety.

Let

$$\mathbb{Z}^S := \text{Fun}(S, \mathbb{Z})$$

be the lattice of integer-valued functions on  $S$ . This is a lattice in  $\mathbb{R}^S$ , the vector space of real-valued functions on  $S$ . We denote elements of  $\mathbb{Z}^S$  by  $\eta = (\eta_\chi)_{\chi \in S}$ , where  $\eta_\chi = \eta(\chi)$ . The

dual lattice is  $(\mathbb{Z}^S)^\vee$ , where elements are similarly denoted  $\mu = (\mu_\chi)_{\chi \in S} \in (\mathbb{Z}^S)^\vee$ . The dual pairing  $(\mathbb{Z}^S)^\vee \times \mathbb{Z}^S \rightarrow \mathbb{Z}$  is given by

$$(\mu, \eta) \mapsto \langle \mu, \eta \rangle := \sum_{\chi \in S} \mu_\chi \eta_\chi.$$

The indicators  $\{e_\chi\}_{\chi \in S}$ , given by  $(e_\chi)_{\chi'} = e_\chi(\chi') = \delta_{\chi, \chi'}$ , are a basis for  $\mathbb{Z}^S$ . The dual basis for  $(\mathbb{Z}^S)^\vee$  is  $\{e_\chi^\vee\}_{\chi \in S}$ , where  $\langle e_\chi^\vee, e_{\chi'} \rangle = \delta_{\chi, \chi'}$ .

We have a map  $\varphi : (\mathbb{Z}^S)^\vee \rightarrow \text{span}_\mathbb{Z}(S) \subseteq X^*(T)$  given by

$$\mu \mapsto \sum_{\chi \in S} \mu_\chi \chi.$$

The kernel of  $\varphi$  is

$$M := \ker \varphi = \left\{ \mu \in (\mathbb{Z}^S)^\vee : \sum_{\chi \in S} \mu_\chi \chi = 0 \right\}.$$

Since a subgroup of a finitely-generated free abelian group is again finitely-generated and free,  $M$  is a finitely-generated free abelian group. It has rank

$$\text{rank}(M) = \text{rank}((\mathbb{Z}^S)^\vee) - \text{rank}(\text{span}_\mathbb{Z}(S)) = |S| - \text{rank}(\text{span}_\mathbb{Z}(S)).$$

There is a corresponding sublattice of  $\mathbb{Z}^S$ ,

$$\begin{aligned} K &:= \{\eta \in \mathbb{Z}^S : \langle \mu, \eta \rangle = 0 \text{ for all } \mu \in M\} \\ &= \{\eta \in \mathbb{Z}^S : \sum_{\chi \in S} \mu_\chi \eta_\chi = 0 \text{ for all } \mu \in (\mathbb{Z}^S)^\vee \text{ such that } \sum_{\chi \in S} \mu_\chi \chi = 0\}. \end{aligned}$$

We have that  $\text{rank}(K) = \text{rank}(\text{span}_\mathbb{Z}(S))$  (You can see this must be the case once you know that  $\mathbb{Z}^S/K \cong M^\vee$ , but I don't know if there is a straightforward way to prove it.) If  $\eta \in \mathbb{Z}^S$ , we use  $\bar{\eta}$  to denote the coset of  $\eta$  in  $\mathbb{Z}^S/K$ , i.e.,  $\bar{\eta} = \eta + K \in \mathbb{Z}^S/K$ .

The following theorem describes the cone of the toric variety  $\mathbb{A}^n//T$ :

**Theorem 47.** *Let*

$$N := \mathbb{Z}^S/K,$$

and

$$\sigma := \text{span}_{\mathbb{R}_{\geq 0}} \{\bar{e}_\chi : \chi \in S\} \subseteq N_\mathbb{R}.$$

Then  $S_\sigma = \sigma^\vee \cap M \cong \mathcal{M}$ , so that  $\mathbb{A}^n//T$  is isomorphic to  $U_\sigma$ .

**Example 48.** *We consider some examples of the lattices  $N$  that arise for different choices of  $S \subseteq X^*(T)$ .*

(1) *Let  $T = \mathbb{C}^\times$  and  $S = \{\chi, -\chi\}$ , where  $\chi(t) = t$ . Explicitly, the action of  $T$  on  $\mathbb{A}^2$  is*

$$t \cdot (z_1, z_2) = (tz_1, t^{-1}z_2).$$

*Then,*

$$M = \{\mu \in (\mathbb{Z}^S)^\vee : \mu_\chi \chi + \mu_{-\chi}(-\chi) = 0\} = \{(\mu_\chi, \mu_{-\chi}) \in (\mathbb{Z}^S)^\vee : \mu_\chi = \mu_{-\chi}\}.$$

*Also,*

$$\begin{aligned} K &= \{\eta \in \mathbb{Z}^S : \mu_\chi \eta_\chi + \mu_{-\chi} \eta_{-\chi} = 0 \text{ whenever } \mu_\chi = \mu_{-\chi}\} \\ &= \{(\eta_\chi, \eta_{-\chi}) \in \mathbb{Z}^S : \eta_\chi = -\eta_{-\chi}\} \end{aligned}$$

$$= \text{span}_{\mathbb{Z}}\{e_{\chi} - e_{-\chi}\}.$$

Therefore,  $N = \mathbb{Z}^S / K$  and  $\sigma = \text{span}_{\mathbb{R}_{\geq 0}}\{\overline{e_{\chi}}, \overline{e_{-\chi}}\} \subseteq N_{\mathbb{R}}$ .

(2) Let  $T$  be a maximal torus in  $G = \text{GL}_3$ , and let  $S$  be the root system of  $G$  with respect to  $T$ . Then  $S = \Phi = \{\alpha, \beta, \alpha + \beta, -\alpha, -\beta, -(\alpha + \beta)\}$ , where  $\alpha$  and  $\beta$  are simple roots. We have that  $\text{rank}(M) = |\Phi| - \text{rank}(\text{span}_{\mathbb{Z}}(\Phi)) = 4$ , and we see

$$M = \text{span}_{\mathbb{Z}}\{e_{\alpha} + e_{-\alpha}, e_{\beta} + e_{-\beta}, e_{\alpha} + e_{\beta} + e_{-(\alpha+\beta)}, e_{\alpha+\beta} + e_{-\alpha} + e_{-\beta}\}.$$

Furthermore,  $\text{rank}(K) = \text{rank}(\text{span}_{\mathbb{Z}}(\Phi))$ , and we see

$$K = \text{span}_{\mathbb{Z}}\{e_{\alpha}^{\vee} + e_{\alpha+\beta}^{\vee} - e_{-\alpha}^{\vee} - e_{-(\alpha+\beta)}^{\vee}, e_{\beta}^{\vee} + e_{\alpha+\beta}^{\vee} - e_{-\beta}^{\vee} - e_{-(\alpha+\beta)}^{\vee}\}.$$

We have that  $N = \mathbb{Z}^{\Phi} / K$  and  $\sigma = \text{span}_{\mathbb{R}_{\geq 0}}\{\overline{e_{\alpha}} : \alpha \in \Phi\}$ .

We now prove Theorem 47. The proof relies on the following lemma, which describes the dual lattice to a sublattice:

**Lemma 49.** Let  $N_1$  and  $N_2$  be lattices such that  $N_2 \leq N_1$ . Letting  $M_i$  denote the dual lattice to  $N_i$ , we have that

$$M_2 \cong M_1 / K,$$

where  $K := \{f \in M_1 : f(n) = 0 \text{ for all } n \in N_2\}$ .

*Proof.* Let  $\phi : M_1 \rightarrow M_2$  be the restriction map  $f \mapsto f|_{N_2}$ . Then  $K = \ker \phi$ , and  $M_1 / K \cong \text{im } \phi$ , so we just need to show  $\phi$  is surjective. Let  $\{n_1, \dots, n_r\}$  be a set of generators for  $N_2$ , which we extend to a set of generators for  $N_1$ ,  $\{n_1, \dots, n_r, n_{r+1}, \dots, n_s\}$ . If  $f \in M_2$ , we can extend  $f$  to a map  $\tilde{f} \in \text{Hom}(N_1, \mathbb{Z})$  by defining  $\tilde{f}$  on the generators of  $N_1$  as

$$\tilde{f}(n_i) := \begin{cases} f(n_i) & \text{if } i = 1, \dots, r, \text{ i.e., if } n_i \in N_2, \\ 0 & \text{if } i = r+1, \dots, s, \text{ i.e., if } n_i \notin N_2. \end{cases}$$

Then  $\phi(\tilde{f}) = \tilde{f}|_{N_2} = f$  and  $\phi$  is surjective. □

*Proof of Theorem 47.* Recall that

$$M = \left\{ \mu \in (\mathbb{Z}^S)^{\vee} : \sum_{\chi \in S} \mu_{\chi} \chi = 0 \right\}, \quad K = \{ \eta \in \mathbb{Z}^S : \langle \mu, \eta \rangle = 0 \text{ for all } \mu \in M \}.$$

Then Lemma 49 implies that  $M^{\vee} \cong \mathbb{Z}^S / K =: N$ . The cone  $\sigma$  in  $N$  defined as

$$\sigma := \{\overline{e_{\chi}} : \chi \in S\} \subseteq N_{\mathbb{R}}.$$

We want to show that  $S_{\sigma} = \sigma^{\vee} \cap M$  is isomorphic to the semigroup

$$\mathcal{M} := \left\{ \eta \in (\mathbb{Z}_{\geq 0})^S : \sum_{\chi \in S} \eta_{\chi} \chi = 0 \right\}.$$

We have that

$$\sigma^{\vee} \cap M = \left\{ \mu \in (\mathbb{Z}^S)^{\vee} : \sum_{\chi \in S} \mu_{\chi} \chi = 0 \text{ and } \langle \mu, \eta \rangle = 0 \text{ for all } \eta \in \sigma \right\}$$



$$\begin{aligned}
&= \left\{ \mu \in (\mathbb{Z}^S)^\vee : \sum_{\chi \in S} \mu_\chi \chi = 0 \text{ and } \langle \mu, \overline{e_\chi} \rangle = 0 \text{ for all } \chi \in S \right\} \\
&= \left\{ \mu \in (\mathbb{Z}^S)^\vee : \sum_{\chi \in S} \mu_\chi \chi = 0 \text{ and } \mu_\chi \geq 0 \text{ for all } \chi \in S \right\} \\
&\cong \left\{ \eta \in (\mathbb{Z}_{\geq 0})^S : \sum_{\chi \in S} \eta_\chi \chi = 0 \right\} \\
&= \mathcal{M}.
\end{aligned}$$

In the second equality above, we used the fact that checking  $\mu \in (\mathbb{Z}^S)^\vee$  is nonnegative on the cone  $\sigma$  is equivalent to checking  $\mu$  is nonnegative on the generators of  $\sigma$ .  $\square$

## 7. ABSTRACT VARIETIES

In this section, we study abstract algebraic varieties — these are spaces obtained by glueing together affine varieties. An algebraic variety is the analogue in algebraic geometry of a manifold in differential geometry (these are spaces obtained by glueing open subsets of Euclidean space). Our interest in algebraic varieties is motivated by the fact that there are toric varieties which are more general than the affine ones encountered in section 5, and these general toric varieties are in particular algebraic varieties. In the next section, we will study general toric varieties using our understanding of algebraic varieties developed here.

In order to motivate the definition of algebraic varieties, we will first study projective varieties, which are an important special cases. Then we will define general algebraic varieties, and see how they are determined by glueing affine varieties.

**7.1. Motivation: projective varieties.** Let  $k$  be a field. Projective  $n$ -space over  $k$ , denoted  $\mathbb{P}_k^n$  or  $\mathbb{P}^n$ , is the quotient space

$$\mathbb{P}_k^n := (k^{n+1} \setminus \{0\}) / \sim,$$

where  $(a_0, \dots, a_n) \sim (b_0, \dots, b_n)$  if  $(a_0, \dots, a_n) = \lambda(b_0, \dots, b_n)$  for some  $\lambda \in k \setminus \{0\}$ . This space can be viewed as the set of lines in  $k^{n+1}$  which pass through 0 in the following way: if  $\ell$  is a line in  $k^{n+1}$  passing through the origin and  $(a_0, \dots, a_n)$  is any nonzero point in  $\ell$ , then the map

$$\ell \mapsto [(a_0, \dots, a_n)]$$

defines a bijection between the set of lines and  $\mathbb{P}^n$ . The map is well-defined since  $[(a_0, \dots, a_n)] = [(b_0, \dots, b_n)]$  if and only if  $(a_0, \dots, a_n) = \lambda(b_0, \dots, b_n)$  for some  $\lambda \in k \setminus \{0\}$ , and it is a bijection since for any nonzero point in  $k^{n+1}$ , there is a unique line passing through the point and the origin. The equivalence class  $[(a_0, \dots, a_n)]$  is usually denoted  $(a_0 : \dots : a_n)$ . If  $P$  is the point  $(a_0 : \dots : a_n)$ , the coordinates  $(a_0 : \dots : a_n)$  are called the homogeneous coordinates of  $P$ . Note that for any  $P$ , at least one homogeneous coordinate  $a_i$  is nonzero.

In analogy with the affine theory, after defining projective space, we should define polynomials on the space so that algebraic subsets can be defined as the zero sets of polynomials. However, the naïve approach of defining polynomials on  $\mathbb{P}^n$  to be polynomials in the homogeneous coordinates is not well-defined, since the homogeneous coordinates are not unique; we could then have that a polynomial takes different values on different homogeneous coordinates for the same point. The solution for defining algebraic subsets of  $\mathbb{P}^n$  is to use *homogeneous* polynomials.

Example: elliptic curves.

**7.2. Abstract varieties.** Recall the definitions of a sheaf of  $k$ -algebras and ringed spaces from section 5. The definition of algebraic prevariety extends the notion of an affine variety. We follow Milne's definition of an algebraic variety [Mil13].

**Definition 50.** A topological space  $V$  is called *quasicompact* if every open covering of  $V$  has a finite subcovering.

The condition to be quasicompact is often known as being compact. The convention of Bourbaki is that a compact topological space is one that is quasicompact and Hausdorff [Mil13, §2 g.]; we follow this convention.

**Definition 51.** An algebraic prevariety over  $k$  is a  $k$ -ringed space  $(V, \mathcal{O}_V)$  such that  $V$  is quasicompact and every point of  $V$  has an open neighbourhood  $U$  for which  $(U, \mathcal{O}_V|_U)$  is isomorphic to the ringed space of regular functions on an algebraic set over  $k$ .

In other words, a ringed space  $(V, \mathcal{O}_V)$  is an algebraic prevariety over  $k$  if there exists a finite open covering  $V = \bigcup V_i$  such that  $(V_i, \mathcal{O}_V|_{V_i})$  is an affine algebraic variety over  $k$  for all  $i$ .

Recall that a topological manifold is required to be Hausdorff, a condition which excludes pathological topological behaviour like non-uniqueness of limits. The analogous condition for prevarieties is called being separated. An algebraic variety will then be a separated prevariety.

**Definition 52.** An algebraic prevariety  $V$  is called separated if for every pair of regular maps  $\varphi_1, \varphi_2 : Z \rightarrow V$ , where  $Z$  is an affine algebraic variety, the set  $\{z \in Z : \varphi_1(z) = \varphi_2(z)\}$  is closed in  $Z$ .

The following lemma tells us how we can obtain a prevariety by patching together ringed spaces:

**Proposition 53.** Suppose that the set  $V$  is a finite union  $V = \bigcup V_i$  of subsets  $V_i$  and that each  $V_i$  is equipped with a ringed space structure. Assume that the following patching condition holds: for all  $i, j$ ,  $V_i \cap V_j$  is open in both  $V_i$  and  $V_j$  and  $\mathcal{O}_{V_i}|_{V_i \cap V_j} = \mathcal{O}_{V_j}|_{V_i \cap V_j}$ . Then there is a unique ringed space structure on  $V$  such that

- (1) each inclusion  $V_i \hookrightarrow V$  is a homeomorphism of  $V_i$  onto an open set, and
- (2) for each  $i$ ,  $\mathcal{O}_V|_{V_i} = \mathcal{O}_{V_i}$ .

If every  $V_i$  is an algebraic variety, then so also is  $V$ , and to give a regular map from  $V$  to a prevariety  $W$  amounts to giving a family of regular maps  $\varphi_i : V_i \rightarrow W$  such that  $\varphi_i|_{V_i \cap V_j} = \varphi_j|_{V_i \cap V_j}$ .

Example: line with two origins

**7.3. The glueing construction.** Let  $\{Y_i\}_{i \in I}$  be a finite set of affine varieties. Suppose that for all  $i, j \in I$ , we have isomorphic open subsets  $Y_{ij} \subseteq Y_i$  and  $Y_{ji} \subseteq Y_j$ . Let  $\{\phi_{ij} : Y_{ij} \rightarrow Y_{ji}\}_{i,j \in I}$  be isomorphism such that for all  $i, j, k \in I$ ,

- (1)  $\phi_{ij} = \phi_{ji}^{-1}$
- (2)  $\phi_{ij}(Y_{ij} \cap Y_{ik}) = Y_{ji} \cap Y_{jk}$ , and
- (3)  $\phi_{ik} = \phi_{jk} \circ \phi_{ij}$  on  $Y_{ij} \cap Y_{ik}$ .

The data of the affine varieties  $\{Y_i\}_{i \in I}$  and isomorphisms  $\{\phi_{ij}\}_{i,j \in I}$  is called *glueing data*. The abstract variety  $X$  which glues together the  $\{Y_i\}_{i \in I}$  varieties is a certain topological space. First, we construct the disjoint union of the varieties,  $\hat{X}$ , which is given by

$$\hat{X} := \bigsqcup_{i \in I} Y_i = \{(x, Y_i) : i \in I, x \in Y_i\}.$$

The set  $\hat{X}$  is endowed with the disjoint union topology, where by definition, a set in  $\hat{X}$  is open if it is a disjoint union of open subsets of the  $Y_i$ . To construct  $X$ , we want to identify points in  $\hat{X}$  if they belong to two isomorphic open subsets. Specifically, define an equivalence relation on  $\hat{X}$ ,  $\sim$ , by declaring  $(x, Y_i) \sim (y, Y_j)$  if  $x \in Y_{ij}$ ,  $y \in Y_{ji}$  and  $\phi_{ij}(x) = y$ . Condition (1) on the glueing isomorphisms ensures that  $\equiv$  is reflexive and symmetric, and conditions

(2) and (3) ensures it is transitive. Now define the abstract variety  $X$  to be  $\hat{X}/\sim$  with the quotient topology; this topology is called the Zariski topology on  $X$ . For each  $i \in I$ , denote

$$U_i := \{[(x, Y_i)] \in X : x \in Y_i\}.$$

This is an open set of  $X$ , and the map  $h_i : Y_i \rightarrow U_i$ ,  $y \mapsto [(y, Y_i)]$  is a homeomorphism. Then  $X$  is locally isomorphic to an affine variety.

**Example 54.** *For an example of glueing affine varieties, we consider how  $\mathbb{P}^1$  can be constructed by glueing together two copies of  $\mathbb{A}^1$ . Let  $Y_1 = \mathbb{A}^1$ ,  $Y_2 = \mathbb{A}^1$  and  $Y_{12} = k^\times \subseteq Y_1$ ,  $Y_{21} = k^\times \subseteq Y_2$ . Define the glueing isomorphisms*

$$\phi_{ij} : Y_{ij} \rightarrow Y_{ji}, \quad t \mapsto t^{-1}.$$

*It is clear that  $\phi_{12} = \phi_{21}^{-1}$  so that axiom (1) for the glueing isomorphisms holds (axioms (2) and (3) are vacuously true). Then, the variety obtained by glueing  $Y_1$  and  $Y_2$  is*

$$X = Y_1 \sqcup Y_2 / \sim,$$

*where  $(x, Y_1) \sim (y, Y_2)$  if  $x \neq 0$  and  $y = x^{-1}$ . To see that  $X$  is  $\mathbb{P}^1$ , we can think of the open sets*

$$U_1 = \{[(a, Y_1)] : a \in Y_1\}, \quad U_2 = \{[(b, Y_2)] : b \in Y_2\}$$

*as the usual affine charts for  $\mathbb{P}^1$ ; these are*

$$U_x := \{(x : y) : x \neq 0\}, \quad U_y := \{(x, y) : y \neq 0\},$$

*and the maps*

$$U_x \rightarrow U_1, (x : y) \mapsto [(x/y, Y_1)], \quad U_y \rightarrow U_2, (x : y) \mapsto [(y/x, Y_2)]$$

*are homeomorphisms. Observe that if  $(x : y) \in U_x \cap U_y$ , the image of  $(x : y)$  in  $X$  of the above two maps are points which are identified.*

Example: line with two origins

## 8. ABSTRACT TORIC VARIETIES

In this section, we present the general definition of an abstract toric variety. While affine toric varieties correspond to cones in a vector space, these abstract toric varieties correspond to a collection of cones which “fit together” in a nice way — these collections of cones are called fans.

**8.1. Fans.** Given a certain collection of cones called a fan, one constructs an abstract toric variety by glueing affine toric varieties  $U_\sigma$  for cones  $\sigma$  in the fan. We now define a fan, and demonstrate how it encodes the glueing data. As in chapter 5, let  $N$  be a lattice,  $M$  the dual lattice, and  $N_{\mathbb{R}}$  and  $M_{\mathbb{R}}$  the corresponding vector spaces.

**Definition 55** ([Ful93, §1.4]). *A fan  $\Delta$  in  $N$  is a set of rational strongly convex polyhedral cones  $\sigma$  in  $N_{\mathbb{R}}$  such that*

- (1) *each face of a cone in  $\Delta$  is also a cone in  $\Delta$ , and*
- (2) *the intersection of two cones in  $\Delta$  is a face of each.*

For simplicity, we will assume that fans only contain a finite number of cones. The idea of constructing the toric variety corresponding to  $\Delta$  is that we consider each cone  $\sigma$ , and glue together the affine toric varieties  $U_\sigma$ . The following lemma shows us that if  $\tau$  is a face of a cone  $\sigma$ , then  $U_\tau$  is an open subset of  $U_\sigma$ ; this will allow us to glue the affine toric varieties corresponding to cones in a fan.

Recall that a homomorphism of semigroups  $S \rightarrow S'$  induces an algebra homomorphism  $\mathbb{C}[S] \rightarrow \mathbb{C}[S']$  and hence a morphism  $\text{Spec}(\mathbb{C}[S]) \rightarrow \text{Spec}(\mathbb{C}[S'])$ . In particular, if  $\tau$  is contained in  $\sigma$ , there is an inclusion  $\sigma^\vee \hookrightarrow \tau^\vee$  which determines a morphism  $U_\tau \rightarrow U_\sigma$ .

**Proposition 56** ([Ful93, §1.3]). *If  $\tau$  is a face of  $\sigma$ , then the map  $U_\tau \rightarrow U_\sigma$  embeds  $U_\tau$  as a principal open subset of  $U_\sigma$ .*

We need the following lemma from convex geometry:

**Lemma 57** ([Ful93, §1.2]). *Let  $\sigma$  be a rational convex polyhedral cone, and suppose  $u \in S_\sigma$ . Then  $\tau = \sigma \cap u^\perp$  is rational convex polyhedral cone. All faces of  $\sigma$  have this form, and*

$$S_\tau = S_\sigma + \mathbb{Z}_{\geq 0} \cdot (-u).$$

*Proof. To do: This is Fulton’s proof, which is sparse on details. It would be good to write this clearer.* If  $\tau$  is a face, then  $\tau = \sigma \cap u^\perp$  for any  $u$  in the relative interior of  $\sigma^\vee \cap \tau^\perp$ , and  $u$  can be taken to be in  $M$  since  $\sigma^\vee \cap \tau^\perp$  is rational. Given  $w \in S_\tau$ , then  $w + p \cdot u \in \sigma^\vee$  for large positive  $p$ , and taking  $p$  to be an integer shows that  $w \in S_\sigma + \mathbb{Z}_{\geq 0} \cdot (-u)$ .  $\square$

*Proof of Proposition 56.* The lemma yields  $u \in S_\sigma$  such that  $\tau = \sigma \cap u^\perp$  and

$$S_\tau = S_\sigma + \mathbb{Z}_{\geq 0} \cdot (-u).$$

Then any basis element in  $\mathbb{C}[S_\tau]$  can be written as  $\chi^{w-pu} = \chi^w / (\chi^u)^p$  for some  $w \in S_\sigma$  and  $p \in \mathbb{Z}_{\geq 0}$ . Then,  $\mathbb{C}[S_\tau] = \mathbb{C}[S_\sigma]_{\chi^u}$ , which is the algebraic version of the assertion.  $\square$

**8.2. Abstract toric varieties.** Let  $\Delta$  be a fan in  $N$ .

**Definition 58.** *The toric variety  $X(\Delta)$  is constructed by glueing the affine toric varieties  $U_\sigma$ , as  $\sigma$  ranges over all elements of  $\Delta$ . For cones  $\sigma, \tau \in \Delta$ , the intersection  $\sigma \cap \tau$  is a face of each and hence  $U_{\sigma \cap \tau}$  is isomorphic to a principal open subset of each  $U_\sigma$  and  $U_\tau$ , and these varieties can be glued along this open subvariety.*

The fact that the glueing is compatible follows from the order-preserving nature of the correspondence between cones and affine varieties. *To do: Understand this better. Can we give explicit glueing maps  $\phi_{ij}$  as we used in the definition of glueing?*

Example: one-dim toric varieties

8.3. **The dense torus.**

8.4. **The orbit-cone correspondence.**

8.5. **Classification of toric surfaces.**

8.6. **Toric morphisms.**

## REFERENCES

- [BV04] Stephen Boyd and Lieven Vandenbergh, *Convex optimization*, Cambridge University Press, Cambridge, 2004. MR2061575
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011. MR2810322
- [DLHK13] Jesús A. De Loera, Raymond Hemmecke, and Matthias Köppe, *Algebraic and geometric ideas in the theory of discrete optimization*, MOS-SIAM Series on Optimization, vol. 14, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA; Mathematical Optimization Society, Philadelphia, PA, 2013. MR3024570
- [Dol03] Igor Dolgachev, *Lectures on invariant theory*, London Mathematical Society Lecture Note Series, vol. 296, Cambridge University Press, Cambridge, 2003. MR2004511
- [Ful93] William Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993. The William H. Roever Lectures in Geometry. MR1234037
- [Gec13] Meinolf Geck, *An introduction to algebraic geometry and algebraic groups*, Oxford Graduate Texts in Mathematics, vol. 20, Oxford University Press, Oxford, 2013. First paperback reprinting of the 2003 original. MR3086289
- [Har77] Robin Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, vol. No. 52, Springer-Verlag, New York-Heidelberg, 1977. MR0463157
- [Hil90] David Hilbert, *Ueber die Theorie der algebraischen Formen*, Math. Ann. **36** (1890), no. 4, 473–534, DOI 10.1007/BF01208503 (German). MR1510634
- [Hum72] James E. Humphreys, *Introduction to Lie algebras and representation theory*, Graduate Texts in Mathematics, vol. Vol. 9, Springer-Verlag, New York-Berlin, 1972. MR0323842
- [Hum75] ———, *Linear algebraic groups*, Graduate Texts in Mathematics, vol. No. 21, Springer-Verlag, New York-Heidelberg, 1975. MR0396773
- [Hum90] ———, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, vol. 29, Cambridge University Press, Cambridge, 1990. MR1066460
- [Kam21] Masoud Kamgarpour, *UQ Maths Stradbroke Island Workshop on Character Varieties* (2021). Available at [https://baileywhitbread.com/files/21\\_character\\_varieties.pdf](https://baileywhitbread.com/files/21_character_varieties.pdf).
- [Lan02] Serge Lang, *Algebra*, 3rd ed., Graduate Texts in Mathematics, vol. 211, Springer-Verlag, New York, 2002. MR1878556
- [Mil13] James S. Milne, *Lie Algebras, Algebraic Groups, and Lie Groups* (2013). Available at [www.jmilne.org/math/](http://www.jmilne.org/math/).
- [Muk03] Shigeru Mukai, *An introduction to invariants and moduli*, Japanese edition, Cambridge Studies in Advanced Mathematics, vol. 81, Cambridge University Press, Cambridge, 2003. MR2004218
- [Mum65] David Mumford, *Geometric invariant theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, (N.F.), vol. Band 34, Springer-Verlag, Berlin-New York, 1965. MR0214602
- [Nag60] Masayoshi Nagata, *On the fourteenth problem of Hilbert*, Proc. Internat. Congress Math. 1958, Cambridge Univ. Press, New York, 1960, pp. 459–462. MR0116056
- [Oda88] Tadao Oda, *Convex bodies and algebraic geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 15, Springer-Verlag, Berlin, 1988. An introduction to the theory of toric varieties; Translated from the Japanese. MR0922894
- [Rei88] Miles Reid, *Undergraduate algebraic geometry*, London Mathematical Society Student Texts, vol. 12, Cambridge University Press, Cambridge, 1988. MR0982494
- [Rei95] ———, *Undergraduate commutative algebra*, London Mathematical Society Student Texts, vol. 29, Cambridge University Press, Cambridge, 1995. MR1458066