


Toric Varieties

Bridges in Algebra and Geometry

Declan Fletcher

October 2024

Two bridges




Overview

1. Affine varieties
2. Polynomial maps
3. The Nullstellensatz
4. Irreducibility and prime ideals
5. Singularities
6. Convex cones
7. Toric varieties
8. Faces and open subsets
9. Singularities of toric varieties

Affine varieties

Let k be a field. The set of n -tuples of elements of k is called affine space, \mathbb{A}^n . Given a polynomial f in $k[X_1, \dots, X_n]$, we can consider its zero set

$$\mathbf{V}(f) := \{(a_1, \dots, a_n) \in \mathbb{A}^n : f(a_1, \dots, a_n) = 0\}.$$



We can also consider spaces cut out by sets of polynomials $I \subseteq k[X_1, \dots, X_n]$:

$$\mathbf{V}(I) := \{(a_1, \dots, a_n) \in \mathbb{A}^n : f(a_1, \dots, a_n) = 0 \text{ for all } f \in I\}.$$

These spaces are called **affine varieties**.

Varieties throughout mathematics

Choose $k = \mathbb{R}$
and consider $\nabla(x^2 + y^2 + z^2 - 1)$



Choose $k = \mathbb{Q}$
and consider $\nabla(x^2 + y^2 - 1)$

Integer solutions
to $(\frac{a}{c})^2 + (\frac{b}{c})^2 = 1$.

The Zariski topology


Affine space is endowed with the **Zariski topology**.

This is the topology defined by declaring $S \subseteq \mathbb{A}^n$ closed if

$$S = \mathbf{V}(I)$$

for some $I \subseteq k[X_1, \dots, X_n]$.

The Zariski topology on a variety V is the induced topology from \mathbb{A}^n .



Polynomial functions

We say a function $V \rightarrow k$ is **polynomial** if it is the restriction of a polynomial function $\mathbb{A}^n \rightarrow k$.

Problem: different polynomials give the same function. For example, on the circle $V = \mathbf{V}(X^2 + Y^2 - 1)$, the polynomials

$$2XY^2, \quad 2XY^2 - 2X(X^2 + Y^2 - 1)$$

define the same function.

Solution: consider functions in the quotient

$$k[V] := k[X_1, \dots, X_n]/\mathbf{I}(V),$$

where

$$\mathbf{I}(V) := \{\text{polynomials vanishing on } V\}.$$


We call $k[V]$ the **coordinate ring** of V , and $\mathbf{I}(V)$ the **ideal** of V .

Polynomial maps

A map $\varphi : V \subseteq \mathbb{A}^n \rightarrow W \subseteq \mathbb{A}^m$ is called **polynomial** if its components are polynomial functions on V , i.e., if

$$\varphi = (\varphi_1, \dots, \varphi_m)$$

for some $\varphi_i \in k[V]$.



We say φ is an isomorphism if it is bijective with polynomial inverse.

The Nullstellensatz

We have maps

$$\{ \text{varieties in } \mathbb{A}^n \} \xrightleftharpoons[\vee]{I} \{ \text{ideals in } k[X_1, \dots, X_n] \}.$$

The Nullstellensatz (Moral)

Let k be **algebraic closed**. There is a bijection

$$\{ \text{varieties in } \mathbb{A}^n \} \leftrightarrow \{ \text{radical ideals in } k[X_1, \dots, X_n] \}.$$


An ideal I is called radical if whenever $f^n \in I$ for some $n \in \mathbb{Z}_{>0}$, we have $f \in I$. These are ideals containing all their square roots, cube roots, etc.

Henceforth, we assume k is algebraically closed.

Maps and homomorphisms

A polynomial map $\varphi : V \rightarrow W$ gives us a rule for turning polynomial functions on W into polynomial functions on V . Specifically, it gives a ring homomorphism

$$\varphi^* : k[W] \rightarrow k[V], \quad f \mapsto \varphi^* f := f \circ \varphi.$$




The map $\varphi \mapsto \varphi^*$ is in fact a bijection

$$\left\{ \text{poly. maps } V \rightarrow W \right\} \leftrightarrow \left\{ \text{ring homs } k[W] \rightarrow k[V] \right\}.$$

Moreover, φ is an isomorphism if and only if φ^* is. Then V and W are isomorphic if and only if $k[V]$ and $k[W]$ are.

Irreducibility and prime ideals

We say a variety is **irreducible** if it is not the union of two proper closed subsets.



The ideal $\mathbf{I}(V)$ is **prime** if and only if $k[V] = k[X_1, \dots, X_n]/\mathbf{I}(V)$ has no zero divisors.

Proposition

The variety V is irreducible if and only if the ideal $\mathbf{I}(V)$ is prime.

Tangent spaces and dimension

Tangent spaces are an important tool for studying smooth surfaces in differential geometry. There is an analog for varieties.

We define tangent spaces for the hypersurface $V = \mathbf{V}(f)$. Let $P = (a_1, \dots, a_n)$ be a point in V . The first-order part of f at P is

$$f_P^{(1)} := \sum_{i=1}^n \frac{\partial f}{\partial X_i}(P)(X_i - a_i).$$

The **tangent space** of V at P is

$$T_P V := \mathbf{V}(f_P^{(1)}).$$

The **dimension** of V is



$$\dim(V) := \min_{P \in V} \dim(T_P V).$$

Singular and non-singular points

We say a point P is **singular** if

$$\dim(T_P V) > \dim(V),$$

and non-singular otherwise.



Convex cones


A **polyhedral cone** in the vector space \mathbb{R}^n is a set of the form

$$\sigma = \text{span}_{\mathbb{R}_{\geq 0}}\{v_1, \dots, v_r\},$$

for some vectors v_1, \dots, v_r in \mathbb{Z}^n .

The **dual cone** to a polyhedral cone σ is

$$\sigma^\vee := \{u \in (\mathbb{R}^n)^*: u(v) \geq 0 \text{ for all } v \in \sigma\}.$$



Cones and their duals



A trivial (but important) example: if $\sigma = \{0\}$, then $\sigma^\vee = (\mathbb{R}^n)^*$.

For a non-trivial example, consider

$$\sigma = \text{span}_{\mathbb{R}_{\geq 0}} \{e_1, -e_1 + 2e_2\}.$$


Then,

$$\sigma^\vee = \text{span}_{\mathbb{R}_{\geq 0}} \{2e_1 + e_2, e_2\}.$$



Toric varieties, the intuition

Given a cone and its dual, we place a monomial $X^i Y^j$ on the integer points (i, j) in the dual space:



We form a ring using the monomials lying in the dual cone:

$$k[1, Y, XY, X^2Y, Y^2, XY^2, \dots] = k[Y, XY, X^2Y].$$

The toric variety U_σ is defined by taking this as its coordinate ring:

$$U_\sigma := \text{Spec}(k[Y, XY, X^2Y]) = \text{Spec}(k[R, S, T]/(RT - S^2)).$$

Toric varieties, the definition

The integer points in σ^\vee form a **semigroup**,

$$S_\sigma := \sigma^\vee \cap (\mathbb{Z}^n)^*.$$

We form the **semigroup algebra** $k[S_\sigma]$. This has the basis of formula symbols

$$\{\chi^u : u \in S_\sigma\}$$

with multiplication

$$\chi^u \chi^{u'} = \chi^{u+u'}.$$

The **toric variety** associated with σ is defined as

$$U_\sigma := \text{Spec}(k[S_\sigma]).$$

The torus in toric varieties

When $\sigma = \{0\}$, we know $\sigma^\vee = (\mathbb{R}^n)^*$. The semigroup S_σ is

$$S_\sigma = (\mathbb{R}^n)^* \cap (\mathbb{Z}^n)^* = (\mathbb{Z}^n)^*.$$

We see

$$\begin{aligned} k[S_\sigma] &= k[\chi^{e_1^*}, \chi^{-e_1^*}, \dots, \chi^{e_n^*}, \chi^{-e_n^*}] \\ &= k[X_1, X_1^{-1}, \dots, X_n, X_n^{-1}] \end{aligned}$$


Then

$$U_\sigma = \text{Spec}(k[X_1, X_1^{-1}, \dots, X_n, X_n^{-1}]) = (k^\times)^n$$

is the **algebraic torus**.

Faces

A **face** of a cone σ is the intersection of σ with a hyperplane which contains σ in a half-space.



If τ is a face of σ , then $\tau \subseteq \sigma$ but $\tau^\vee \supseteq \sigma^\vee$ and $S_\tau \supseteq S_\sigma$. We then have an inclusion

$$k[S_\sigma] \hookrightarrow k[S_\tau].$$

Corresponding to this, there is a map


$$U_\tau \rightarrow U_\sigma.$$

Faces and open subsets

Theorem

Let τ be a face of σ . Then the map $U_\tau \rightarrow U_\sigma$ embeds U_τ as a principal open subset of U_σ .

Smaller faces correspond to smaller open subsets.



Singularities of toric varieties


Theorem

An toric variety U_σ is non-singular if and only if σ is generated by a subset of a basis for \mathbb{Z}^n . In this case,

$$U_\sigma \cong k^r \times (k^\times)^{n-r},$$

where $r = \dim(\sigma)$.

Cones detect singularities



References

- David Cox, John Little, and Henry Schenck, *Toric varieties*, American Mathematical Society, 2011.
- William Fulton, *Introduction to toric varieties*, Princeton University Press, 1993.
- James Milne, *Algebraic Geometry* (2023), www.jmilne.org/math/.
- Miles Reid, *Undergraduate algebraic geometry*, Cambridge Univeristy Press, 1988.