

# Toric Varieties

---

Declan Fletcher

October 2024

# What are toric varieties, and why study them?

Algebraic varieties are geometric spaces defined by solutions to polynomial equations.


In general, these are complicated and difficult to study.

Toric varieties are a special class of varieties determined by a convex cone.

# Algebraic varieties




$$x^2 + y^2 = 1$$



$$x^2 + y^2 + z^2 = 1$$



$$y^2 = x^3 + 2x^2$$



$$xy = z^2$$

# The definition of a variety

Varieties are sets of solutions  $(a_1, \dots, a_n) \in \mathbb{C}^n$  to poly. equations

$$f_1(a_1, \dots, a_n) = 0, \quad \dots, \quad f_s(a_1, \dots, a_n) = 0.$$

Choose the zero polynomial:


$$\rightsquigarrow \mathbb{C}^n.$$

Choose  $y^2 = x^3 + 2x^2$ :

$\rightsquigarrow$  singular curve.

Choose  $xy = 1$ :

$$\rightsquigarrow \{(t, t^{-1}) : t \in \mathbb{C}^\times\} \cong \mathbb{C}^\times.$$



$\mathbb{C}^\times$  is called an algebraic **torus**. The  $d$ -dimensional torus is  $(\mathbb{C}^\times)^d$ .



# Convex cones

To understand toric varieties, we need to understand convex cones.

We consider **polyhedral cones**. These are sets in  $\mathbb{R}^n$  of the form

$$\sigma = \text{span}_{\mathbb{R}_{\geq 0}} \{v_1, \dots, v_r\},$$


for some  $v_1, \dots, v_r \in \mathbb{R}^n$ .  $\sigma$  is **rational** if we can take each  $v_i \in \mathbb{Z}^n$ .



# Dual cones


**Fix:** cone  $\sigma$ . The **dual cone** is the set of linear functionals which are non-negative on  $\sigma$ :

$$\sigma^\vee := \{u \in (\mathbb{R}^n)^* : u(v) \geq 0 \text{ for all } v \in \sigma\}.$$



## An example of a toric variety

**Fix:** cone  $\sigma$ , dual  $\sigma^\vee$ . Monomials live on integer points in  $(\mathbb{R}^n)^*$ :



We create an algebra using the monomials in  $\sigma^\vee$ :

$$\begin{aligned}\mathbb{C}[1, s, r^2s, rs, s^2, rs^2, \dots] &= \mathbb{C}[s, r^2s, rs] \\ &\cong \mathbb{C}[x, y, z]/(xy - z^2).\end{aligned}$$

The toric variety  $U_\sigma$  is the set of solutions to  $xy - z^2 = 0$  in  $\mathbb{C}^3$ :

$$xy = z^2.$$

# The definition of a toric variety

**Fix:** cone  $\sigma$ , dual  $\sigma^\vee$ . The previous construction generalises.

We associate monomials  $x_1^{i_1} \cdots x_n^{i_n}$  to integer points in  $(\mathbb{R}^n)^*$ .

We create an algebra using the monomials lying in  $\sigma^\vee$ , called  $\mathbb{C}[S_\sigma]$ .

$\mathbb{C}[S_\sigma]$  is finitely generated, so it's given by generators and relations:

$$\mathbb{C}[S_\sigma] = \mathbb{C}[y_1, \dots, y_m] / (f_1, \dots, f_s).$$

The **toric variety**  $U_\sigma$  is the subset of  $\mathbb{C}^m$  defined by the equations


$$f_1(a_1, \dots, a_m) = 0, \quad \dots, \quad f_s(a_1, \dots, a_m) = 0.$$




# Cones detect singularities

## Theorem

*The toric variety  $U_\sigma$  is non-singular if and only if  $\sigma$  is generated by a subset of a basis for  $\mathbb{Z}^n$ .*



$$xy = z^2$$





## References

Stephen Boyd and Lieven Vandenberghe, *Convex optimization*, Cambridge University Press, 2004.

William Fulton, *Introduction to toric varieties*, Princeton University Press, 1993.

Shigeru Mukai, *An introduction to invariants and moduli*, Cambridge University Press, 2003.

Miles Reid, *Undergraduate algebraic geometry*, Cambridge University Press, 1988.