

Toric Varieties

Declan Fletcher

October 2024

Algebraic varieties




$$x^2 + y^2 = 1$$

Algebraic varieties




$$x^2 + y^2 = 1$$




$$x^2 + y^2 + z^2 = 1$$

Algebraic varieties



$$x^2 + y^2 = 1$$



$$x^2 + y^2 + z^2 = 1$$




$$y^2 = x^3 + 2x^2$$

Algebraic varieties




$$x^2 + y^2 = 1$$



$$x^2 + y^2 + z^2 = 1$$



$$y^2 = x^3 + 2x^2$$



$$xy = z^2$$

What are toric varieties, and why study them?

What are toric varieties, and why study them?

Toric varieties are a special class of varieties determined by elementary convex geometry.

What are toric varieties, and why study them?

Toric varieties are a special class of varieties determined by elementary convex geometry.

They serve as testing grounds for conjectures and theorems in algebraic geometry.

What are toric varieties, and why study them?

Toric varieties are a special class of varieties determined by elementary convex geometry.

They serve as testing grounds for conjectures and theorems in algebraic geometry.

For example, the Hodge conjecture—a Millennium Prize Problem—is proven for toric varieties but unresolved in general.

What are toric varieties, and why study them?

Toric varieties are a special class of varieties determined by elementary **convex** geometry.

They serve as testing grounds for conjectures and theorems in algebraic geometry.

For example, the **Hodge conjecture**—a Millennium Prize Problem—is proven for toric varieties but unresolved in general.

Goal of the talk: introduce toric varieties and explain some of their properties.

The definition of a variety

Varieties are sets of solutions $(a_1, \dots, a_n) \in \mathbb{C}^n$ to poly. equations

$$f_1(a_1, \dots, a_n) = 0, \quad \dots, \quad f_s(a_1, \dots, a_n) = 0.$$

The definition of a variety

Varieties are sets of solutions $(a_1, \dots, a_n) \in \mathbb{C}^n$ to poly. equations

$$f_1(a_1, \dots, a_n) = 0, \quad \dots, \quad f_s(a_1, \dots, a_n) = 0.$$

Choose the zero polynomial:

$$\rightsquigarrow \mathbb{C}^n.$$

The definition of a variety

Varieties are sets of solutions $(a_1, \dots, a_n) \in \mathbb{C}^n$ to poly. equations


$$f_1(a_1, \dots, a_n) = 0, \quad \dots, \quad f_s(a_1, \dots, a_n) = 0.$$

Choose the zero polynomial:

$$\rightsquigarrow \mathbb{C}^n.$$

Choose $y^2 = x^3 + 2x^2$:

\rightsquigarrow singular curve.



The definition of a variety

Varieties are sets of solutions $(a_1, \dots, a_n) \in \mathbb{C}^n$ to poly. equations

$$f_1(a_1, \dots, a_n) = 0, \quad \dots, \quad f_s(a_1, \dots, a_n) = 0.$$

Choose the zero polynomial:


$$\rightsquigarrow \mathbb{C}^n.$$

Choose $y^2 = x^3 + 2x^2$:

\rightsquigarrow singular curve.

Choose $xy = 1$:

$$\rightsquigarrow \{(t, t^{-1}) : t \in \mathbb{C}^\times\} \cong \mathbb{C}^\times.$$



The definition of a variety

Varieties are sets of solutions $(a_1, \dots, a_n) \in \mathbb{C}^n$ to poly. equations

$$f_1(a_1, \dots, a_n) = 0, \quad \dots, \quad f_s(a_1, \dots, a_n) = 0.$$

Choose the zero polynomial:


$$\rightsquigarrow \mathbb{C}^n.$$

Choose $y^2 = x^3 + 2x^2$:

\rightsquigarrow singular curve.

Choose $xy = 1$:

$$\rightsquigarrow \{(t, t^{-1}) : t \in \mathbb{C}^\times\} \cong \mathbb{C}^\times.$$



\mathbb{C}^\times is called the algebraic **torus**. The d -dimensional torus is $(\mathbb{C}^\times)^d$.

To study toric varieties, we need to understand convex cones.

Convex cones

To study toric varieties, we need to understand convex cones.

We consider **polyhedral cones**. These are sets in \mathbb{R}^n of the form

$$\sigma = \operatorname{span}_{\mathbb{R}_{\geq 0}} \{v_1, \dots, v_r\},$$

for some $v_1, \dots, v_r \in \mathbb{R}^n$.



Convex cones

To study toric varieties, we need to understand convex cones.

We consider **polyhedral cones**. These are sets in \mathbb{R}^n of the form

$$\sigma = \text{span}_{\mathbb{R}_{\geq 0}}\{v_1, \dots, v_r\},$$

for some $v_1, \dots, v_r \in \mathbb{R}^n$.



Convex cones



To study toric varieties, we need to understand convex cones.

We consider **polyhedral cones**. These are sets in \mathbb{R}^n of the form

$$\sigma = \operatorname{span}_{\mathbb{R}_{\geq 0}} \{v_1, \dots, v_r\},$$

for some $v_1, \dots, v_r \in \mathbb{R}^n$.

We call σ **rational** if we can take each $v_i \in \mathbb{Z}^n$.



Fix: cone σ .

Dual cones


Fix: cone σ . The **dual cone** is the set of linear functionals which are non-negative on σ :

$$\sigma^\vee := \{u \in (\mathbb{R}^n)^* : u(v) \geq 0 \text{ for all } v \in \sigma\}.$$

Dual cones

Fix: cone σ . The **dual cone** is the set of linear functionals which are non-negative on σ :

$$\sigma^\vee := \{u \in (\mathbb{R}^n)^* : u(v) \geq 0 \text{ for all } v \in \sigma\}.$$




An example of a toric variety

Fix: cone σ , dual σ^\vee .


An example of a toric variety

Fix: cone σ , dual σ^\vee . Monomials live on integer points in $(\mathbb{R}^n)^*$:



An example of a toric variety

Fix: cone σ , dual σ^\vee . Monomials live on integer points in $(\mathbb{R}^n)^*$:




We create a ring using the monomials in σ^\vee :

$$\begin{aligned}\mathbb{C}[1, s, r^2s, rs, s^2, rs^2, \dots] &= \mathbb{C}[s, r^2s, rs] \\ &\cong \mathbb{C}[x, y, z]/(xy - z^2).\end{aligned}$$

An example of a toric variety

Fix: cone σ , dual σ^\vee . Monomials live on integer points in $(\mathbb{R}^n)^*$:



We create a ring using the monomials in σ^\vee :

$$\begin{aligned}\mathbb{C}[1, s, r^2s, rs, s^2, rs^2, \dots] &= \mathbb{C}[s, r^2s, rs] \\ &\cong \mathbb{C}[x, y, z]/(xy - z^2).\end{aligned}$$

The toric variety U_σ is the set of solutions to $xy - z^2 = 0$ in \mathbb{C}^3 :

$$xy = z^2.$$

The definition of a toric variety

Fix: cone σ , dual σ^\vee . The previous construction generalises.

The definition of a toric variety

Fix: cone σ , dual σ^\vee . The previous construction generalises.

We associate monomials $x_1^{i_1} \cdots x_n^{i_n}$ to integer points in $(\mathbb{R}^n)^*$.

The definition of a toric variety

Fix: cone σ , dual σ^\vee . The previous construction generalises.

We associate monomials $x_1^{i_1} \cdots x_n^{i_n}$ to integer points in $(\mathbb{R}^n)^*$.

We create a ring using the monomials lying in σ^\vee , called R_σ .

The definition of a toric variety

Fix: cone σ , dual σ^\vee . The previous construction generalises.

We associate monomials $x_1^{i_1} \cdots x_n^{i_n}$ to integer points in $(\mathbb{R}^n)^*$.

We create a ring using the monomials lying in σ^\vee , called R_σ .

R_σ is finitely generated, so it's given by generators and relations:

$$R_\sigma = \mathbb{C}[y_1, \dots, y_m] / (f_1, \dots, f_s).$$

The definition of a toric variety

Fix: cone σ , dual σ^\vee . The previous construction generalises.

We associate monomials $x_1^{i_1} \cdots x_n^{i_n}$ to integer points in $(\mathbb{R}^n)^*$.

We create a ring using the monomials lying in σ^\vee , called R_σ .

R_σ is finitely generated, so it's given by generators and relations:

$$R_\sigma = \mathbb{C}[y_1, \dots, y_m] / (f_1, \dots, f_s).$$

The **toric variety** U_σ is the subset of \mathbb{C}^m defined by the equations

$$f_1(a_1, \dots, a_m) = 0, \quad \dots, \quad f_s(a_1, \dots, a_m) = 0.$$

Cones detect singularities

Cones detect singularities


Theorem

The toric variety U_σ is non-singular if and only if σ is generated by a subset of a basis for \mathbb{Z}^n .


Cones detect singularities

Theorem

The toric variety U_σ is non-singular if and only if σ is generated by a subset of a basis for \mathbb{Z}^n .




$$xy = z^2$$




Cones detect singularities

Theorem

The toric variety U_σ is non-singular if and only if σ is generated by a subset of a basis for \mathbb{Z}^n .



$$xy = z^2$$




Torus quotients

Suppose $T = (\mathbb{C}^\times)^d$ acts linearly on \mathbb{C}^n . **Goal:** understand \mathbb{C}^n/T .

Torus quotients

Suppose $T = (\mathbb{C}^\times)^d$ acts linearly on \mathbb{C}^n . **Goal:** understand \mathbb{C}^n/T .


Example: $\mathbb{C}^\times \curvearrowright \mathbb{C}^2$ by $t \cdot (x, y) = (tx, t^{-1}y)$.



Torus quotients

Suppose $T = (\mathbb{C}^\times)^d$ acts linearly on \mathbb{C}^n . **Goal:** understand \mathbb{C}^n/T .

Example: $\mathbb{C}^\times \curvearrowright \mathbb{C}^2$ by $t \cdot (x, y) = (tx, t^{-1}y)$.



Problem: determine invariant polynomials

$$\mathbb{C}[x, y]^T = \{\text{polys } f : f(p) = f(T \cdot p)\}.$$

Torus quotients as toric varieties

Torus quotients as toric varieties

Theorem

\mathbb{C}^n/T is a toric variety whose cone can be explicitly computed.

Idea:

Torus quotients as toric varieties

Theorem

\mathbb{C}^n/T is a toric variety whose cone can be explicitly computed.

Idea: Invariant monomials are parametrised by points in \mathbb{Z}^n satisfying linear equations.


Torus quotients as toric varieties

Theorem

\mathbb{C}^n/T is a toric variety whose cone can be explicitly computed.

Idea: Invariant monomials are parametrised by points in \mathbb{Z}^n satisfying linear equations. For $\mathbb{C}^\times \curvearrowright \mathbb{C}^2$:

$$x^k y^\ell = t^{k-\ell} x^k y^\ell \quad \Longleftrightarrow \quad k = \ell.$$



Torus quotients as toric varieties


Theorem

\mathbb{C}^n/T is a toric variety whose cone can be explicitly computed.

Idea: Invariant monomials are parametrised by points in \mathbb{Z}^n satisfying linear equations. For $\mathbb{C}^\times \curvearrowright \mathbb{C}^2$:

$$x^k y^\ell = t^{k-\ell} x^k y^\ell \iff k = \ell.$$

Choose σ so σ^\vee only sees points with non-negative coordinates.



References

Stephen Boyd and Lieven Vandenberghe, *Convex optimization*, Cambridge University Press, 2004.

William Fulton, *Introduction to toric varieties*, Princeton University Press, 1993.

Shigeru Mukai, *An introduction to invariants and moduli*, Cambridge University Press, 2003.

Miles Reid, *Undergraduate algebraic geometry*, Cambridge University Press, 1988.