

Marine Fish and Invertebrate Atlas: Summarizing Geographic Distribution and Population Indices in the Scotian Shelf and Bay of Fundy (1970-2020)

Daniel Ricard, Catalina Gomez, Jamie Emberley and Catriona Regnier-McKellar

Science Branch
Gulf Region
Fisheries and Oceans Canada
Moncton, New Brunswick, E1C 5K4, Canada

Science Branch
Maritimes Region
Fisheries and Oceans Canada
Dartmouth, Nova Scotia, B2Y 4A2, Canada

2021

Canadian Technical Report of
Fisheries and Aquatic Sciences ####



Fisheries and Oceans
Canada Pêches et Océans
Canada

Canada

Canadian Technical Report of Fisheries and Aquatic Sciences

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Rapport technique canadien des sciences halieutiques et aquatiques

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Canadian Technical Report of
Fisheries and Aquatic Sciences nnn

2021

5

MARINE FISH AND INVERTEBRATE ATLAS: GEOGRAPHIC DISTRIBUTION, POPULATION INDICES AND ENVIRONMENTAL PREFERENCES OF MARINE SPECIES IN THE SCOTIAN SHELF AND BAY OF FUNDY DERIVED FROM THE ANNUAL MARITIMES SUMMER SURVEY (1970-2020)

by

10

Daniel Ricard¹ Catalina Gomez² Jamie Emberley² Catriona Regnier-McKellar²

¹Science Branch

Gulf Region

Fisheries and Oceans Canada

Moncton, New Brunswick, E1C 5K4, Canada

^2Science Branch

Maritimes Region

Fisheries and Oceans Canada

Dartmouth, Nova Scotia, B2Y 4A2, Canada

© Her Majesty the Queen in Right of Canada, 2021
Cat. No. Fs97-6/nnnE-PDF ISBN ISSN 1488-5379

22 Correct citation for this publication:

23 Ricard, D., Gomez, C., Emberley, J. and Regnier-McKellar, C. 2021. Marine Fish and
24 Invertebrate Atlas: Geographic Distribution, Population Indices and Environmental Preferences
25 of marine species in the Scotian Shelf and Bay of Fundy derived from the annual Maritimes
26 Summer Survey (1970-2020). Can. Tech. Rep. Fish. Aquat. Sci. nnn: viii + 192 p.

27 CONTENTS

28	ABSTRACT	vii
29	RÉSUMÉ	viii
30	1 Introduction	1
31	2 Methods	1
32	2.1 Survey Description	1
33	2.2 Sampling Design	2
34	2.3 Taxonomic Levels	5
35	2.4 Analyses	15
36	2.4.1 Geographic distribution of catches	15
37	2.4.2 Biomass indices	15
38	2.4.3 Distribution indices	15
39	2.4.4 Length frequencies	15
40	2.4.5 Length-weight relationship and condition factor	16
41	2.4.6 Depth, temperature and salinity distribution of catches	16
42	2.4.7 Density-dependent habitat selection	16
43	2.5 Description of Figures	17
44	2.5.1 Type A	17
45	2.5.2 Type B	17
46	2.5.3 Type C.	17
47	2.5.4 Type D.	17
48	2.5.5 Type E.	18
49	2.5.6 Type F.	18
50	3 Results	18
51	3.1 Summary of successful tows by year and stratum	18
52	3.2 Distribution of depth, bottom temperature and bottom salinity from survey tows	23
53	4 Discussion	23
54	4.1 Diversity of approaches used for mapping fish and invertebrates in the Scotian	
55	Shelf bioregion	23
56	4.2 Interpreting spatial results for marine spatial planning purposes	25
57	5 Acknowledgements	26
58	6 References	28
59	7 Appendix	32
60	7.1 Atlantic cod (<i>Morue franche</i>) - species code 10 (category LF)	33
61	7.2 Haddock (<i>Aiglefin</i>) - species code 11 (category LF)	36
62	7.3 White hake (<i>Merluche blanche</i>) - species code 12 (category LF)	39
63	7.4 Red hake (<i>Merluche écureuil</i>) - species code 13 (category LF)	42
64	7.5 Silver hake (<i>Merlu argenté</i>) - species code 14 (category LF)	45
65	7.6 Pollock (<i>Goberge</i>) - species code 16 (category LF)	48
66	7.7 Atlantic redfishes (<i>Sébastes de l'Atlantique</i>) - species code 23 (category LF)	51

67	7.8	Atlantic halibut (Flétan de l'Atlantique) - species code 30 (category LF)	54
68	7.9	American plaice (Plie canadienne) - species code 40 (category LF)	57
69	7.10	Witch flounder (Plie grise) - species code 41 (category LF)	60
70	7.11	Yellowtail flounder (Limande à queue jaune) - species code 42 (category LF)	63
71	7.12	Winter flounder (Limande-plie rouge) - species code 43 (category LF)	66
72	7.13	Atlantic wolffish (Loup atlantique) - species code 50 (category LF)	69
73	7.14	Atlantic herring (Hareng de l'Atlantique) - species code 60 (category LF)	72
74	7.15	Longhorn sculpin (Chabosseau à dix-huit épines) - species code 300 (category LF)	75
75	7.16	Moustache sculpin (Faux-trigle armé) - species code 304 (category LF)	78
76	7.17	Sea raven (Hémithriptère atlantique) - species code 320 (category LF)	81
77	7.18	Alligatorfish (Poisson-alligator atlantique) - species code 340 (category LF)	84
78	7.19	Monkfish (Baudroie d'Amérique) - species code 400 (category LF)	87
79	7.20	Ocean pout (Loquette d'Amérique) - species code 640 (category LF)	90
80	7.21	Thorny skate (Raie épineuse) - species code 201 (category LF)	93
81	7.22	Smooth skate (Raie lisse) - species code 202 (category LF)	96
82	7.23	Winter skate (Raie tachetée) - species code 204 (category LF)	99
83	7.24	Picked dogfish (Aiguillat commun) - species code 220 (category LF)	102
84	7.25	Northern shortfin squid (Encornet rouge nordique) - species code 4511 (category LF)	105
85	7.26	Atlantic hagfish (Myxine du nord) - species code 241 (category LI)	108
86	7.27	Cusk (Brosme) - species code 15 (category LI)	109
87	7.28	Greenland halibut (Flétan noir) - species code 31 (category LI)	110
88	7.29	Gulf Stream flounder (Plie du Gulf Stream) - species code 44 (category LI)	111
89	7.30	American shad (Alose savoureuse) - species code 61 (category LI)	112
90	7.31	Alewife (Gaspareau) - species code 62 (category LI)	113
91	7.32	Capelin (Capelan) - species code 64 (category LI)	114
92	7.33	Atlantic mackerel (Maquereau commun) - species code 70 (category LI)	115
93	7.34	Longfin hake (Merluche à longues nageoires) - species code 112 (category LI)	116
94	7.35	Fourbeard rockling (Motelle à quatre barbillons) - species code 114 (category LI)	117
95	7.36	Blackbelly rosefish (Sébaste chèvre) - species code 123 (category LI)	118
96	7.37	Greater argentine (Grande argentine) - species code 160 (category LI)	119
97	7.38	Arctic hookear sculpin (Hameçon neigeux) - species code 306 (category LI)	120
98	7.39	Atlantic poacher (Agone atlantique) - species code 350 (category LI)	121
99	7.40	Marlin-spike grenadier (Grenadier du Grand Banc) - species code 410 (category LI)	122
100	7.41	Lumpfish (Lompe) - species code 501 (category LI)	123
101	7.42	Atlantic spiny lumpucker (Petite poule de mer atlantique) - species code 502 (category LI)	124
102	7.43	Sand lance (Lançon) - species code 610 (category LI)	125
103	7.44	Snakeblenny (Lompénie-serpent) - species code 622 (category LI)	126
104	7.45	Daubed shanny (Lompénie tachetée) - species code 623 (category LI)	127
105	7.46	Vahl's eelpout (Lycode à carreaux) - species code 647 (category LI)	128
106	7.47	Atlantic butterfish (Stromaté fossette) - species code 701 (category LI)	129
107	7.48	Atlantic hookear sculpin (Hameçon atlantique) - species code 880 (category LI)	130
108	7.49	Barndoor skate (Grande raie) - species code 200 (category LI)	131
109	7.50	Little skate (Raie hérisson) - species code 203 (category LI)	132
110	7.51	Northern prawn (Crevette nordique) - species code 2211 (category SF)	133
111	7.52	Jonah crab (Tourteau jona) - species code 2511 (category SF)	134

114	7.53	Atlantic rock crab (Tourteau poïnclos) - species code 2513 (category SF)	135
115	7.54	Arctic lyre crab (Crabe Hyas coarctatus) - species code 2521 (category SF)	136
116	7.55	Atlantic king crab (Crabe épineux du nord) - species code 2523 (category SF)	137
117	7.56	Queen crab (Crabe des neiges) - species code 2526 (category SF)	138
118	7.57	Great spider crab (Crabe lyre araignée) - species code 2527 (category SF)	139
119	7.58	American lobster (Homard américain) - species code 2550 (category SF)	140
120	7.59	Sea lamprey (Lamproie marine) - species code 240 (category LR)	141
121	7.60	Atlantic tomcod (Poulamon atlantique) - species code 17 (category LR)	142
122	7.61	Offshore silver hake (Merlu argenté du large) - species code 19 (category LR)	143
123	7.62	Spotted wolffish (Loup tacheté) - species code 51 (category LR)	144
124	7.63	Northern wolffish (Loup à tête large) - species code 52 (category LR)	145
125	7.64	Rainbow smelt (Éperlan arc-en-ciel) - species code 63 (category LR)	146
126	7.65	Cunner (Tanche-tautogue) - species code 122 (category LR)	147
127	7.66	Fourspot flounder (Cardeau à quatre ocelles) - species code 142 (category LR)	148
128	7.67	Windowpane flounder (Turbot de sable) - species code 143 (category LR)	149
129	7.68	Longnose greeneye (Oeil-vert à long nez) - species code 149 (category LR)	150
130	7.69	Lanternfishes (Poissons-lanternes) - species code 150 (category LR)	151
131	7.70	Shortnose greeneye (Éperlan du large) - species code 156 (category LR)	152
132	7.71	Silvery lightfish (Brossé améthyste) - species code 158 (category LR)	153
133	7.72	Boa dragonfish (Dragon-boa) - species code 159 (category LR)	154
134	7.73	Shorthorn sculpin (Chabosseau à épines courtes) - species code 301 (category LR)	155
135	7.74	Grubby (Chabosseau bronzé) - species code 303 (category LR)	156
136	7.75	Polar sculpin (Cotte polaire) - species code 307 (category LR)	157
137	7.76	Spatulate sculpin (Icèle spatulée) - species code 314 (category LR)	158
138	7.77	Arctic alligatorfish (Poisson-alligator arctique) - species code 341 (category LR)	159
139	7.78	Alligatorfishes (Poissons-alligator) - species code 351 (category LR)	160
140	7.79	Roughnose grenadier (Grenadier-scie) - species code 412 (category LR)	161
141	7.80	Roundnose grenadier (Grenadier de roche) - species code 414 (category LR)	162
142	7.81	Atlantic seasnail (Limace atlantique) - species code 503 (category LR)	163
143	7.82	Gelatinous snailfish (Limace gélantineuse) - species code 505 (category LR)	164
144	7.83	Variegated snailfish (Limace marbée) - species code 512 (category LR)	165
145	7.84	Sea tadpole (Petite limace de mer) - species code 520 (category LR)	166
146	7.85	Wolf eelpout (Lycode à tête longue) - species code 603 (category LR)	167
147	7.86	Slender snipe eel (Avocette ruban) - species code 604 (category LR)	168
148	7.87	Newfoundland eelpout (Lycode du Labrador) - species code 619 (category LR)	169
149	7.88	Newfoundland eelpout (Lycode du Labrador) - species code 620 (category LR)	170
150	7.89	Rock gunnel (Sigouine de roche) - species code 621 (category LR)	171
151	7.90	Radiated shanny (Ulvaire deux-lignes) - species code 625 (category LR)	172
152	7.91	Fourline snakeblenny (Quatre-lignes atlantique) - species code 626 (category LR)	173
153	7.92	Wrymouth (Terrassier tacheté) - species code 630 (category LR)	174
154	7.93	Spotfin dragonet (Dragonnet tacheté) - species code 637 (category LR)	175
155	7.94	Arctic eelpout (Lycode arctique) - species code 641 (category LR)	176
156	7.95	Atlantic soft pout (Molasse atlantique) - species code 646 (category LR)	177
157	7.96	Silvery John dory (Saint Pierre argenté) - species code 704 (category LR)	178
158	7.97	White barracudina (Lussion blanc) - species code 712 (category LR)	179
159	7.98	Atlantic saury (Balaou atlantique) - species code 720 (category LR)	180
160	7.99	Hatchetfishes (Haches d'argent) - species code 741 (category LR)	181

161	7.100 Atlantic batfish (Malthe atlantique) - species code 742 (category LR)	182
162	7.101 Spottedfin tonguefish (Langue fil noir) - species code 816 (category LR)	183
163	7.102 Black dogfish (Aiguillat noir) - species code 221 (category LR)	184
164	7.103 Longfin inshore squid (Calmar totam) - species code 4512 (category LR)	185
165	7.104 Red deepsea crab (Crabe rouge) - species code 2532 (category SR)	186
166	INDEX	187

ABSTRACT

168 Ricard, D., Gomez, C., Emberley, J. and Regnier-McKellar, C. 2021. Marine Fish and
169 Invertebrate Atlas: Geographic Distribution, Population Indices and Environmental Preferences
170 of marine species in the Scotian Shelf and Bay of Fundy derived from the annual Maritimes
171 Summer Survey (1970-2020). Can. Tech. Rep. Fish. Aquat. Sci. nnn: viii + 192 p.

172 The summer groundfish research vessel survey on the Scotian Shelf and in the Bay of
173 Fundy started in 1970 and was designed to measure the distribution and abundance of
174 major commercial fish species. Over time, additional information on non-commercial species
175 was collected, and allowed considerable insight into ecosystem function and structure, as
176 documented in many primary publications whose analyses used the survey data. The same
177 groundfish survey database has also been used to produce species status reports, atlases of
178 species distribution and remains an essential source of information for stock assessments in the
179 Maritimes Region of Fisheries and Oceans Canada. This report builds on previous work and
180 former atlases by updating a comprehensive suite of indices to assess population status and
181 environmental preferences of 104 species. For each species, trends in geographic distribution
182 and biomass or abundance were plotted. The spatial extent of distribution was plotted over
183 time to gauge how the area occupied has changed. The relationship between abundance or
184 biomass and spatial extent reflected whether the species distribution expands when abundance
185 or biomass increases. Length frequencies over time depicted any changes in mean size. The
186 plots of condition over time revealed whether individual fish are fatter or thinner than their long
187 term mean. Depth, temperature and salinity preferences were estimated to gauge the range
188 of suitable environmental parameters for each species. Finally, for each stratum, the slope
189 describing how local density varies with regional abundance was estimated. The reproducible
190 set of tools provided in this report constitutes a stepping stone to conduct other ecological
191 analyses using the summer groundfish research vessel survey data by fostering reproducibility
192 and transparency of ecological information collected and reported annually. Recognizing the
193 diversity of approaches for visualizing and mapping fish and invertebrates in the Scotian Shelf
194 bioregion, we recommend the development of a regional community of practice to compare and
195 evaluate approaches for mapping, interpolating and/or modelling fish and invertebrates so future
196 publications and advice can lead to more comparable work and consistent science advice to
197 support processes such as marine spatial planning.

RÉSUMÉ

199 Ricard, D., Gomez, C., Emberley, J. and Regnier-McKellar, C. 2021. Marine Fish and
200 Invertebrate Atlas: Geographic Distribution, Population Indices and Environmental Preferences
201 of marine species in the Scotian Shelf and Bay of Fundy derived from the annual Maritimes
202 Summer Survey (1970-2020). Can. Tech. Rep. Fish. Aquat. Sci. nnn: viii + 192 p.

203 Le relevé estival par navires de recherche sur le poisson de fond sur le plateau néo-écossais
204 et dans la baie de Fundy a débuté en 1970 et visait à mesurer la répartition et l'abondance
205 des principales espèces de poissons commerciales. Au fil du temps, des informations
206 supplémentaires sur les espèces non commerciales ont été recueillies et ont permis de mieux
207 comprendre la fonction et la structure de l'écosystème, comme le montrent de nombreuses
208 publications primaires dont les analyses ont utilisé les données d'enquête. La même base
209 de données sur les relevés du poisson de fond a également été utilisée pour produire des
210 rapports sur la situation des espèces, des atlas de la répartition des espèces et demeure une
211 source essentielle d'information pour les évaluations des stocks dans la région des Maritimes
212 de Pêches et Océans Canada. Ce rapport s'appuie sur des travaux antérieurs et d'anciens
213 atlas en mettant à jour une série complète d'indices pour évaluer l'état de la population et les
214 préférences environnementales de 104 espèces. Pour chaque espèce, les tendances de la
215 répartition géographique et de la biomasse ou de l'abondance ont été tracées. L'étendue spatiale
216 de la distribution a été tracée au fil du temps pour évaluer comment la zone occupée a changé.
217 La relation entre l'abondance ou la biomasse et l'étendue spatiale indique si la répartition
218 des espèces augmente lorsque l'abondance ou la biomasse augmente. Les fréquences de
219 longueur au fil du temps représentaient tout changement dans la taille moyenne. Les graphiques
220 de l'état au fil du temps ont révélé si les poissons individuels sont plus gros ou plus minces
221 que leur moyenne à long terme. Les préférences en matière de profondeur, de température
222 et de salinité ont été estimées pour évaluer la gamme de paramètres environnementaux
223 appropriés pour chaque espèce. Enfin, pour chaque strate, la pente décrivant comment la
224 densité locale varie avec l'abondance régionale a été estimée. L'ensemble d'outils reproductibles
225 fournis dans ce rapport constitue un tremplin pour effectuer d'autres analyses écologiques
226 à l'aide des données du relevé estival des navires de recherche sur les poissons de fond
227 en favorisant la reproductibilité et la transparence de l'information écologique recueillie et
228 rapportée annuellement. Reconnaissant la diversité des approches de visualisation et de
229 cartographie des poissons et des invertébrés dans la biorégion du plateau néo-écossais, nous
230 recommandons le développement d'une communauté de pratique régionale pour comparer et
231 évaluer les approches de cartographie, d'interpolation et / ou de modélisation des poissons
232 et des invertébrés afin conduire à des travaux plus comparables et à des avis scientifiques
233 cohérents pour soutenir des processus tels que la planification de l'espace marin.

234

1 Introduction

235 The summer (July-August) groundfish research vessel survey on the Scotian Shelf and in the
236 Bay of Fundy was started in 1970 by Fisheries and Oceans Canada Maritimes Region. The
237 survey was originally designed to measure the distribution and abundance of major commercial
238 fish species. Over time, information on non-commercial species was also collected. The
239 groundfish survey database storing the information collected during the annual survey provides
240 the main source of fisheries-independent information for marine species in the region. This
241 information is routinely used to support stock assessments, to produce species status reports
242 and has been previously used to publish atlases of species distribution.

243 The current document is an update of an earlier report (Ricard and Shackell 2013) that built
244 on former atlases by updating a comprehensive suite of derived indices for 104 species to
245 assess population status and, when feasible, environmental preferences. The information
246 collected during the survey is stored in a relational database management system archived
247 at Fisheries and Oceans Canada Maritimes Region which contains detailed information about
248 the sampling locations and the associated catch. Tow-level survey data is also publicly available
249 from the Ocean Biogeographic Information System (DFO 2016) and from the Open data portal
250 supported by the federal government (DFO 2021). The present atlas builds upon the work done
251 by Fisheries and Oceans colleagues from the northern Gulf of St. Lawrence (Bourdages and
252 Ouellet 2012), southern Gulf of St. Lawrence (Benoît et al. 2003) and on earlier work in the
253 Scotian Shelf (Simon and Comeau 1994; Horsman and Shackell 2009).

254 To facilitate updates and foster collaboration on the analyses of the survey data, the computer
255 code necessary to extract the data, to perform the analyses presented herein, and to reproduce
256 and update the current document is made available in a git repository (Ricard and Gomez 2021).

257 The survey area covers three major Northwest Atlantic Fisheries Organization (NAFO) zones
258 that divide the Scotian Shelf into the colder east 4V and 4W (strata 440-466) and warmer
259 west 4X (strata 470-495). For each species, temporal trends in geographic distribution and,
260 when possible, biomass are plotted. Some caution is required in interpreting the results
261 obtained for several taxa due to low sample size as explained later in the text. A full ecological
262 interpretation of trends is beyond the scope of this report. Other documents stemming from peer-
263 reviewed scientific processes under the auspices of the [Canadian Science Advisory Secretariat](#)
264 (CSAS) provide further descriptions of spatio-temporal trends in different indicators and put the
265 information collected during the summer groundfish research vessel survey in a more focused
266 context (see for example Clark and Emberley (2011)).

267

2 Methods

268 2.1 Survey Description

269 The survey is conducted annually in July-August and covers the Scotian Shelf and the Bay of
270 Fundy (Figure 2). It normally involves two separate two-week trips on board an offshore fisheries
271 vessel from the Canadian Coast Guard.

272 A number of changes in fishing gear type and vessels used occurred since the onset of sampling
273 activities (Clark and Emberley 2011). Comparative fishing experiments were conducted when
274 those changes in survey platforms took place. A timeline of the survey platforms can be found in
275 Figure 1.




Figure 1. Timeline of vessels and gears used in the Maritimes Region summer survey. The y axis identifies the vessel used and the gear used is overlaid on the polygon representing the time window where each vessel was used.

276 2.2 Sampling Design

277 The summer survey covers divisions 4V, 4W and 4X of the Northwest Atlantic Fisheries
278 Organization (NAFO) which includes the Scotian Shelf and the Bay of Fundy. The eastern limit of
279 the survey is the Laurentian Channel and the western limit is the Fundian Channel (Figure 2).

280 The survey follows a stratified random design (Doubleday and Rivard 1981; Lohr 1999)
281 (Figure 3). The number of tows conducted in each stratum is approximately proportional to the
282 surface area of the stratum. The targeted area covered by the survey has remained constant
283 since its inception, with the exception of additional deeper strata that were only sampled a few
284 times since 2000. Because the sampling of the deeper strata is opportunistic and irregular, the
285 analyses presented herein only include strata 440 to 495 which cover NAFO Divisions 4V, 4W
286 and 4X (Figure 3 and Table 1).

287 The basic sampling unit of the survey is a 30-minute fishing tow conducted at a speed of 3.5
288 knots. This yields a distance towed of 1.75 nautical miles.




Figure 2. Map of the Scotian Shelf and Bay of Fundy where the DFO Maritimes summer survey takes place. The bathymetry presented here is the 15 arc-second gridded data set from the General Bathymetric Chart of the Oceans ([GEBCO](#)). Geographical locations of interest and the boundaries of relevant NAFO Divisions are also shown on the map.

Table 1. Summer survey strata details. The strata used in the analyses are presented separately for NAFO Divisions 4Vn, 4VsW and 4X. For each stratum, the depth range in fathoms and the surface area in square kilometers are reported.

NAFO Div.	Stratum	Depth range (fathom)	Area (km ²)
4Vn	440	101-200	924
	441	51-100	1000
	442	11-49	1437

NAFO Div.	Stratum	Depth range (fathom)	Area (km²)
4VsW	443	11-49	1318
	444	51-100	3925
	445	101-200	1023
	446	101-200	491
	447	11-49	1616
	448	11-49	1449
	449	51-100	144
	450	51-100	383
	451	101-200	147
	452	101-200	345
	453	101-200	259
	454	51-100	499
	455	11-49	2122
	456	11-49	955
	457	51-100	811
	458	11-49	658
	459	11-200	3148
	460	51-100	1344
	461	101-200	1154
	462	51-100	2116
	463	11-49	302
	464	11-50	1297
	465	51-100	2383
	466	101-200	226

NAFO Div.	Stratum	Depth range (fathom)	Area (km ²)
4X	470	51-100	920
	471	101-200	1004
	472	51-100	1249
	473	11-49	265
	474	11-49	161
	475	11-49	156
	476	51-100	1478
	477	51-100	1232
	478	101-200	233
	480	11-49	655
	481	51-100	1875
	482	101-200	1042
	483	101-200	532
	484	101-200	2264
	485	51-100	1582
	490	11-49	601
	491	51-100	687
	492	51-100	1086
	493	11-49	533
	494	11-49	417
	495	11-49	584

289 After each tow the catch is sorted by species and weighed. Each fish caught is then measured,
 290 and further sampling of individual fish weight, maturity status and age are performed for different
 291 length classes. When catches exceed 300 individuals, a random sub-sample is used to obtain
 292 the length and weight measurements.

293 2.3 Taxonomic Levels

294 Fish species caught during the surveys are identified by trained scientific personnel and their
 295 scientific name is determined. An internal species code used in the relational database is
 296 reported for each species (Losier and Waite 1989).

297 By its nature as a bottom trawl, the fishing gear used in the survey catches certain species
 298 better than others. To ensure that meaningful ecological information can be extracted from
 299 catch samples, we report the catch records for the subset of species that are caught reliably
 300 by the gear. To appear in this atlas, a species must have had a minimum of 10 observations over
 301 the duration of the survey activities. While both catch abundance and weight are recorded, the
 302 weight of species that appear at low abundances is often recorded as zero in the earlier parts of
 303 the survey when scales of appropriate precision were not available.

304 We divided the species caught into five categories based on 1) their taxonomic classification,




Figure 3. Map of the summer survey strata 440 to 495.

305 2) the number of recorded observations, and 3) their period of valid identification (Table 2).
 306 Category "LF", for "long frequent", was assigned to species that have more than 1000 records
 307 since 1970 and have been consistently identified since the onset of the survey. Category
 308 "LI", for "long intermediate", was assigned to species that had between 1000 and 200 catch
 309 records. Rare and elusive species (those with less than 200 catch records over the duration
 310 of the survey) are also reported but to a lower level of analytical details (Category "LR", for
 311 "long rare"). Category "SF", for "short frequent", was assigned to invertebrate species that were
 312 consistently sampled only since 1999 (Tremblay M. J. 2007). And category "SR", for "short rare"
 313 for invertebrate species consistently sampled only since 1999 and with less than 200 catch
 314 records. To ensure concordance with authoritative taxonomic information, the AphiaID from the

Table 2. Taxonomic levels used to determine the analytical treatment for each species.

Category	Name	Description
L	long - consistently identified since the onset of the survey in 1970	
LF	long frequent	species that have more than 1000 catch records
LI	long intermediate	species that had between 1000 and 200 catch records
LR	long rare	species with less than 200 catch records
S	short - invertebrate species that were consistently sampled only since 1999	
SF	short frequent	species with more than 200 catch records
SR	short rare	species with less than 200 catch records

³¹⁵ World Register of Marine Species (Appeltans et al. 2012) is included for the different species
³¹⁶ presented in this document (Table 3) .

Table 3. List of species included in the Atlas. For each taxonomic order and class, each species is listed in the table, its taxonomic family and scientific name is provided, along with its French and English common names, the species code used in the survey database, its AphiaID with a link to the World Registry of Marine Species, its number of catch records in the survey database and its classification category as defined in section 2.3.

	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Actinopterygii								
<i>Anguilliformes</i>								
	Nemichthyidae	<i>Nemichthys scolopaceus</i>	Slender snipe eel	Avocette ruban	604	126306	28	LR
<i>Argentiniformes</i>								
	Argentinidae	<i>Argentina silus</i>	Greater argentine	Grande argentine	160	126715	963	LI
<i>Aulopiformes</i>								
	Chlorophthalmidae	<i>Chlorophthalmus agassizi</i>	Shortnose greeneye	Éperlan du large	156	126336	78	LR
		<i>Parasudis truculenta</i>	Longnose greeneye	Oeil-vert à long nez	149	158868	45	LR
	Paralepididae	<i>Arctozenus risso</i>	White barracudina	Lussion blanc	712	126352	196	LR
<i>Beloniformes</i>								
	Scomberesocidae	<i>Scomberesox saurus</i>	Atlantic saury	Balaou atlantique	720	126392	37	LR
<i>Clupeiformes</i>								
	Clupeidae	<i>Alosa pseudoharengus</i>	Alewife	Gaspareau	62	158669	977	LI
		<i>Alosa sapidissima</i>	American shad	Alose savoureuse	61	158670	468	LI
		<i>Clupea harengus</i>	Atlantic herring	Hareng de l'Atlantique	60	126417	3487	LF
<i>Gadiformes</i>								
	Gadidae	<i>Gadus morhua</i>	Atlantic cod	Morue franche	10	126436	5451	LF
		<i>Melanogrammus aeglefinus</i>	Haddock	Aiglefin	11	126437	5827	LF
		<i>Microgadus tomcod</i>	Atlantic tomcod	Poulamon atlantique	17	158928	44	LR
		<i>Pollachius virens</i>	Pollock	Goberge	16	126441	2787	LF
	Lotidae	<i>Brosme brosme</i>	Cusk	Brosme	15	126447	688	LI

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Perciformes	Macrouridae		<i>Enchelyopus cimbricus</i>	Fourbeard rockling	Motelle à quatre barbillons	114	126450	693	LI
			<i>Coryphaenoides rupestris</i>	Roundnose grenadier	Grenadier de roche	414	158960	17	LR
			<i>Nezumia bairdii</i>	Marlin-spike grenadier	Grenadier du Grand Banc	410	183289	529	LI
			<i>Trachyrincus murrayi</i>	Roughnose grenadier	Grenadier-scie	412	126481	18	LR
	Merlucciidae		<i>Merluccius albidus</i>	Offshore silver hake	Merlu argenté du large	19	158748	161	LR
			<i>Merluccius bilinearis</i>	Silver hake	Merlu argenté	14	158962	4936	LF
	Phycidae		<i>Phycis chesteri</i>	Longfin hake	Merluche à longues nageoires	112	158988	784	LI
			<i>Urophycis chuss</i>	Red hake	Merluche écureuil	13	126503	2195	LF
			<i>Urophycis tenuis</i>	White hake	Merluche blanche	12	126504	3524	LF
<i>Lophiiformes</i>	Lophiidae	<i>Lophius americanus</i>	Monkfish	Baudroie d'Amérique	400	159184	1970	LF	
	Ogcocephalidae	<i>Dibranchus atlanticus</i>	Atlantic batfish	Malthe atlantique	742	126558	18	LR	
	Myctophidae	<i>Myctophidae</i>	Lanternfishes	Poissons-lanternes	150	125498	160	LR	
<i>Osmeriformes</i>	Osmeridae		<i>Mallotus villosus</i>	Capelin	Capelan	64	126735	540	LI
			<i>Osmerus mordax</i>	Rainbow smelt	Éperlan arc-en-ciel	63	126737	59	LR
	Ammodytidae	<i>Ammodytes dubius</i>	Sand lance	Lançon	610	151520	1283	LI	
<i>Anarhichadidae</i>			<i>Anarhichas denticulatus</i>	Northern wolffish	Loup à tête large	52	126757	17	LR
			<i>Anarhichas lupus</i>	Atlantic wolffish	Loup atlantique	50	126758	1572	LF

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
			<i>Anarhichas minor</i>	Spotted wolffish	Loup tacheté	51	126759	20	LR
		Callionymidae	<i>Foetorepus agassizii</i>	Spotfin dragonet	Dragonnet tacheté	637	276339	20	LR
		Cryptacanthodidae	<i>Cryptacanthodes maculatus</i>	Wrymouth	Terrassier tacheté	630	159675	120	LR
		Labridae	<i>Tautogolabrus adspersus</i>	Cunner	Tanche-tautogue	122	159785	82	LR
		Pholidae	<i>Pholis gunnellus</i>	Rock gunnel	Sigouine de roche	621	126996	21	LR
		Scombridae	<i>Scomber scombrus</i>	Atlantic mackerel	Maquereau commun	70	127023	696	LI
		Stichaeidae	<i>Eumesogrammus praecisus</i>	Fourline snakeblenny	Quatre-lignes atlantique	626	159817	40	LR
			<i>Leptoclinus maculatus</i>	Daubed shanny	Lompénie tachetée	623	127072	443	LI
			<i>Lumpenus lampretaeformis</i>	Snakeblenny	Lompénie-serpent	622	154675	423	LI
			<i>Ulvaria subbifurcata</i>	Radiated shanny	Ulvaire deux-lignes	625	159821	145	LR
		Stromateidae	<i>Peprilus triacanthus</i>	Atlantic butterfish	Stromaté fossette	701	159828	487	LI
		Zoarcidae	<i>Lycenchelys verrillii</i>	Wolf eelpout	Lycode à tête longue	603	159258	40	LR
			<i>Lycodes lavalaei</i>	Newfoundland eelpout	Lycode du Labrador	620	127107	72	LR
			<i>Lycodes reticulatus</i>	Arctic eelpout	Lycode arctique	641	127112	70	LR
			<i>Lycodes terraenovae</i>	Newfoundland eelpout	Lycode du Labrador	619	127117	64	LR
			<i>Lycodes vahlii</i>	Vahl's eelpout	Lycode à carreaux	647	127118	565	LI
			<i>Melanostigma atlanticum</i>	Atlantic soft pout	Molasse atlantique	646	127120	43	LR
			<i>Zoarces americanus</i>	Ocean pout	Loquette d'Amérique	640	159267	1478	LF
<i>Pleuronectiformes</i>		Cynoglossidae	<i>Syphurus diomedeanus</i>	Spottedfin tonguefish	Langue fil noir	816	159358	24	LR

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
		Paralichthyidae	<i>Citharichthys arctifrons</i>	Gulf Stream flounder	Plie du Gulf Stream	44	158791	382	LI
			<i>Hippoglossina oblonga</i>	Fourspot flounder	Cardeau à quatre ocelles	142	158833	76	LR
		Pleuronectidae	<i>Glyptocephalus cynoglossus</i>	Witch flounder	Plie grise	41	127136	4301	LF
			<i>Hippoglossoides platessoides</i>	American plaice	Plie canadienne	40	127137	6023	LF
			<i>Hippoglossus hippoglossus</i>	Atlantic halibut	Flétan de l'Atlantique	30	127138	1634	LF
			<i>Limanda ferruginea</i>	Yellowtail flounder	Limande à queue jaune	42	158879	3233	LF
			<i>Pseudopleuronectes americanus</i>	Winter flounder	Limande-plie rouge	43	158885	1632	LF
			<i>Reinhardtius hippoglossoides</i>	Greenland halibut	Flétan noir	31	127144	736	LI
		Scophthalmidae	<i>Scophthalmus aquosus</i>	Windowpane flounder	Turbot de sable	143	158907	115	LR
Scorpaeniformes	Agonidae		<i>Agonidae</i>	Alligatorfishes	Poissons-alligator	351	125588	43	LR
			<i>Aspidophoroides monopterygius</i>	Alligatorfish	Poisson-alligator atlantique	340	159459	1029	LF
			<i>Leptagonus decagonus</i>	Atlantic poacher	Agone atlantique	350	127191	266	LI
			<i>Ulcina olrikii</i>	Arctic alligatorfish	Poisson-alligator arctique	341	274356	13	LR
	Cottidae		<i>Artediellus atlanticus</i>	Atlantic hookear sculpin	Hameçon atlantique	880	127193	258	LI
			<i>Artediellus uncinatus</i>	Arctic hookear sculpin	Hameçon neigeux	306	127195	306	LI

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Perciformes	Triglidae		<i>Icelus spatula</i>	Spatulate sculpin	lcèle spatulée	314	127200	40	LR
			<i>Myoxocephalus aenaeus</i>	Grubby	Chabosseau bronzé	303	159519	40	LR
			<i>Myoxocephalus octodecemspinosus</i>	Longhorn sculpin	Chabosseau à dix-huit épines	300	159520	3292	LF
			<i>Myoxocephalus scorpius</i>	Shorthorn sculpin	Chabosseau à épines courtes	301	127203	131	LR
			<i>Triglops murrayi</i>	Moustache sculpin	Faux-trigle armé	304	127205	1182	LF
	Cyclopteridae		<i>Cyclopterus lumpus</i>	Lumpfish	Lompe	501	127214	216	LI
			<i>Eumicrotremus spinosus</i>	Atlantic spiny lumpucker	Petite poule de mer atlantique	502	127217	226	LI
	Hemitripteridae		<i>Hemitripterus americanus</i>	Sea raven	Hémithriptère atlantique	320	159518	2126	LF
	Liparidae		<i>Careproctus reinhardtii</i>	Sea tadpole	Petite limace de mer	520	127212	18	LR
			<i>Liparis atlanticus</i>	Atlantic seasnail	Limace atlantique	503	159524	34	LR
			<i>Liparis fabricii</i>	Gelatinous snailfish	Limace gélatineuse	505	127218	27	LR
			<i>Liparis gibbus</i>	Variegated snailfish	Limace marbée	512	159526	41	LR
Stomiiformes	Psychrolutidae		<i>Cottunculus microps</i>	Polar sculpin	Cotte polaire	307	127235	29	LR
	Sebastidae		<i>Helicolenus dactylopterus</i>	Blackbelly rosefish	Sébaste chèvre	123	127251	610	LI
			<i>Sebastes</i>	Atlantic redfishes	Sébastes de l'Atlantique	23	126175	4152	LF
	Sternopychidae		<i>Maurolicus muelleri</i>	Silvery lightfish	Brossé améthyste	158	127312	52	LR
			<i>Sternopychidae</i>	Hatchetfishes	Haches d'argent	741	125603	21	LR
	Stomiidae		<i>Stomias boa</i>	Boa dragonfish	Dragon-boa	159	127374	20	LR
<i>Zeiformes</i>									

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
		Zeidae	<i>Zenopsis conchifer</i>	Silvery John dory	Saint Pierre argenté	704	127426	39	LR
Cephalopoda									
<i>Myopsida</i>									
		Loliginidae	<i>Doryteuthis pealeii</i>	Longfin inshore squid	Calmar totam	4512	574541	96	LR
<i>Oegopsida</i>									
		Ommastrephidae	<i>Illex illecebrosus</i>	Northern shortfin squid	Encornet rouge nordique	4511	153087	4836	LF
Elasmobranchii									
<i>Rajiformes</i>									
		Rajidae	<i>Amblyraja radiata</i>	Thorny skate	Raie épineuse	201	105865	3937	LF
			<i>Dipturus laevis</i>	Barndoor skate	Grande raie	200	158548	246	LI
			<i>Leucoraja erinacea</i>	Little skate	Raie hérisson	203	158551	712	LI
			<i>Leucoraja ocellata</i>	Winter skate	Raie tachetée	204	158553	1180	LF
			<i>Malacoraja senta</i>	Smooth skate	Raie lisse	202	158554	1773	LF
<i>Squaliformes</i>									
		Etmopteridae	<i>Centroscyllium fabricii</i>	Black dogfish	Aiguillat noir	221	105906	31	LR
		Squalidae	<i>Squalus acanthias</i>	Piked dogfish	Aiguillat commun	220	105923	1985	LF
Malacostraca									
<i>Decapoda</i>									
		Cancridae	<i>Cancer borealis</i>	Jonah crab	Tourteau jona	2511	158056	1387	SF
			<i>Cancer irroratus</i>	Atlantic rock crab	Tourteau poïnclos	2513	158057	788	SF
		Geryonidae	<i>Chaceon quinquedens</i>	Red deepsea crab	Crabe rouge	2532	158407	33	SR
		Lithodidae	<i>Lithodes maja</i>	Atlantic king crab	Crabe épineux du nord	2523	107205	531	SF
		Nephropidae	<i>Homarus americanus</i>	American lobster	Homard américain	2550	156134	1623	SF

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Oregoniidae			<i>Chionoecetes opilio</i>	Queen crab	Crabe des neiges	2526	107315	1546	SF
			<i>Hyas araneus</i>	Great spider crab	Crabe lyre araignée	2527	107322	625	SF
			<i>Hyas coarctatus</i>	Arctic lyre crab	Crabe Hyas coarctatus	2521	107323	711	SF
		Pandalidae	<i>Pandalus borealis</i>	Northern prawn	Crevette nordique	2211	107649	718	SF
Myxini									
<i>Myxiniformes</i>									
		Myxinidae	<i>Myxine glutinosa</i>	Atlantic hagfish	Myxine du nord	241	101170	804	LI
Petromyzonti									
<i>Petromyzontiformes</i>									
		Petromyzontidae	<i>Petromyzon marinus</i>	Sea lamprey	Lamproie marine	240	101174	16	LR

317 **2.4 Analyses**

318 The Oracle relational database where all survey data are stored and archived is accessible from
319 the Bedford Institute of Oceanography in Dartmouth, Nova Scotia. Queries written in Structured
320 Query Language (SQL) are used to extract the data from the production server and to create the
321 data products used in all subsequent analyses. Catch records classified as "valid" (i.e. coming
322 from a representative tow without damage to the net) are used in the current analyses. To make
323 the available samples comparable, catch number and weight for each species was standardized
324 for the distance towed.

325 All data processing and analyses were conducted using the R software (R Core Team 2020)
326 using packages gstat (Pebesma 2004), PBSmapping (Schnute et al. 2019), RODBC (Ripley
327 and Lapsley 2019), spatstat (Baddeley 2015), maptools (Bivand and Lewin-Koh 2020), rgeos
328 (Bivand and Rundel 2020), classInt(Bivand 2020), RColorBrewer(Neuwirth 2014), MASS (Ripley
329 et al. 2020), worms (Holstein 2018), and tidyverse (Wickham 2019). The present document is
330 rendered as a Technical Report using the csasdown R package developed and maintained by
331 Fisheries and Oceans Canada scientists (Anderson et al. 2021).

332 **2.4.1 Geographic distribution of catches**

333 Spatial interpolation of catch biomass (kg/tow) was done using a weighting inversely proportional
334 to the distance (inverse-distance weighted, IDW), using function "idw" of the spatstat R package
335 (Baddeley 2015). The IDW method was used with a power parameter value of 10.

336 **2.4.2 Biomass indices**

337 For each species, stratified random estimates of catch biomass (Smith 1996) were computed for
338 each year. Yearly estimates of the standard error were also computed.

339 **2.4.3 Distribution indices**

340 For each Category L, I and S fish species, the minimum area required to account for 75% and
341 95% of the total biomass were computed (D75% and D95%). These measures of distributions
342 were computed for each year by using the Lorenz curve of mean stratum-level catch estimates
343 and the area of occupied strata (Swain and Sinclair 1994; Swain and Morin 1996).

344 **2.4.4 Length frequencies**

345 The length frequency distribution of catch (the stratified numbers-at-length) is tabulated for each
346 seven-year period (1970-2009), and last ten-year period (2010-2020).

347 **2.4.5 Length-weight relationship and condition factor**

348 The relationship between the weight and the length of fish was estimated using the following
349 non-linear isometric relationship:

$$W = \alpha L^\beta$$

350 where W is the total weight (g), L is the length (cm), and, α and β are the parameters to be
351 estimated.

352 Average fish condition (C) was computed as:

$$C = \frac{W}{\alpha L^\beta}$$

353

354 **2.4.6 Depth, temperature and salinity distribution of catches**

355 For each category L species, We followed the methods developed by (Perry and Smith 1994)
356 and generated cumulative frequency distributions of depth, temperature and salinity of survey
357 catches.

358 **2.4.7 Density-dependent habitat selection**

359 We followed the methods of (Myers and Stokes 1989) to evaluate how fish abundance in each
360 stratum varied with overall temporal fluctuations of population abundance.

361 For each category L species, we fitted a model of the relationship between stratum-level density
362 and overall abundance (the yearly stratified random estimate of abundance, defined above).
363 To properly use the observations of zero catch while accounting for the logarithmic distribution
364 of catch abundance, we implemented the model as a generalised linear using a log link and a
365 Poisson error distribution:

$$Y_{h,i} = \alpha_{h,i} Y_i^{\beta_{h,i}}$$

366 where, $y_{h,i}$ is the average abundance of stratum h in year i , and $\alpha_{h,i}$ and $\beta_{h,i}$ are the fitted
367 parameters. The estimated parameter $\beta_{h,i}$ is referred to as the “slope parameter” and indicates
368 whether stratum-level density is positively ($\beta_{h,i} <= 0$), negatively ($\beta_{h,i} >= 0$) or negligibly
369 ($\beta_{h,i} \approx 0$) related to population abundance.

370 To estimate the suitability of each stratum, the median abundance observed during the years
371 that are in the top 25% of yearly estimates is used. We combine the slope parameter estimates
372 from the above model with the median abundance to identify strata that have consistently high
373 abundance and whose local density is weakly related to fluctuation in population abundance
374 ($\beta_{h,i} \approx 0$). Preferred strata are identified for each category L species.

375 **2.5 Description of Figures**

376 **2.5.1 Type A**

377 For Category L and S species:

378 Spatial distribution of catch-per unit of effort, (CPUE, kilograms per tow) in July-August for the
379 Bay of Fundy and Scotian Shelf in five-year periods. Spatial interpolation between tows was
380 done using Inverse Distance Weight (IDW). The probability of occurrence (proportion of tows with
381 catch records for a given species) was also reported for each five-year period.

382 For Category LR and SR:

383 Location of tows with catch over the period 1970-2020 (Type LR) or the period 1999-2020 (Type
384 SR). Location of tows with catch over the period 1970-2020 (Type LR) or the period 1999-2020
385 (Type SR).

386 **2.5.2 Type B**

387 For Category L, S and I species:

388 Stratified random estimate of CPUE (left panel), distribution indices (D75% and D95%, the
389 minimum area containing 75% and 95% of biomass, middle panel), and distribution vs. weight
390 per tow (right panel). The stratified random mean is plotted as a solid line with the 95%
391 confidence region indicated by the solid grey line. The overall mean is plotted as a grey
392 horizontal line and the overall mean plus or minus 50% of the standard deviation appear as
393 horizontal dashed lines. In all three panels, the early years appear in blue and the last years
394 appear in red. The predictions from a loess estimator are overlaid on the distribution indices
395 (middle panel). The Pearson correlation coefficient between D75% and biomass, and its
396 statistical significance, are also reported in the right panel.

397 **2.5.3 Type C.**

398 Length frequency distribution for NAFO divisions 4X and 4VW. A smoothed length frequency
399 distribution is shown for each 7-year periods covered by the surveys.

400 **2.5.4 Type D.**

401 Average fish condition for all fish lengths (black dots and black line), large fish (thick gray line),
402 and small fish (thin gray line). Fish condition is presented for NAFO divisions 4VW (right panel)
403 and 4X (left panel).

404 **2.5.5 Type E.**

405 Cumulative frequency distributions of depth, temperature and salinity at all sampled locations
406 (thick solid line) and at fishing locations with catch records (thin dashed line). The depth,
407 temperature and salinity associated with 5%, 25%, 50%, 75% and 95% of the cumulative catch is
408 shown in tabular fashion on the bottom right panel.

409 **2.5.6 Type F.**

410 Slopes estimates from the density-dependent habitat selection model (y axis) plotted versus
411 the median abundance during the top 25% of years. The red box indicates strata of particular
412 importance for a species by identifying slopes that are within a standard error from zero and that
413 are within the top 25% of median abundance. Each stratum is identified on the plot by the last
414 two digits of its number.

415 **3 Results**

416 The plots generated for each species are presented in the Appendix.

417 **3.1 Summary of successful tows by year and stratum**

418 A total of 9080 representative tows were conducted for the period spanning from 1970 to 2020
419 (Figure 4).

420 Tables 4, 5 and 6 present the number of tows conducted in each stratum and year.




Figure 4. Map of the 9080 representative tows in the Summer survey from 1970 to 2020.

Table 4. Number of representative tows conducted in each stratum during the period 1970 to 1989.

Stratum	NAFO Div.	Area (km ²)	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
440	4VN	3173.016	4	2	2	3	3	3	3	3	3	3	3	3	3	3	3	4	5	5	6	4
441	4VN	3434.000	4	2	2	3	3	3	1	3	3	3	3	3	3	3	3	5	5	4	4	4
442	4VN	4934.658	3	2	2	2	3	3	2	3	3	3	3	3	3	3	3	3	5	6	7	5
443	4VSW	4526.012	4	2	4	4	8	3	1	2	4	4	4	3	3	5	4	4	6	6	5	2
444	4VSW	13478.450	3	2	5	4	6	4	6	7	4	4	4	5	5	6	4	4	6	6	3	6
445	4VSW	3512.982	5	2	5	4	5	5	1	3	4	4	4	5	5	3	4	5	6	4	4	4
446	4VSW	1686.094	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	4	3	3	3	3
447	4VSW	5549.344	4	2	6	5	7	4	4	3	4	4	4	5	4	4	4	4	5	7	6	6
448	4VSW	4975.866	5	2	5	4	5	4	4	4	4	4	4	4	6	4	4	4	5	5	5	5
449	4VSW	494.496	2	2	2	2	3	2	2	2	1	2	2	2	1	2	2	2	2	2	2	2
450	4VSW	1315.222	2	2	3	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
451	4VSW	504.798	1	2	2	2	2	2	2	2	2	3	2	2	3	2	2	2	2	2	2	2
452	4VSW	1184.730	2	2	2	2	2	2	2	2	2	2	2	4	2	2	2	2	2	2	2	2
453	4VSW	889.406	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	2	2	2
454	4VSW	1713.566	3	2	3	3	3	3	3	3	3	3	3	2	3	3	3	3	3	2	2	2
455	4VSW	7286.948	7	6	7	6	7	6	6	7	7	7	7	7	7	7	7	8	7	7	7	7
456	4VSW	3279.470	5	4	6	5	5	6	4	6	6	6	7	6	6	6	6	6	7	6	6	6
457	4VSW	2784.974	2	2	2	2	3	2	2	2	2	2	2	3	2	2	2	2	2	4	2	2
458	4VSW	2259.572	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	5	5	3	3
459	4VSW	10810.232	3	2	4	4	4	4	4	4	4	4	4	4	3	4	4	6	6	5	6	5
460	4VSW	4615.296	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	4	3	3	3
461	4VSW	3962.836	3	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3	3	2	2
462	4VSW	7266.344	3	3	4	3	4	4	4	4	4	4	4	6	4	4	4	4	6	5	4	4
463	4VSW	1037.068	2	2	2	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2
464	4VSW	4453.898	4	3	5	3	3	6	5	5	5	5	5	4	5	5	5	7	6	5	5	5
465	4VSW	8183.222	6	5	5	4	5	4	5	5	5	5	5	7	6	5	5	5	8	8	8	8
466	4VSW	776.084	2	2	3	2	3	3	3	3	3	3	3	2	3	3	3	3	3	2	2	2
470	4X	3159.280	1	2	2	2	2	3	2	2	2	2	2	2	2	2	2	2	2	3	3	3
471	4X	3447.736	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
472	4X	4289.066	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	4	4	4
473	4X	910.010	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
474	4X	552.874	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	0	2	2	2	2
475	4X	535.704	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
476	4X	5075.452	2	2	2	2	2	2	3	2	2	2	1	2	2	2	2	2	2	4	4	4
477	4X	4230.688	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2	5	4	4
478	4X	800.122	2	2	3	2	3	3	3	3	2	3	3	3	3	3	3	3	3	2	2	2
480	4X	2249.270	4	4	4	3	3	3	4	4	3	4	3	3	4	4	4	4	4	4	4	4
481	4X	6438.750	5	3	4	4	4	3	4	4	5	4	3	4	4	4	4	4	4	6	7	6
482	4X	3578.228	2	1	2	2	2	2	3	2	2	3	2	2	2	2	2	2	2	3	3	3
483	4X	1826.888	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2	2
484	4X	7774.576	2	2	3	3	3	3	3	3	2	3	3	3	4	3	3	3	4	4	4	4
485	4X	5432.588	2	2	2	3	3	3	3	3	3	2	3	4	3	3	3	3	6	7	6	6
490	4X	2063.834	2	2	2	2	2	3	3	3	3	2	3	3	3	3	3	3	3	4	4	4
491	4X	2359.158	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	4	4	4
492	4X	3729.324	3	2	4	3	3	3	3	3	3	3	3	3	3	3	3	3	3	4	4	4
493	4X	1830.322	1	2	3	3	3	3	3	3	3	3	2	3	3	3	3	3	3	3	3	3
494	4X	1431.978	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
495	4X	2005.456	2	2	2	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2
		171809.888	134	110	146	134	153	143	135	144	141	147	145	150	150	146	143	152	171	188	177	170

Table 5. Number of representative tows conducted in each stratum during the period 1990 to 2009.

Stratum	NAFO Div.	Area (km ²)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	
440	4VN	3173.016	4	4	4	3	4	4	4	4	4	4	6	4	4	4	4	4	4	4	3	4	
441	4VN	3434.000	6	5	5	5	5	5	5	5	6	7	6	6	7	6	7	6	6	5	6		
442	4VN	4934.658	5	5	6	5	6	6	6	6	7	6	6	5	6	6	7	5	5	5	6		
443	4VSW	4526.012	4	2	4	3	3	4	4	5	5	4	4	5	5	5	5	4	4	4	5	4	
444	4VSW	13478.450	7	8	8	9	6	8	8	7	8	8	9	10	9	9	9	8	10	8	6	9	
445	4VSW	3512.982	4	4	4	5	7	4	4	4	3	3	6	5	5	5	5	6	5	4	3	6	
446	4VSW	1686.094	3	3	3	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	2	3	
447	4VSW	5549.344	8	7	7	7	7	7	6	7	7	6	7	7	7	7	7	7	6	6	4	6	
448	4VSW	4975.866	9	6	6	7	7	7	6	7	6	7	8	8	8	8	7	8	8	6	5	7	
449	4VSW	494.496	2	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2	
450	4VSW	1315.222	3	3	3	3	3	3	3	3	2	3	3	3	3	3	3	3	3	3	3	3	
451	4VSW	504.798	2	2	2	2	2	2	4	2	2	2	2	2	2	2	2	2	2	2	2	2	
452	4VSW	1184.730	3	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
453	4VSW	889.406	3	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	1	2	
454	4VSW	1713.566	3	2	2	2	2	2	3	2	2	2	2	2	2	2	2	3	2	2	2	2	
455	4VSW	7286.948	12	10	10	9	10	10	10	13	8	11	11	11	11	11	8	12	11	7	5	8	
456	4VSW	3279.470	10	7	7	8	8	8	8	8	6	8	10	8	8	8	8	8	8	6	2	7	
457	4VSW	2784.974	4	2	2	2	2	2	2	2	1	4	2	2	2	2	2	2	2	2	2	2	
458	4VSW	2259.572	9	8	8	8	8	8	7	8	5	6	10	8	7	8	8	10	8	5	2	7	
459	4VSW	10810.232	5	5	6	4	6	6	4	5	6	6	8	6	6	6	6	6	6	5	3	6	
460	4VSW	4615.296	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	4	3	2	3	3	
461	4VSW	3962.836	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	4	2	2	2	2	
462	4VSW	7266.344	5	5	4	4	4	4	4	4	4	4	4	4	4	4	4	5	4	3	4	4	
463	4VSW	1037.068	3	2	2	2	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	
464	4VSW	4453.898	9	7	7	7	7	7	7	4	7	7	7	7	7	7	5	8	7	6	4	5	
465	4VSW	8183.222	12	9	10	10	10	10	10	10	9	10	10	10	10	10	10	10	10	7	8	7	
466	4VSW	776.084	3	2	2	2	2	3	2	2	3	2	2	2	2	2	2	2	2	1	3	2	
470	4X	3159.280	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
471	4X	3447.736	2	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2	
472	4X	4289.066	6	4	4	4	4	4	4	3	4	4	4	4	4	4	4	4	4	3	4	3	
473	4X	910.010	3	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
474	4X	552.874	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
475	4X	535.704	2	2	2	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2	
476	4X	5075.452	4	4	4	4	4	4	4	4	4	4	4	4	4	5	4	4	4	4	4	4	
477	4X	4230.688	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	
478	4X	800.122	2	2	2	2	2	3	3	2	2	2	2	2	2	2	2	2	3	2	2	2	
480	4X	2249.270	8	8	8	8	8	8	8	8	8	8	7	8	8	8	7	9	8	8	8	8	
481	4X	6438.750	8	9	9	9	9	7	9	9	9	9	8	9	8	9	8	9	6	12	9	7	8
482	4X	3578.228	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	2	4	3	3	3	
483	4X	1826.888	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
484	4X	7774.576	3	3	3	3	3	3	3	3	3	3	3	3	3	4	3	3	4	4	3	4	
485	4X	5432.588	2	3	3	3	3	3	3	3	3	3	3	4	3	5	5	3	2	5	4	5	
490	4X	2063.834	4	4	4	4	4	5	4	4	4	3	4	4	4	4	4	4	3	3	3	4	
491	4X	2359.158	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	5	3	3	4	4	
492	4X	3729.324	3	3	3	3	3	2	3	3	3	3	3	3	3	3	5	2	3	4	4	4	
493	4X	1830.322	3	3	3	3	3	2	3	3	2	3	3	3	4	5	2	4	4	3	3	4	
494	4X	1431.978	2	2	2	2	2	2	2	2	2	2	2	2	2	3	4	2	2	4	3	4	
495	4X	2005.456	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	5	3	3	4	
		171809.888	213	189	193	190	195	195	191	193	186	191	213	201	208	216	188	222	209	177	165	196	

Table 6. Number of representative tows conducted in each stratum during the period 2010 to 2020 and for the whole 1970 to 2020 period.

Stratum	NAFO Div.	Area (km2)	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	Total
440	4VN	3173.016	4	5	4	4	4	4	4	4	0	5	4	190
441	4VN	3434.000	6	7	6	6	6	6	6	6	0	7	4	238
442	4VN	4934.658	5	6	6	6	6	6	6	6	0	6	5	240
443	4VSW	4526.012	4	6	5	5	3	7	4	5	0	9	4	214
444	4VSW	13478.450	11	13	9	8	9	9	11	10	0	6	8	352
445	4VSW	3512.982	4	7	2	4	3	4	4	4	0	6	3	215
446	4VSW	1686.094	3	4	3	3	3	2	3	2	0	3	2	145
447	4VSW	5549.344	6	8	6	7	7	7	7	7	0	6	5	291
448	4VSW	4975.866	7	10	8	8	8	7	6	6	0	7	4	299
449	4VSW	494.496	2	4	2	2	2	2	2	2	0	2	2	100
450	4VSW	1315.222	3	3	3	3	3	3	3	2	0	3	2	144
451	4VSW	504.798	2	2	2	2	2	2	2	2	0	2	2	104
452	4VSW	1184.730	2	2	2	2	1	4	3	3	0	3	3	110
453	4VSW	889.406	2	1	3	2	3	2	2	1	0	2	2	116
454	4VSW	1713.566	2	4	2	2	2	2	2	2	0	3	2	121
455	4VSW	7286.948	10	10	10	11	11	9	9	8	0	9	6	429
456	4VSW	3279.470	7	9	8	8	6	5	6	6	0	6	4	331
457	4VSW	2784.974	2	4	2	2	2	3	3	3	0	3	2	113
458	4VSW	2259.572	6	9	8	6	4	5	5	5	0	6	3	269
459	4VSW	10810.232	6	7	6	6	6	7	7	6	0	9	7	262
460	4VSW	4615.296	3	4	4	3	3	5	5	5	3	6	5	151
461	4VSW	3962.836	2	3	3	2	2	3	3	3	2	3	3	113
462	4VSW	7266.344	4	6	4	4	5	5	5	5	0	5	5	212
463	4VSW	1037.068	2	3	2	2	2	3	2	2	0	2	2	107
464	4VSW	4453.898	6	7	7	7	7	6	6	4	0	6	4	288
465	4VSW	8183.222	8	10	10	10	10	10	9	7	3	10	7	397
466	4VSW	776.084	2	2	2	2	2	2	2	3	0	3	2	118
470	4X	3159.280	2	2	3	2	2	3	3	3	4	3	2	112
471	4X	3447.736	2	2	3	2	2	3	3	3	4	4	3	110
472	4X	4289.066	4	6	4	4	4	4	4	4	4	4	4	172
473	4X	910.010	2	2	2	2	2	2	2	2	2	2	2	104
474	4X	552.874	2	2	2	2	2	2	2	2	2	2	2	100
475	4X	535.704	2	2	2	2	2	2	2	2	2	2	2	103
476	4X	5075.452	4	4	4	4	4	5	5	5	5	5	5	177
477	4X	4230.688	5	4	5	5	6	5	5	4	4	6	4	204
478	4X	800.122	2	2	2	2	2	2	2	3	2	2	2	119
480	4X	2249.270	8	7	8	8	6	7	7	7	5	7	5	306
481	4X	6438.750	8	10	9	9	9	8	10	9	6	9	6	350
482	4X	3578.228	3	4	3	3	3	3	4	4	3	4	3	141
483	4X	1826.888	2	3	2	2	2	2	3	3	2	3	2	105
484	4X	7774.576	3	5	5	5	4	6	5	7	7	7	7	186
485	4X	5432.588	5	6	5	5	5	6	6	6	4	6	5	196
490	4X	2063.834	3	4	2	4	3	4	4	4	3	4	3	173
491	4X	2359.158	4	4	4	4	4	4	4	4	3	4	3	168
492	4X	3729.324	4	6	4	4	4	3	4	4	3	4	4	171
493	4X	1830.322	3	4	4	4	3	3	4	6	3	3	3	159
494	4X	1431.978	4	4	4	4	3	4	4	3	2	4	3	128
495	4X	2005.456	3	4	4	4	2	4	4	4	3	4	3	127
		171809.888	196	243	210	208	196	212	214	208	81	227	175	9080

421 **3.2 Distribution of depth, bottom temperature and bottom salinity from survey tows**

422 The depth, bottom temperature and bottom salinity cumulative frequency distribution for the
423 survey are presented in Figure 5.

424 **4 Discussion**

425 This report builds on previous work and former atlases by updating a comprehensive suite of
426 indices to give a snapshot of population status and environmental preferences of 104 fish and
427 invertebrate species. The current document is not meant to replace stock assessments, species-
428 specific analyses of abundance, biomass and distribution, or any targeted attempts to integrate
429 information about species or group of species from the wide and disparate sources of data about
430 marine organisms in the area covered by the DFO Maritimes summer trawl survey. It is rather
431 meant to provide a reproducible set of tools to extract and visualize the information collected
432 in the summer groundfish research vessel survey. It is hoped that this document can provide a
433 stepping stone to conduct other ecological analyses using the trawl survey data and increase
434 reproducibility and transparency of ecological information collected annually.

435 **4.1 Diversity of approaches used for mapping fish and invertebrates in the Scotian Shelf
436 bioregion**

437 Different methods have been applied in the Northwest Atlantic, and specifically on the Scotian
438 Shelf bioregion, to map fish and invertebrate species distribution. The present report, for
439 example, builds upon the atlas of important habitat developed to map the persistence of relatively
440 high biomass for key fish species using the summer groundfish research vessel survey (Horsman
441 and Shackell 2009). Important habitat was obtained by interpolating observed weight per each
442 species using an inverse-distance weighted (IDW) methodology, and calculating areas with
443 relatively persistent high biomass for periods representing different fishery management eras.
444 To compliment information from this atlas, including additional representations of biomass and
445 diversity, a similar IDW interpolation mapping procedure was followed by Smith et al. (2015),
446 Ward-Paige and Bundy (2015), and Bundy et al. (2017). The summer groundfish research vessel
447 survey is typically conducted during the month of July. However, from the fall of 1978 through to
448 the spring of 1985, DFO also conducted spring and fall surveys using the same sampling design.
449 This unique seasonal data was used to map the seasonal spatial distribution of key demersal
450 and other fish species using IDW interpolation on the Scotian Shelf from the spring, summer
451 and fall between 1978 and 1985 (Smith et al. 2015). Following recommendations provided by
452 Kenchington and Kenchington (2017), the spatial distribution of three indicators of biodiversity
453 for fish and invertebrates were mapped using IDW interpolation to identify areas with persistently
454 high values across fishery management eras, and compared with areas of persistently high
455 abundance for selected species (Ward-Paige and Bundy 2015). This analysis revealed a lack of
456 consistent relationships between areas of persistent high diversity and persistent high biomass,
457 suggesting that both can be used as independent and important spatial indicators of the system
458 (Ward-Paige and Bundy 2015). Groupings of fishes and invertebrates based on size, habitat
459 and feeding guild, were also mapped using IDW interpolations to identify hotspots of functional

460 group diversity (Bundy et al. 2017). This analysis revealed a spatially and temporally variable
461 distribution of functional diversity across the Scotian Shelf with notable areas of high and low
462 diversity (Bundy et al. 2017). Top quintiles of each functional group using the IDW approach
463 were used as representative layers for fish and invertebrates in the MPA Network design in the
464 Scotian Shelf Bioregion (Serdynska et al. In press). IDW interpolation methods have also been
465 used to map the distribution of individual species such as sea cucumbers (*Cucumaria frondosa*)
466 in the Scotian Shelf bioregion (Shackell et al. 2013), and sea scallop (*Placopecten magellanicus*)
467 in Georges and Browns Bank (Hubley et al. 2014).

468 Species Distribution Modelling (SDM), instead of IDW, can also be used to evaluate spatio-
469 temporal dynamics by predicting and understanding past, present and future distribution
470 of species using environmental predictors (Robinson et al. 2017). A variety of modelling
471 approaches are being implemented in Maritimes Region to map and predict fish and invertebrate
472 species distribution by incorporating environmental predictors to account for seasonal and
473 temporal variability. For example, a stock assessment of snow crab (*Chionoecetes opilio*) on
474 the Scotian Shelf used data from the snow crab survey from 2005 to 2018 to map spatial data
475 products for this stock, including annual predicted interpolations of potential habitat using
476 Generalized Additive Models (GAM) and several environmental covariates including depth,
477 curvature, slope, species composition, and annual temperature (Zisserson et al. 2019). Sea
478 scallop predicted habitat using Maximum Entropy (MaxEnt) models were computed for German
479 Bank using data compiled via benthic habitat mapping and seafloor geotechnical surveys in
480 2006, 2009, and 2010 (Brown et al. 2012). Predictions in the Scotian Shelf bioregion and the
481 Northeast United States using datasets from DFO and the National Oceanic and Atmospheric
482 Administration from 1993 to 2012 also predicted sea scallop habitat at a wider scale based
483 on three scenarios of seasonal temperature and salinity climatologies (NOAA) (Lowen et
484 al. 2019). Offshore American lobster stock assessments (*Homarus americanus*) used data
485 from the RV, DFO Georges Bank, and National Marine Fisheries Service (NMFS) Northeast
486 Fisheries Science Center (NEFSC) bottom trawl surveys (1970 to 2015) to predict species
487 distribution using boosted regression trees and several environmental predictors (bathymetry,
488 slope, curvature, and annual temperature interpolations) (Cook et al. 2017). Information on
489 the potential for recovery of cusk (*Brosme brosme*) used data from the bottom longline Halibut
490 industry survey and Cusk absences in the Summer groundfish research vessel survey from
491 1998-2013 to predict suitable habitat using GAM, MaxEnt, and random forest models and
492 several physical environmental variables (e.g. complexity, benthic current stress and complexity,
493 temperature, salinity, primary production, chlorophyll, suspended matter) (Harris et al. 2018).
494 Atlantic halibut (*Hippoglossus hippoglossus*) assessments using Summer groundfish research
495 vessel survey and NOAA survey data from 2001 to 2013 predicted juvenile habitat using MaxEnt
496 model and environmental predictors (bathymetry, slope, bottom temperature) (French et al.
497 2018). Persistent areas of high Atlantic halibut juvenile abundance were predicted using data
498 from 27 bottom trawl surveys combined (NMFS and DFO) from 1978 to 2013 and applying
499 Bayesian hierarchical spatiotemporal models with two environmental predictors (depth and
500 temperature) (Boudreau et al. 2017).

501 These examples of mapping efforts in Maritimes Region showcase the diversity of approaches
502 relevant to a variety of important research questions and management applications. Approaches,
503 methods, datasets, and environmental predictors are selected based on individual project
504 research questions, and considerations for each species, communities or stock. This allows
505 research groups to maintain innovation and keep up with emerging methods and technologies to

506 improve assessments, predictions, and ultimately, science advice. The diversity of approaches
507 also leads to complexity when looking across studies as each data compilation and predictive
508 method carries its own independent assumptions and can lead to different spatial outputs.

509 **4.2 Interpreting spatial results for marine spatial planning purposes**

510 Fisheries and Oceans Canada is leading a marine spatial planning process that brings together
511 relevant authorities and stakeholders to better coordinate how we use and manage marine
512 spaces to achieve ecological, economic and social objectives. Operationalizing marine spatial
513 planning includes a series of steps, including the process of analyzing existing conditions
514 by collecting and mapping information about ecological, environmental and oceanographic
515 conditions (Ehler and Douvere 2009; Agardy et al. 2011). Mapping the distribution of species
516 is critical for the implementation of spatial management and as a first step in marine spatial
517 planning processes. Species distribution have supported the identification of important sites for
518 a given species or areas of high richness and diversity, which in turn can be used to inform siting
519 decisions of new activities such as Marine Protected Areas (MPA), aquaculture sites or wind
520 turbines. In the Scotian Shelf bioregion, mapping species distributions has been used to highlight
521 areas of high biological diversity to support the identification of Ecologically or Biologically
522 Significant Areas (Ricard and Shackell 2013; Ward-Paige and Bundy 2015), to distinguish
523 important and persistent habitat of significant species and functional groups to support MPA and
524 conservation planning (Horsman and Shackell 2009; Smith et al. 2015; Ward-Paige and Bundy
525 2015; Bundy et al. 2017), to identify important habitat for Species at Risk (Harris et al. 2018) and
526 to highlight reserves for data-poor invertebrate fisheries (Shackell et al. 2013). Mapping species
527 distribution has also been used to illustrate multi-decadal scale projections of changes in species
528 distribution in the context of climate change and adaption (Stanley et al. 2018; Greenan et al.
529 2019).

530 In support of the marine spatial planning process, a public web-based atlas with relevant
531 geospatial information is being developed to support decision-making. This Atlantic Canada-
532 wide compilation of data and information will be a web-based, public platform with interactive
533 maps of ocean ecosystems, human uses and management areas. The current document cannot
534 present the full diversity of data and mapping products that can be produced for the Maritimes
535 Region. Consequently, we recommend that the data and mapping products presented in this
536 report not be used blindly for the planned atlas, until an evaluation of what spatial information is
537 available and what was used in the past is conducted.

538 This diverse portfolio of approaches and applications is not unique to the Maritimes Region. A
539 recent review of global distribution modelling efforts recommended the adoption of a consistent
540 framework that integrates multi-model approaches and a clear expression of errors and
541 uncertainties (Robinson et al. 2017). In this context, Pacific Region has developed two initiatives
542 to enable consistency and frequent publication, reproducibility, and transparency. One initiative
543 developed a reproducible report to give a synthesis of data availability, population trends, fishing
544 trends, growth and maturity patterns for 113 groundfish species in British Columbia to support
545 stock assessment (Anderson et al. 2019). The second initiative developed a SDM framework
546 that was applied to twelve species on Canada's Pacific coast as part of the Regional Response
547 Plan (Nephin et al. 2019). The Maritimes and Gulf region, through this and past reports, are also

548 using similar reproducible approaches to facilitate annual updates and transparency (Ricard and
549 Shackell 2013; Ricard et al. 2021).

550 Recognizing the diversity of approaches for mapping fish and invertebrates in the Scotian Shelf
551 bioregion, we recommend the development of a regional community of practice to compare
552 and evaluate approaches for mapping, interpolating and/or modelling fish and invertebrates
553 so future publications and advice related to spatial outputs can lead to more comparable work
554 and consistent science advice to support processes such as marine spatial planning. At the
555 international level, guidelines and standards related to appropriate variables and methods
556 for mapping and modeling species and communities of deep-sea habitats were proposed to
557 encourage the production of publications that will lead to more comparable work (Kenchington
558 et al. 2019). Similar general guidance for how groups approach mapping activities would be a
559 worthwhile product in Maritimes Region. Until then, we propose the use of the Open Data record
560 for the Maritimes RV surveys (DFO 2021) as a precursor to the public web-based marine spatial
561 planning atlas.

562 **5 Acknowledgements**

563 We thank all the dedicated personnel involved in running trawl surveys in the Maritimes Region
564 and the numerous colleagues in Maritimes Region that have shared information and advice in
565 support of this report. The assistance of the Gulf Region secondary publications coordinators,
566 Alicia Cassidy and Jeff Clements, in getting this report published is well appreciated. This
567 document greatly benefited from the constructive comments of two reviewers.




Figure 5. Cumulative frequency distribution of bottom salinity (top panel), bottom temperature (middle panel) and depth (bottom panel) of representative sets from the DFO Maritimes summer survey.

6 References

- 569 Agardy, T., Notarbartolo di Sciara, G., and Christie, P. 2011. [Addressing the shortcomings of](#)
 571 [marine protected areas through large scale marine spatial planning](#). Marine Policy 35(2):
 572 226–232.
- 573 Anderson, S.C., Grandin, C., Edwards, A.M., Grinnell, M.H., Ricard, D., and Haigh, R. 2021.
 574 Csasdown: Reproducible CSAS Reports with Bookdown.
- 575 Anderson, S.C., Keppel, E.A., and Edwards, A.M. 2019. [A reproducible data synopsis for](#)
 576 [over 100 species of British Columbia groundfish](#). DFO Can. Sci. Advis. Sec. Res. Doc.
 577 2019/041. vii + 321 p.
- 578 Appeltans, W., Bouchet, P., Boxshall, G.A., De Broyer, C., Voogd, N.J. de, Gordon, D.P.,
 579 Hoeksema, B.W., Horton, T., Kennedy, M., J., M., Poore, G.C.B., Read, G., Stöhr,
 580 S., Walter, T.C., and Costello, M.J. (*Editors*). 2012. World register of marine species.
 581 Accessed at <http://www.marinespecies.org> on 2020-12-1.
- 582 Baddeley, R., A. 2015. Spatial point patterns: Methodology and applications
 583 with R. Chapman; Hall/CRC Press, London [http://www.crcpress.com/
 584 Spatial-Point-Patterns-Methodology-and-Applications-with-R/
 585 Baddeley-Rubak-Turner/9781482210200/](http://www.crcpress.com/Spatial-Point-Patterns-Methodology-and-Applications-with-R-Baddeley-Rubak-Turner/9781482210200/).
- 586 Benoît, H.P., Abgrall, M.-J., and Swain, D.P. 2003. [An assessment of the general status of](#)
 587 [marine and diadromous fish species in the southern Gulf of St. Lawrence based on annual](#)
 588 [bottom trawl surveys \(1971-2002\)](#). Can. Tech. Rep. Fish. Aquat. Sci. 2472: iv + 183 p.
- 589 Bivand, R. 2020. classInt: Choose univariate class intervals. R package version 0.4-3 <https://CRAN.R-project.org/package=classInt>.
- 590 Bivand, R., and Lewin-Koh, N. 2020. Maptools: Tools for handling spatial objects. R package
 591 version 1.0-2 <https://CRAN.R-project.org/package=maptools>.
- 592 Bivand, R., and Rundel, C. 2020. Rgeos: Interface to geometry engine - open source
 593 ('GEOS'). R package version 0.5-5 <http://CRAN.R-project.org/package=rgeos>.
- 594 Boudreau, S.A., Shackell, N.L., Carson, S., and Heyer C. E., den. 2017. [Connectivity,](#)
 595 [persistence, and loss of high abundance areas of a recovering marine fish population in](#)
 596 [the Northwest Atlantic Ocean](#). Ecol. Evol. 7: 9739–9749.
- 597 Bourdages, H., and Ouellet, J.-F. 2012. [Geographic distribution and abundance indices of](#)
 598 [marine fish in the northern Gulf of St. Lawrence \(1990-2009\)](#). Can. Tech. Rep. Fish.
 599 Aquat. Sci. 2963: vi + 171 p.
- 600 Brown, C., Sameoto, J., and Smith, S. 2012. [Multiple methods, maps, and management](#)
 601 [applications: Purpose made seafloor maps in support of ocean management](#). Journal of
 602 Sea research 72: 113.
- 603 Bundy, A., Will, E., Serdynska, A., Cook, A., and Ward-Paige, C.A. 2017. [Defining and](#)
 604 [mapping functional groups for fishes and invertebrates in the Scotian Shelf bioregion](#). Can.
 605 Tech. Rep. Fish. Aquat. Sci. 3186: iv + 49 p.

- 607 Clark, D.W., and Emberley, J. 2011. Update of the 2010 summer scotian shelf and bay of
608 fundy research vessel survey. Can. Tech. Rep. Fish. Aquat. Sci.: 1238: ix + 98 p.
- 609 Cook, A.M., Cassista Da-Ros, M., and Denton, C. 2017. [Framework Assessment of the](#)
610 [Offshore American Lobster in Lobster Fishing Area \(LFA\) 41.](#) ICES Journal of Marine
611 Science 2017/065 viii + 186 p.
- 612 DFO. 2016. DFO maritimes research vessel trawl surveys invertebrate observations. Version
613 7 in OBIS canada digital collections. Bedford Institute of Oceanography, Dartmouth, NS,
614 Canada, Published by OBIS, Digital 2016.
- 615 DFO. 2021. Maritimes research vessel surveys [dataset]. Retrieved from
616 <https://open.canada.ca/data/en/dataset/8ddcaeeaa-b806-4958-a79f-ba9ab645f53b>.
- 617 Doubleday, W.G., and Rivard, D. 1981. Bottom trawl surveys. Can. Spec. Publ. Fish. Aquat.
618 Sci: 58: 237 pp.
- 619 Ehler, C., and Douvere, F. 2009. Marine spatial planning: A step-by-step approach.
- 620 French, K., Shackell, N., and Heyer, N. den. 2018. [Information on the Potential for Recovery](#)
621 [of Cusk in Canadian Waters.](#) Fish. Bull. 116: 107–121.
- 622 Greenan, B.J.W., Shackell, N.L., Ferguson, K., Greyson, P., Cogswell, A., Brickman, D.,
623 Wang, Z., Cook, A., Brennan, C.E., and Saba, V.S. 2019. Climate change vulnerability
624 of American lobster fishing communities in Atlantic Canada. Frontiers in Marine Science 6:
625 579.
- 626 Harris, L.E., Greenlaw, M., McCurdy, D., and MacDonald, D. 2018. [Information on the](#)
627 [Potential for Recovery of Cusk in Canadian Waters.](#) DFO Can. Sci. Advis. Sec. Res. Doc.
628 2018/002. vi + 62 p.
- 629 Holstein, J. 2018. Worms: Retrieving aphia information from world register of marine species.
630 R package version 0.2.2 <https://CRAN.R-project.org/package=worms>.
- 631 Horsman, T.L., and Shackell, N.L. 2009. [Atlas of important habitat for key fish species of the](#)
632 [Scotian Shelf, Canada.](#) Can. Tech. Rep. Fish. Aquat. Sci. 2835: viii + 82 p.
- 633 Hubley, P.B., Reeves, A., Smith, S.J., and Nasmyth, L. 2014. [Georges Bank 'a' and Browns](#)
634 [Bank 'North' Scallop \(*Placopecten magellanicus*\) Stock Assessment.](#) DFO Can. Sci.
635 Advis. Sec. Res. Doc. 2013/079. vi + 58 p.
- 636 Kenchington, E., Callery, O., Davidson, F., Grehan, A., Morato, T., Appiott, J., Davis, A.,
637 Dunstan, P., Du Preez, C., Finney, J., González-Irusta, J.M., Howell, K., Knudby, A.,
638 Lacharité, M., Lee, J., Murillo, F.J., Beazley, L., Roberts, J.M., Roberts, M., Rooper, C.,
639 Rowden, A., Rubidge, E., Stanley, R., Stirling, D., Tanaka, K.R., Vanhatalo, J., Weigel, B.,
640 Woolley, S., and Yesson, C. 2019. [Use of Species Distribution Modeling in the Deep Sea.](#)
641 Can. Tech. Rep. Fish. Aquat. Sci. 3296: ix + 76 p.
- 642 Kenchington, T.J., and Kenchington, E.L.R. 2017. [Biodiversity metrics for use in the](#)
643 [ecosystem approach to oceans management.](#) Can. Tech. Rep. Fish. Aquat. Sci. 3186:
644 iv + 49 p.

- 645 Lohr, S. 1999. Sampling: Design and analysis. Pacific Grove, CA: Brooks/Cole Publishing
646 Company.
- 647 Losier, R.J., and Waite, L.E. 1989. Systematic listing of scientific and/or common names of
648 invertebrates and marine plants and their respective codes used by marine fish division,
649 Fisheries and Oceans, Scotia-Fundy Region. Canadian Data Report of Fisheries and
650 Aquatic Sciences (721).
- 651 Lowen, B., Hart, D., Stanley, R., Lehnert, S., Bradbury, I., and C., D. 2019. [Assessing effects](#)
652 [of genetic, environmental, and biotic gradients in species distribution modelling](#). ICES
653 Journal of Marine Science 76(6): 1762–1775.
- 654 Myers, R.A., and Stokes, K. 1989. Density-dependent habitat utilization of groundfish and the
655 improvement of research surveys. (D:15). International Council for the Exploration of the
656 Sea Council Meeting.
- 657 Nephin, J., Gregr, E.J., St. Germain, C., Fields, C., and Finney, J.L. and. 2019. [Development](#)
658 [of a Species Distribution Modelling Framework and its Application to Twelve Species on](#)
659 [Canada's Pacific Coast](#). DFO Can. Sci. Advis. Sec. Res. Doc. 2020/004. xii + 107 p.
- 660 Neuwirth, E. 2014. RColorBrewer:ColorBrewer palettes. R package version 1.1-2 <https://CRAN.R-project.org/package=RColorBrewer>.
- 661
- 662 Pebesma, E. 2004. Multivariable geostatistics in S: The gstat package. *In* Computers and
663 Geosciences.
- 664 Perry, R.I., and Smith, S.J. 1994. Identifying habitat associations of marine fishes using
665 survey data: An application to the northwest atlantic. Canadian Journal of Fisheries and
666 Aquatic Sciences (51(3)): 589–602.
- 667 R Core Team. 2020. R: A language and environment for statistical computing. R Foundation
668 for Statistical Computing, Vienna, Austria.
- 669 Ricard, D., and Gomez, C. 2021. Maritimes-SUMMER-atlas. <https://github.com/dfo-gulf-science/Maritimes-SUMMER-Atlas>; GitHub.
- 670
- 671 Ricard, D., Rolland, N., and Swain, D. 2021. Occurrence, geographic distribution and
672 abundance indices of marine organisms caught in the september research vessel trawl
673 survey in the southern Gulf of St. Lawrence (1971-2020). Can. Tech. Rep. Fish. Aquat.
674 Sci. xxxx: viii + 180 p.
- 675 Ricard, D., and Shackell, N.L. 2013. [Population status \(abundance/biomass, geographic](#)
676 [extent, body size and condition\), important habitat, depth, temperature and salinity of](#)
677 [marine fish and invertebrates on the Scotian Shelf and Bay of Fundy \(1970-2012\)](#). Can.
678 Tech. Rep. Fish. Aquat. Sci. 3012: viii + 180 p.
- 679 Ripley, B., and Lapsley, M. 2019. RODBC: ODBC database access. R package version 1.3-16
680 <http://CRAN.R-project.org/package=RODBC>.
- 681 Ripley, B., Venables, B., Bates, D., Hornik, K., Gebhardt, A., and Firth, D. 2020. Modern
682 applied statistics with s. R package version 7.3-53 <https://cran.r-project.org/web/packages/MASS/index.html>.
- 683

- 684 Robinson, N.M., Nelson, W.A., Costello, M.J., Sutherland, J.E., and Lundquist, C.J.
685 2017. Systematic review of marine-based species distribution models (SDMs) with
686 recommendations for best practice. *Front. Mar. Sci* (4): 421.
- 687 Schnute, J.T., Boers, N., and Haigh, R. 2019. PBSmapping: Mapping fisheries data and
688 spatial analysis tools. R package version 2.72.1 <https://cran.r-project.org/web/packages/PBSmapping/index.html>.
- 689
- 690 Shackell, N., Brickman, D., and Frank, K. 2013. [Reserve site selection for data-poor
691 invertebrate fisheries using patch scale and dispersal dynamics: A case study of sea
692 cucumber](#). *Aquatic Conserv: Mar. Freshw. Ecosyst.* 23: 723–731.
- 693 Simon, J.E., and Comeau, P.A. 1994. [Summer distribution and abundance trends of species
694 caught on the Scotian Shelf from 1970-92, by the research vessel groundfish survey](#). *Can.
695 Tech. Rep. Fish. Aquat. Sci.* 1953.
- 696 Smith, C.D., Serdynska, A.R., King, M.C., and Shackell, N.L. 2015. [Spring, summer and
697 fall distribution of common demersal fishes on the Scotian Shelf between 1978 and 1985](#).
698 *Can. Tech. Rep. Fish. Aquat. Sci.* 3068: vi + 38 p.
- 699 Smith, S.J. 1996. Assessment of groundfish stocks based on bottom trawl survey results.
700 *NAFO Scientific Council Studies* 28: 25–53.
- 701 Stanley, R.E., DiBacco, C., Lowen, B., Beiko, R., Jeffery, N., Wyngaarden, M., Bentzen, P.,
702 Brickman, D., Benestan, L., Bernatchez, L., Johnson, C., Snelgrove, P., Wang, Z., and
703 Wringe, I., B. Bradbury. 2018. A climate-associated multispecies cryptic cline in the
704 northwest Atlantic. *Science Advances* 4(3): 1–7.
- 705 Swain, D.P., and Morin, R. 1996. Relationships between geographic distribution and
706 abundance of American plaice (*Hippoglossoides platessoides*) in the southern Gulf of
707 St. Lawrence. *Canadian Journal of Fisheries and Aquatic Sciences* 53(1): 106–119.
- 708 Swain, D.P., and Sinclair, A.F. 1994. Fish distribution and catchability: What is the appropriate
709 measure of distribution? *Canadian Journal of Fisheries and Aquatic Sciences* 51(5):
710 1046–1054.
- 711 Tremblay M. J., B.R., Black G. A. P. 2007. The distribution of common decapod crustaceans
712 and other invertebrates recorded in annual ecosystem surveys of the scotian shelf 1999-
713 2006, by the research vessel groundfish survey. *Can. Tech. Rep. Fish. Aquat. Sci.* 74.
- 714 Ward-Paige, C.A., and Bundy, A. 2015. [Mapping Biodiversity on the Scotian Shelf and in the
715 Bay of Fundy](#). *Can. Tech. Rep. Fish. Aquat. Sci.* 3068: vi + 38 p.
- 716 Wickham, H. 2019. Tidyverse: Easily install and load the 'tidyverse'. R package version 1.3.0
717 <https://cran.r-project.org/web/packages/tidyverse/index.html>.
- 718 Zisserson, B.M., Cameron, B.J., Glass, A.C., and Choi, J.S. 2019. [Assessment of Scotian
719 Shelf Snow Crab in 2017](#). DFO Can. Sci. Advis. Sec. Res. Doc. 2018/051. ix + 147 p.

7 Appendix

721

7.1 Atlantic cod (*Morue franche*) - species code 10 (category LF)

722

Scientific name: [Gadus morhua](#)

723




Figure 7.1A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic cod.

724




Figure 7.1B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic cod.




Figure 7.1C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic cod.






Figure 7.1D. Average fish condition in NAFO units 4X and 4VW for Atlantic cod.



727



Freq	Depth	Temp	Sal
F5	26	1.2	31.00
F25	43	3.5	32.47
F50	60	6.0	33.27
F75	82	8.1	34.40
F95	126	10.0	35.03

Figure 7.1E. Catch distribution by depth, temperature and salinity of Atlantic cod.

728




Figure 7.1F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic cod.

729

7.2 Haddock (Aiglefin) - species code 11 (category LF)

730

Scientific name: [Melanogrammus aeglefinus](#)

731




Figure 7.2A. Inverse distance weighted distribution of catch biomass (kg/tow) for Haddock.

732




Figure 7.2B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Haddock.




Figure 7.2C. Length frequency distribution in NAFO units 4X and 4VW for Haddock.






Figure 7.2D. Average fish condition in NAFO units 4X and 4VW for Haddock.



735



Freq	Depth	Temp	Sal
F5	27	1.1	31.00
F25	49	3.0	32.45
F50	66	5.5	33.14
F75	87	7.9	34.36
F95	127	10.0	35.03

Figure 7.2E. Catch distribution by depth, temperature and salinity of Haddock.




Figure 7.2F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Haddock.

737

7.3 White hake (Merluche blanche) - species code 12 (category LF)

738

Scientific name: [Urophycis tenuis](#)

739




Figure 7.3A. Inverse distance weighted distribution of catch biomass (kg/tow) for White hake.

740




Figure 7.3B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of White hake.




Figure 7.3C. Length frequency distribution in NAFO units 4X and 4VW for White hake.






Figure 7.3D. Average fish condition in NAFO units 4X and 4VW for White hake.




743



Freq	Depth	Temp	Sal
F5	40	1.1	31.00
F25	70	3.0	32.46
F50	89	5.5	33.20
F75	115	7.9	34.39
F95	163	10.0	35.04

Figure 7.3E. Catch distribution by depth, temperature and salinity of White hake.



744

Figure 7.3F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for White hake.

745

7.4 Red hake (Merluche écureuil) - species code 13 (category LF)

746

Scientific name: [Urophycis chuss](#)

747




Figure 7.4A. Inverse distance weighted distribution of catch biomass (kg/tow) for Red hake.

748




Figure 7.4B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Red hake.




Figure 7.4C. Length frequency distribution in NAFO units 4X and 4VW for Red hake.






Figure 7.4D. Average fish condition in NAFO units 4X and 4VW for Red hake.



751



Freq	Depth	Temp	Sal
F5	35	1.1	31.00
F25	62	2.9	32.43
F50	82	5.4	33.12
F75	103	7.9	34.32
F95	130	10.0	35.08

Figure 7.4E. Catch distribution by depth, temperature and salinity of Red hake.

752




Figure 7.4F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Red hake.

753

7.5 Silver hake (*Merlu argenté*) - species code 14 (category LF)

754

Scientific name: [Merluccius bilinearis](#)

755




Figure 7.5A. Inverse distance weighted distribution of catch biomass (kg/tow) for Silver hake.

756




Figure 7.5B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Silver hake.




Figure 7.5C. Length frequency distribution in NAFO units 4X and 4VW for Silver hake.






Figure 7.5D. Average fish condition in NAFO units 4X and 4VW for Silver hake.



759



Freq	Depth	Temp	Sal
F5	28	1.2	31.00
F25	55	3.1	32.46
F50	77	5.5	33.20
F75	104	8.0	34.37
F95	137	10.0	35.07

Figure 7.5E. Catch distribution by depth, temperature and salinity of Silver hake.

760




Figure 7.5F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Silver hake.

761

7.6 Pollock (Goberge) - species code 16 (category LF)

762

Scientific name: [Pollachius virens](#)

763




Figure 7.6A. Inverse distance weighted distribution of catch biomass (kg/tow) for Pollock.

764




Figure 7.6B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Pollock.




Figure 7.6C. Length frequency distribution in NAFO units 4X and 4VW for Pollock.






Figure 7.6D. Average fish condition in NAFO units 4X and 4VW for Pollock.



767



Freq	Depth	Temp	Sal
F5	37	1.1	31.00
F25	60	2.8	32.45
F50	82	5.3	33.14
F75	108	7.7	34.33
F95	137	10.0	35.03

Figure 7.6E. Catch distribution by depth, temperature and salinity of Pollock.

768




Figure 7.6F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Pollock.

769 **7.7 Atlantic redfishes (Sébastes de l'Atlantique) - species code 23 (category LF)**

770 Scientific name: [Sebastes](#)

771




Figure 7.7A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic redfishes.

772




Figure 7.7B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic redfishes.




Figure 7.7C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic redfishes.






Figure 7.7D. Average fish condition in NAFO units 4X and 4VW for Atlantic redfishes.



775



Freq	Depth	Temp	Sal
F5	47	1.2	31.00
F25	68	3.4	32.48
F50	86	5.9	33.29
F75	114	8.1	34.41
F95	166	10.0	35.05

Figure 7.7E. Catch distribution by depth, temperature and salinity of Atlantic redfishes.




Figure 7.7F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic redfishes.

777

7.8 Atlantic halibut (Flétan de l'Atlantique) - species code 30 (category LF)

778

Scientific name: [Hippoglossus hippoglossus](#)

779




Figure 7.8A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic halibut.

780




Figure 7.8B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic halibut.




Figure 7.8C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic halibut.






Figure 7.8D. Average fish condition in NAFO units 4X and 4VW for Atlantic halibut.



783



Freq	Depth	Temp	Sal
F5	31	1.3	31.00
F25	49	3.2	32.45
F50	60	5.8	33.16
F75	75	8.3	34.34
F95	122	10.0	35.08

Figure 7.8E. Catch distribution by depth, temperature and salinity of Atlantic halibut.

784




Figure 7.8F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic halibut.

785

7.9 American plaice (*Ple canadienne*) - species code 40 (category LF)

786

Scientific name: [Hippoglossoides platessoides](#)

787




Figure 7.9A. Inverse distance weighted distribution of catch biomass (kg/tow) for American plaice.

788




Figure 7.9B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of American plaice.




Figure 7.9C. Length frequency distribution in NAFO units 4X and 4VW for American plaice.






Figure 7.9D. Average fish condition in NAFO units 4X and 4VW for American plaice.



791



Freq	Depth	Temp	Sal
F5	29	1.3	31.00
F25	46	3.7	32.48
F50	67	6.1	33.27
F75	87	8.1	34.41
F95	133	10.0	35.05

Figure 7.9E. Catch distribution by depth, temperature and salinity of American plaice.

792




Figure 7.9F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for American plaice.

793

7.10 Witch flounder (*Ple grise*) - species code 41 (category LF)

794

Scientific name: [Glyptocephalus cynoglossus](#)

795




Figure 7.10A. Inverse distance weighted distribution of catch biomass (kg/tow) for Witch flounder.

796




Figure 7.10B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Witch flounder.




Figure 7.10C. Length frequency distribution in NAFO units 4X and 4VW for Witch flounder.






Figure 7.10D. Average fish condition in NAFO units 4X and 4VW for Witch flounder.



799



Freq	Depth	Temp	Sal
F5	32	1.3	31.00
F25	57	3.5	32.49
F50	77	6.1	33.30
F75	102	8.2	34.45
F95	152	10.0	35.06

Figure 7.10E. Catch distribution by depth, temperature and salinity of Witch flounder.




Figure 7.10F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Witch flounder.

801 **7.11 Yellowtail flounder (Limande à queue jaune) - species code 42 (category LF)**

802 Scientific name: [Limanda ferruginea](#)

803




Figure 7.11A. Inverse distance weighted distribution of catch biomass (kg/tow) for Yellowtail flounder.

804




Figure 7.11B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Yellowtail flounder.




Figure 7.11C. Length frequency distribution in NAFO units 4X and 4VW for Yellowtail flounder.






Figure 7.11D. Average fish condition in NAFO units 4X and 4VW for Yellowtail flounder.




807



Freq	Depth	Temp	Sal
F5	22	1.4	31.00
F25	32	3.7	32.52
F50	43	6.3	33.25
F75	56	8.2	34.41
F95	81	10.0	35.06

Figure 7.11E. Catch distribution by depth, temperature and salinity of Yellowtail flounder.



808

Figure 7.11F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Yellowtail flounder.


809

7.12 Winter flounder (Limande-plie rouge) - species code 43 (category LF)

810

Scientific name: [Pseudopleuronectes americanus](#)

811



812




Figure 7.12B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Winter flounder.




Figure 7.12C. Length frequency distribution in NAFO units 4X and 4VW for Winter flounder.






Figure 7.12D. Average fish condition in NAFO units 4X and 4VW for Winter flounder.



815



Freq	Depth	Temp	Sal
F5	22	1.1	31.00
F25	31	3.0	32.48
F50	42	5.5	33.17
F75	54	8.3	34.47
F95	84	10.0	35.10

Figure 7.12E. Catch distribution by depth, temperature and salinity of Winter flounder.

816




Figure 7.12F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Winter flounder.

817

7.13 Atlantic wolffish (*Loup atlantique*) - species code 50 (category LF)

818

Scientific name: [Anarhichas lupus](#)

819




Figure 7.13A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic wolffish.

820




Figure 7.13B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic wolffish.




Figure 7.13C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic wolffish.







Figure 7.13D. Average fish condition in NAFO units 4X and 4VW for Atlantic wolffish.



823



Freq	Depth	Temp	Sal
F5	34	1.4	31.00
F25	49	3.6	32.48
F50	62	6.0	33.25
F75	77	8.1	34.33
F95	112	10.0	35.05

Figure 7.13E. Catch distribution by depth, temperature and salinity of Atlantic wolffish.

824




Figure 7.13F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic wolffish.

825 **7.14 Atlantic herring (Hareng de l'Atlantique) - species code 60 (category LF)**

826 Scientific name: [Clupea harengus](#)

827




Figure 7.14A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic herring.

828




Figure 7.14B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic herring.




Figure 7.14C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic herring.






Figure 7.14D. Average fish condition in NAFO units 4X and 4VW for Atlantic herring.



831



Freq	Depth	Temp	Sal
F5	32	1.2	31.00
F25	54	3.2	32.45
F50	74	5.6	33.22
F75	95	8.3	34.38
F95	132	10.0	35.10

Figure 7.14E. Catch distribution by depth, temperature and salinity of Atlantic herring.

832




Figure 7.14F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic herring.

833 **7.15 Longhorn sculpin (Chaboisseau à dix-huit épines) - species code 300 (category**
 834 **LF)**

835 Scientific name: [Myoxocephalus octodecemspiniferus](#)

836




Figure 7.15A. Inverse distance weighted distribution of catch biomass (kg/tow) for Longhorn sculpin.

837




Figure 7.15B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Longhorn sculpin.




Figure 7.15C. Length frequency distribution in NAFO units 4X and 4VW for Longhorn sculpin.




Figure 7.15D. Average fish condition in NAFO units 4X and 4VW for Longhorn sculpin.




Figure 7.15E. Catch distribution by depth, temperature and salinity of Longhorn sculpin.




Figure 7.15F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Longhorn sculpin.

842

7.16 Moustache sculpin (Faux-trigle armé) - species code 304 (category LF)

843

Scientific name: [Triglops murrayi](#)

844




Figure 7.16A. Inverse distance weighted distribution of catch biomass (kg/tow) for Moustache sculpin.

845




Figure 7.16B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Moustache sculpin.




Figure 7.16C. Length frequency distribution in NAFO units 4X and 4VW for Moustache sculpin.






Figure 7.16D. Average fish condition in NAFO units 4X and 4VW for Moustache sculpin.



848



Freq	Depth	Temp	Sal
F5	30	1.5	31.00
F25	40	4.0	32.48
F50	50	6.5	33.31
F75	63	8.2	34.39
F95	88	10.0	35.06

Figure 7.16E. Catch distribution by depth, temperature and salinity of Moustache sculpin.

849




Figure 7.16F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Moustache sculpin.

850 **7.17 Sea raven (Hémithriptère atlantique) - species code 320 (category LF)**

851 Scientific name: [Hemitripterus americanus](#)




Figure 7.17A. Inverse distance weighted distribution of catch biomass (kg/tow) for Sea raven.




Figure 7.17B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Sea raven.




Figure 7.17C. Length frequency distribution in NAFO units 4X and 4VW for Sea raven.






Figure 7.17D. Average fish condition in NAFO units 4X and 4VW for Sea raven.



856



Freq	Depth	Temp	Sal
F5	22	1.1	31.00
F25	32	3.1	32.43
F50	45	5.4	33.15
F75	61	8.0	34.35
F95	89	10.0	35.05

Figure 7.17E. Catch distribution by depth, temperature and salinity of Sea raven.

857




Figure 7.17F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Sea raven.

858

7.18 Alligatorfish (Poisson-alligator atlantique) - species code 340 (category LF)

859

Scientific name: [Aspidophoroides monopterygius](#)

860




Figure 7.18A. Inverse distance weighted distribution of catch biomass (kg/tow) for Alligatorfish.

861




Figure 7.18B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Alligatorfish.




Figure 7.18C. Length frequency distribution in NAFO units 4X and 4VW for Alligatorfish.






Figure 7.18D. Average fish condition in NAFO units 4X and 4VW for Alligatorfish.



864



Freq	Depth	Temp	Sal
F5	32	1.5	31.00
F25	44	3.7	32.53
F50	57	6.1	33.28
F75	72	8.2	34.45
F95	102	10.0	35.10

Figure 7.18E. Catch distribution by depth, temperature and salinity of Alligatorfish.




Figure 7.18F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Alligatorfish.

866

7.19 Monkfish (Baudroie d'Amérique) - species code 400 (category LF)

867

Scientific name: [Lophius americanus](#)

868




Figure 7.19A. Inverse distance weighted distribution of catch biomass (kg/tow) for Monkfish.

869




Figure 7.19B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Monkfish.




Figure 7.19C. Length frequency distribution in NAFO units 4X and 4VW for Monkfish.







Figure 7.19D. Average fish condition in NAFO units 4X and 4VW for Monkfish.



872



Freq	Depth	Temp	Sal
F5	33	1.1	31.00
F25	60	2.9	32.43
F50	84	5.3	33.07
F75	110	7.8	34.31
F95	148	10.0	35.03

Figure 7.19E. Catch distribution by depth, temperature and salinity of Monkfish.

873




Figure 7.19F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Monkfish.

874

7.20 Ocean pout (Loquette d'Amérique) - species code 640 (category LF)

875

Scientific name: [Zoarces americanus](#)

876




Figure 7.20A. Inverse distance weighted distribution of catch biomass (kg/tow) for Ocean pout.

877




Figure 7.20B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Ocean pout.




Figure 7.20C. Length frequency distribution in NAFO units 4X and 4VW for Ocean pout.




Figure 7.20D. Average fish condition in NAFO units 4X and 4VW for Ocean pout.




Figure 7.20E. Catch distribution by depth, temperature and salinity of Ocean pout.




Figure 7.20F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Ocean pout.

882

7.21 Thorny skate (Raie épineuse) - species code 201 (category LF)

883

Scientific name: [Amblyraja radiata](#)

884




Figure 7.21A. Inverse distance weighted distribution of catch biomass (kg/tow) for Thorny skate.

885




Figure 7.21B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Thorny skate.




Figure 7.21C. Length frequency distribution in NAFO units 4X and 4VW for Thorny skate.






Figure 7.21D. Average fish condition in NAFO units 4X and 4VW for Thorny skate.



888



Freq	Depth	Temp	Sal
F5	29	1.3	31.00
F25	45	3.7	32.50
F50	67	6.2	33.30
F75	91	8.1	34.40
F95	148	9.9	35.03

Figure 7.21E. Catch distribution by depth, temperature and salinity of Thorny skate.




Figure 7.21F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Thorny skate.

890

7.22 Smooth skate (Raie lisse) - species code 202 (category LF)

891

Scientific name: [Malacoraja senta](#)

892




Figure 7.22A. Inverse distance weighted distribution of catch biomass (kg/tow) for Smooth skate.

893




Figure 7.22B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Smooth skate.




Figure 7.22C. Length frequency distribution in NAFO units 4X and 4VW for Smooth skate.






Figure 7.22D. Average fish condition in NAFO units 4X and 4VW for Smooth skate.



896



Freq	Depth	Temp	Sal
F5	33	1.2	31.00
F25	59	3.5	32.47
F50	80	5.9	33.23
F75	110	8.1	34.45
F95	171	10.0	35.06

Figure 7.22E. Catch distribution by depth, temperature and salinity of Smooth skate.

897




Figure 7.22F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Smooth skate.

898 **7.23 Winter skate (Raie tachetée) - species code 204 (category LF)**

899 Scientific name: [Leucoraja ocellata](#)

900




Figure 7.23A. Inverse distance weighted distribution of catch biomass (kg/tow) for Winter skate.

901




Figure 7.23B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Winter skate.




Figure 7.23C. Length frequency distribution in NAFO units 4X and 4VW for Winter skate.






Figure 7.23D. Average fish condition in NAFO units 4X and 4VW for Winter skate.



904



Freq	Depth	Temp	Sal
F5	21	1.1	31.00
F25	32	3.5	32.44
F50	45	5.9	33.19
F75	71	8.3	34.42
F95	140	10.0	35.03

Figure 7.23E. Catch distribution by depth, temperature and salinity of Winter skate.




Figure 7.23F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Winter skate.

906

7.24 Picked dogfish (Aiguillat commun) - species code 220 (category LF)

907

Scientific name: [Squalus acanthias](#)

908




Figure 7.24A. Inverse distance weighted distribution of catch biomass (kg/tow) for Picked dogfish.

909




Figure 7.24B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Picked dogfish.




Figure 7.24C. Length frequency distribution in NAFO units 4X and 4VW for Picked dogfish.






Figure 7.24D. Average fish condition in NAFO units 4X and 4VW for Picked dogfish.



912



Freq	Depth	Temp	Sal
F5	35	1.1	31.00
F25	60	2.8	32.47
F50	83	5.2	33.28
F75	108	7.7	34.37
F95	139	10.0	35.02

Figure 7.24E. Catch distribution by depth, temperature and salinity of Picked dogfish.




Figure 7.24F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Picked dogfish.

914 **7.25 Northern shortfin squid (*Encornet rouge nordique*) - species code 4511 (category**
 915 **LF)**

916 Scientific name: [Illex illecebrosus](#)

917




Figure 7.25A. Inverse distance weighted distribution of catch biomass (kg/tow) for Northern shortfin squid.

918




Figure 7.25B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Northern shortfin squid.




Figure 7.25C. Length frequency distribution in NAFO units 4X and 4VW for Northern shortfin squid.






Figure 7.25D. Average fish condition in NAFO units 4X and 4VW for Northern shortfin squid.



921



Freq	Depth	Temp	Sal
F5	31	1.3	31.00
F25	53	3.5	32.48
F50	71	5.9	33.28
F75	93	8.1	34.39
F95	139	10.0	35.05

Figure 7.25E. Catch distribution by depth, temperature and salinity of Northern shortfin squid.

922




Figure 7.25F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Northern shortfin squid.

923

7.26 Atlantic hagfish (*Myxine du nord*) - species code 241 (category LI)

924

Scientific name: *Myxine glutinosa*

925




Figure 7.26A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic hagfish.

926




Figure 7.26B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic hagfish.

927

7.27 Cusk (Brosme) - species code 15 (category LI)

928

Scientific name: [Brosme brosme](#)

929




Figure 7.27A. Inverse distance weighted distribution of catch biomass (kg/tow) for Cusk.

930




Figure 7.27B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Cusk.

931

7.28 Greenland halibut (Flétan noir) - species code 31 (category LI)

932

Scientific name: [Reinhardtius hippoglossoides](#)

933




Figure 7.28A. Inverse distance weighted distribution of catch biomass (kg/tow) for Greenland halibut.

934




Figure 7.28B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Greenland halibut.

935 **7.29 Gulf Stream flounder (Plie du Gulf Stream) - species code 44 (category LI)**

936 Scientific name: [Citharichthys arctifrons](#)




Figure 7.29A. Inverse distance weighted distribution of catch biomass (kg/tow) for Gulf Stream flounder.




Figure 7.29B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Gulf Stream flounder.

939

7.30 American shad (*Alose savoureuse*) - species code 61 (category LI)

940

Scientific name: [Alosa sapidissima](#)

941




Figure 7.30A. Inverse distance weighted distribution of catch biomass (kg/tow) for American shad.

942




Figure 7.30B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of American shad.

943

7.31 Alewife (Gaspareau) - species code 62 (category LI)

944

Scientific name: *Alosa pseudoharengus*

945




Figure 7.31A. Inverse distance weighted distribution of catch biomass (kg/tow) for Alewife.

946




Figure 7.31B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Alewife.

947

7.32 Capelin (Capelan) - species code 64 (category LI)

948

Scientific name: [Mallotus villosus](#)

949




Figure 7.32A. Inverse distance weighted distribution of catch biomass (kg/tow) for Capelin.

950




Figure 7.32B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Capelin.

951 **7.33 Atlantic mackerel (*Maquereau commun*) - species code 70 (category LI)**

952 Scientific name: *Scomber scombrus*

953




Figure 7.33A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic mackerel.

954




Figure 7.33B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic mackerel.

955 **7.34 Longfin hake (*Merluche à longues nageoires*) - species code 112 (category LI)**

956 Scientific name: [Phycis chesteri](#)

957




Figure 7.34A. Inverse distance weighted distribution of catch biomass (kg/tow) for Longfin hake.

958





Figure 7.34B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Longfin hake.


959 **7.35 Fourbeard rockling (Motelle à quatre barbillons) - species code 114 (category LI)**

960 Scientific name: [Enchelyopus cimbrius](#)



961

Figure 7.35A. Inverse distance weighted distribution of catch biomass (kg/tow) for Fourbeard rockling.



962

Figure 7.35B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Fourbeard rockling.

963 **7.36 Blackbelly rosefish (Sébaste chèvre) - species code 123 (category LI)**

964 Scientific name: [Helicolenus dactylopterus](#)




Figure 7.36A. Inverse distance weighted distribution of catch biomass (kg/tow) for Blackbelly rosefish.




Figure 7.36B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Blackbelly rosefish.

967

7.37 Greater argentine (Grande argentine) - species code 160 (category LI)

968

Scientific name: [Argentina silus](#)

969




Figure 7.37A. Inverse distance weighted distribution of catch biomass (kg/tow) for Greater argentine.

970




Figure 7.37B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Greater argentine.

971 **7.38 Arctic hookear sculpin (*Hameçon neigeux*) - species code 306 (category LI)**

972 Scientific name: [Artediellus uncinatus](#)




Figure 7.38A. Inverse distance weighted distribution of catch biomass (kg/tow) for Arctic hookear sculpin.




Figure 7.38B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Arctic hookear sculpin.

975

7.39 Atlantic poacher (*Agone atlantique*) - species code 350 (category LI)

976

Scientific name: [Leptagonus decagonus](#)

977




Figure 7.39A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic poacher.

978




Figure 7.39B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic poacher.

979 **7.40 Marlin-spike grenadier (Grenadier du Grand Banc) - species code 410 (category**
 980 **LI)**

981 Scientific name: [Nezumia bairdii](#)




Figure 7.40A. Inverse distance weighted distribution of catch biomass (kg/tow) for Marlin-spike grenadier.




Figure 7.40B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Marlin-spike grenadier.

984

7.41 Lumpfish (Lompe) - species code 501 (category LI)

985

Scientific name: [Cyclopterus lumpus](#)

986




Figure 7.41A. Inverse distance weighted distribution of catch biomass (kg/tow) for Lumpfish.

987




Figure 7.41B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Lumpfish.

988 **7.42 Atlantic spiny lumpsucker (Petite poule de mer atlantique) - species code 502**
 989 (**category LI**)

990 Scientific name: [Eumicrotremus spinosus](#)

991




Figure 7.42A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic spiny lumpsucker.

992




Figure 7.42B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic spiny lumpsucker.

993

7.43 Sand lance (Lançon) - species code 610 (category LI)

994

Scientific name: [Ammodytes dubius](#)

995




Figure 7.43A. Inverse distance weighted distribution of catch biomass (kg/tow) for Sand lance.

996




Figure 7.43B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Sand lance.

997

7.44 Snakeblenny (Lompénie-serpent) - species code 622 (category LI)

998

Scientific name: [Lumpenus lampretaeformis](#)

999




Figure 7.44A. Inverse distance weighted distribution of catch biomass (kg/tow) for Snakeblenny.

1000




Figure 7.44B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Snakeblenny.

1001

7.45 Daubed shanny (Lompénie tachetée) - species code 623 (category LI)

1002

Scientific name: [Leptoclinus maculatus](#)

1003




Figure 7.45A. Inverse distance weighted distribution of catch biomass (kg/tow) for Daubed shanny.

1004




Figure 7.45B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Daubed shanny.

1005

7.46 Vahl's eelpout (*Lycodes vahlii*) - species code 647 (category LI)

1006

Scientific name: [Lycodes vahlii](#)

1007




Figure 7.46A. Inverse distance weighted distribution of catch biomass (kg/tow) for Vahl's eelpout.

1008




Figure 7.46B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Vahl's eelpout.

1009

7.47 Atlantic butterfish (*Stromaté fossette*) - species code 701 (category LI)

1010

Scientific name: [Peprilus triacanthus](#)

1011




Figure 7.47A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic butterfish.

1012




Figure 7.47B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic butterfish.


1013

7.48 Atlantic hookear sculpin (*Hameçon atlantique*) - species code 880 (category LI)

1014

Scientific name: [Artediellus atlanticus](#)

1015



1016

Figure 7.48A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic hookear sculpin.




Figure 7.48B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic hookear sculpin.

1017

7.49 Barndoor skate (Grande raie) - species code 200 (category LI)

1018

Scientific name: [Dipturus laevis](#)

1019




Figure 7.49A. Inverse distance weighted distribution of catch biomass (kg/tow) for Barndoor skate.

1020




Figure 7.49B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Barndoor skate.

1021

7.50 Little skate (Raie hérisson) - species code 203 (category LI)

1022

Scientific name: [Leucoraja erinacea](#)

1023




Figure 7.50A. Inverse distance weighted distribution of catch biomass (kg/tow) for Little skate.

1024




Figure 7.50B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Little skate.

1025

7.51 Northern prawn (Crevette nordique) - species code 2211 (category SF)

1026

Scientific name: [Pandalus borealis](#)

1027




Figure 7.51A. Inverse distance weighted distribution of catch biomass (kg/tow) for Northern prawn.

1028




Figure 7.51B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Northern prawn.

1029

7.52 Jonah crab (*Tourteau jona*) - species code 2511 (category SF)

1030

Scientific name: [Cancer borealis](#)

1031




Figure 7.52A. Inverse distance weighted distribution of catch biomass (kg/tow) for Jonah crab.

1032




Figure 7.52B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Jonah crab.

1033

7.53 Atlantic rock crab (Tourteau poïnclos) - species code 2513 (category SF)

1034

Scientific name: [Cancer irroratus](#)

1035




Figure 7.53A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic rock crab.

1036




Figure 7.53B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic rock crab.

1037

7.54 Arctic lyre crab (*Crabe Hyas coarctatus*) - species code 2521 (category SF)

1038

Scientific name: [Hyas coarctatus](#)

1039




Figure 7.54A. Inverse distance weighted distribution of catch biomass (kg/tow) for Arctic lyre crab.

1040




Figure 7.54B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Arctic lyre crab.

1041

7.55 Atlantic king crab (Crabe épineux du nord) - species code 2523 (category SF)

1042

Scientific name: [Lithodes maja](#)

1043




Figure 7.55A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic king crab.

1044




Figure 7.55B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic king crab.

1045

7.56 Queen crab (Crabe des neiges) - species code 2526 (category SF)

1046

Scientific name: [Chionoecetes opilio](#)

1047




Figure 7.56A. Inverse distance weighted distribution of catch biomass (kg/tow) for Queen crab.

1048




Figure 7.56B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Queen crab.

1049

7.57 Great spider crab (Crabe lyre araignée) - species code 2527 (category SF)

1050

Scientific name: [Hyas araneus](#)

1051




Figure 7.57A. Inverse distance weighted distribution of catch biomass (kg/tow) for Great spider crab.

1052




Figure 7.57B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Great spider crab.

1053

7.58 American lobster (Homard américain) - species code 2550 (category SF)

1054

Scientific name: [Homarus americanus](#)

1055




Figure 7.58A. Inverse distance weighted distribution of catch biomass (kg/tow) for American lobster.

1056




Figure 7.58B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of American lobster.

1057

7.59 Sea lamprey (*Lamproie marine*) - species code 240 (category LR)

1058

Scientific name: [Petromyzon marinus](#)

1059




Figure 7.59A. Catch distribution for Sea lamprey.

1060 **7.60 Atlantic tomcod (*Poulamon atlantique*) - species code 17 (category LR)**

1061 Scientific name: [Micogadus tomcod](#)




Figure 7.60A. Catch distribution for Atlantic tomcod.

1063

7.61 Offshore silver hake (Merlu argenté du large) - species code 19 (category LR)

1064

Scientific name: [Merluccius albidus](#)

1065




Figure 7.61A. Catch distribution for Offshore silver hake.

1066

7.62 Spotted wolffish (Loup tacheté) - species code 51 (category LR)

1067

Scientific name: [Anarhichas minor](#)

1068




Figure 7.62A. Catch distribution for Spotted wolffish.

1069

7.63 Northern wolffish (Loup à tête large) - species code 52 (category LR)

1070

Scientific name: [Anarhichas denticulatus](#)

1071




Figure 7.63A. Catch distribution for Northern wolffish.

1072

7.64 Rainbow smelt (Éperlan arc-en-ciel) - species code 63 (category LR)

1073

Scientific name: [Osmerus mordax](#)

1074




Figure 7.64A. Catch distribution for Rainbow smelt.

1075

7.65 Cunner (Tanche-tautogue) - species code 122 (category LR)

1076

Scientific name: [Tautogolabrus adspersus](#)

1077




Figure 7.65A. Catch distribution for Cunner.

1078

7.66 Fourspot flounder (Cardeau à quatre ocelles) - species code 142 (category LR)

1079

Scientific name: [Hippoglossina oblonga](#)

1080




Figure 7.66A. Catch distribution for Fourspot flounder.

1081

7.67 Windowpane flounder (Turbot de sable) - species code 143 (category LR)

1082

Scientific name: [Scophthalmus aquosus](#)

1083




Figure 7.67A. Catch distribution for Windowpane flounder.

1084

7.68 Longnose greeneye (Oeil-vert à long nez) - species code 149 (category LR)

1085

Scientific name: [Parasudis triculenta](#)

1086




Figure 7.68A. Catch distribution for Longnose greeneye.

1087

7.69 Lanternfishes (Poissons-lanternes) - species code 150 (category LR)

1088

Scientific name: [Myctophidae](#)

1089




Figure 7.69A. Catch distribution for Lanternfishes.

1090 **7.70 Shortnose greeneye (Éperlan du large) - species code 156 (category LR)**

1091 Scientific name: [Chlorophthalmus agassizi](#)




Figure 7.70A. Catch distribution for Shortnose greeneye.

1093

7.71 Silvery lightfish (Brossé améthyste) - species code 158 (category LR)

1094

Scientific name: [Maurolicus muelleri](#)

1095




Figure 7.71A. Catch distribution for Silvery lightfish.

1096

7.72 Boa dragonfish (Dragon-boa) - species code 159 (category LR)

1097

Scientific name: [Stomias boa](#)

1098




Figure 7.72A. Catch distribution for Boa dragonfish.

1100

7.73 Shorthorn sculpin (*Chabotisca* à épines courtes) - species code 301 (category LR)

1101

Scientific name: *Myoxocephalus scorpius*




Figure 7.73A. Catch distribution for Shorthorn sculpin.

1103

7.74 Grubby (Chabosseau bronzé) - species code 303 (category LR)

1104

Scientific name: [Myoxocephalus aenaeus](#)

1105




Figure 7.74A. Catch distribution for Grubby.

1106

7.75 Polar sculpin (Cotte polaire) - species code 307 (category LR)

1107

Scientific name: [Cottunculus microps](#)

1108




Figure 7.75A. Catch distribution for Polar sculpin.

1109

7.76 Spatulate sculpin (Icèle spatulée) - species code 314 (category LR)

1110

Scientific name: *Icelus spatula*

1111




Figure 7.76A. Catch distribution for Spatulate sculpin.

1112

7.77 Arctic alligatorfish (Poisson-alligator arctique) - species code 341 (category LR)

1113

Scientific name: [Ulcina olrikii](#)

1114




Figure 7.77A. Catch distribution for Arctic alligatorfish.


1115

7.78 Alligatorfishes (Poissons-alligator) - species code 351 (category LR)

1116

Scientific name: [Agonidae](#)

1117



1118

7.79 Roughnose grenadier (Grenadier-scie) - species code 412 (category LR)

1119

Scientific name: [Trachyrincus murrayi](#)

1120




Figure 7.79A. Catch distribution for Roughnose grenadier.

1121 **7.80 Roundnose grenadier (Grenadier de roche) - species code 414 (category LR)**

1122 Scientific name: [Coryphaenoides rupestris](#)




Figure 7.80A. Catch distribution for Roundnose grenadier.

1124

7.81 Atlantic seasnail (*Limace atlantique*) - species code 503 (category LR)

1125

Scientific name: [Liparis atlanticus](#)

1126




Figure 7.81A. Catch distribution for Atlantic seasnail.

1127 **7.82 Gelatinous snailfish (Limace gélatineuse) - species code 505 (category LR)**

1128 Scientific name: [Liparis fabricii](#)




Figure 7.82A. Catch distribution for Gelatinous snailfish.

1130 **7.83 Variegated snailfish (*Limace marbée*) - species code 512 (category LR)**

1131 Scientific name: [Liparis gibbus](#)




Figure 7.83A. Catch distribution for Variegated snailfish.

1133

7.84 Sea tadpole (Petite limace de mer) - species code 520 (category LR)

1134

Scientific name: [Careproctus reinhardtii](#)

1135




Figure 7.84A. Catch distribution for Sea tadpole.

1136

7.85 Wolf eelpout (*Lycodes à tête longue*) - species code 603 (category LR)

1137

Scientific name: *Lycenchelys verrillii*

1138




Figure 7.85A. Catch distribution for Wolf eelpout.

1139

7.86 Slender snipe eel (*Avocette ruban*) - species code 604 (category LR)

1140

Scientific name: [Nemichthys scolopaceus](#)

1141




Figure 7.86A. Catch distribution for Slender snipe eel.

1142

7.87 Newfoundland eelpout (*Lycodes* du Labrador) - species code 619 (category LR)

1143

Scientific name: [Lycodes terraenovae](#)

1144




Figure 7.87A. Catch distribution for Newfoundland eelpout.

1145 **7.88 Newfoundland eelpout (*Lycodes* du Labrador) - species code 620 (category LR)**

1146 Scientific name: [Lycodes lavalaei](#)




Figure 7.88A. Catch distribution for Newfoundland eelpout.

1148

7.89 Rock gunnel (Sigouine de roche) - species code 621 (category LR)

1149

Scientific name: [Pholis gunnellus](#)

1150




Figure 7.89A. Catch distribution for Rock gunnel.

1151

7.90 Radiated shanny (*Ulvaria deux-lignes*) - species code 625 (category LR)

1152

Scientific name: [Ulvaria subbifurcata](#)

1153




Figure 7.90A. Catch distribution for Radiated shanny.

1154

7.91 Fourline snakeblenny (Quatre-lignes atlantique) - species code 626 (category LR)

1155

Scientific name: [Eumesogrammus praecisus](#)

1156




Figure 7.91A. Catch distribution for Fourline snakeblenny.

1157

7.92 Wrymouth (Terrassier tacheté) - species code 630 (category LR)

1158

Scientific name: [Cryptacanthodes maculatus](#)

1159




Figure 7.92A. Catch distribution for Wrymouth.

1160 7.93 Spotfin dragonet (Dragonnet tacheté) - species code 637 (category LR)

1161 Scientific name: [Foetorepus agassizii](#)




Figure 7.93A. Catch distribution for Spotfin dragonet.

1163

7.94 Arctic eelpout (*Lycodes arctique*) - species code 641 (category LR)

1164

Scientific name: [Lycodes reticulatus](#)

1165




Figure 7.94A. Catch distribution for Arctic eelpout.

1166

7.95 Atlantic soft pout (*Molasse atlantique*) - species code 646 (category LR)

1167

Scientific name: [Melanostigma atlanticum](#)

1168




Figure 7.95A. Catch distribution for Atlantic soft pout.

1169

7.96 Silvery John dory (Saint Pierre argenté) - species code 704 (category LR)

1170

Scientific name: [Zenopsis conchifer](#)

1171




Figure 7.96A. Catch distribution for Silvery John dory.

1172

7.97 White barracudina (*Lussion blanc*) - species code 712 (category LR)

1173

Scientific name: [Arctozenus risso](#)

1174




Figure 7.97A. Catch distribution for White barracudina.

1175

7.98 Atlantic saury (*Balaou atlantique*) - species code 720 (category LR)

1176

Scientific name: [Scomberesox saurus](#)

1177




Figure 7.98A. Catch distribution for Atlantic saury.

1178

7.99 Hatchetfishes (Haches d'argent) - species code 741 (category LR)

1179

Scientific name: [Sternopychidae](#)

1180




Figure 7.99A. Catch distribution for Hatchetfishes.

1181

7.100 Atlantic batfish (*Malthe atlantique*) - species code 742 (category LR)

1182

Scientific name: [Dibranchus atlanticus](#)

1183




Figure 7.100A. Catch distribution for Atlantic batfish.

1184

7.101 Spottedfin tonguefish (Langue fil noir) - species code 816 (category LR)

1185

Scientific name: [Symphurus diomedeanus](#)

1186




Figure 7.101A. Catch distribution for Spottedfin tonguefish.

1187

7.102 Black dogfish (Aiguillat noir) - species code 221 (category LR)

1188

Scientific name: [Centroscyllium fabricii](#)

1189




Figure 7.102A. Catch distribution for Black dogfish.

1190

7.103 Longfin inshore squid (*Calmar totam*) - species code 4512 (category LR)

1191

Scientific name: [Doryteuthis pealeii](#)

1192




Figure 7.103A. Catch distribution for Longfin inshore squid.

1193

7.104 Red deepsea crab (Crabe rouge) - species code 2532 (category SR)

1194

Scientific name: [Chaceon quinquedens](#)

1195




Figure 7.104A. Catch distribution for Red deepsea crab.

INDEX

1196	Agone atlantique, 121	1242	Atlantic hookear sculpin, 130
1197	Agonidae, 160	1243	Atlantic king crab, 137
1198	Agonidae, 160	1244	Atlantic mackerel, 115
1199	Aspidophoroides monopterygius, 84	1245	Atlantic poacher, 121
1200	Leptagonus decagonus, 121	1246	Atlantic redfishes, 51
1201	Ulcina olrikii, 159	1247	Atlantic rock crab, 135
1202	Aiglefin, 36	1248	Atlantic saury, 180
1203	Aiguillat commun, 102	1249	Atlantic seasnail, 163
1204	Aiguillat noir, 184	1250	Atlantic soft pout, 177
1205	Alewife, 113	1251	Atlantic spiny lumpsucker, 124
1206	Alligatorfish, 84	1252	Atlantic tomcod, 142
1207	Alligatorfishes, 160	1253	Atlantic wolffish, 69
1208	Alosa pseudoharengus, 113	1254	Avocette ruban, 168
1209	Alosa sapidissima, 112	1255	Balaou atlantique, 180
1210	Alose savoureuse, 112	1256	Barndoor skate, 131
1211	Amblyraja radiata, 93	1257	Baudroie d'Amérique, 87
1212	American lobster, 140	1258	Black dogfish, 184
1213	American plaice, 57	1259	Blackbelly rosefish, 118
1214	American shad, 112	1260	Boa dragonfish, 154
1215	Ammodytes dubius, 125	1261	Brosme, 109
1216	Ammodytidae	1262	Brosme brosme, 109
1217	Ammodytes dubius, 125	1263	Brossé améthyste, 153
1218	Anarhichadidae	1264	Callionymidae
1219	Anarhichas denticulatus, 145	1265	Foetorepus agassizii, 175
1220	Anarhichas lupus, 69	1266	Calmar totam, 185
1221	Anarhichas minor, 144	1267	Cancer borealis, 134
1222	Anarhichas denticulatus, 145	1268	Cancer irroratus, 135
1223	Anarhichas lupus, 69	1269	Cancridae
1224	Anarhichas minor, 144	1270	Cancer borealis, 134
1225	Arctic alligatorfish, 159	1271	Cancer irroratus, 135
1226	Arctic eelpout, 176	1272	Capelan, 114
1227	Arctic hookear sculpin, 120	1273	Capelin, 114
1228	Arctic lyre crab, 136	1274	Cardeau à quatre ocelles, 148
1229	Arctozenus risso, 179	1275	Careproctus reinhardti, 166
1230	Argentina silus, 119	1276	Centroscyllium fabricii, 184
1231	Argentinidae	1277	Chabosseau bronzé, 156
1232	Argentina silus, 119	1278	Chabosseau à dix-huit épines, 75
1233	Artediellus atlanticus, 130	1279	Chabosseau à épines courtes, 155
1234	Artediellus uncinatus, 120	1280	Chaceon quinquedens, 186
1235	Aspidophoroides monopterygius, 84	1281	Chionoecetes opilio, 138
1236	Atlantic batfish, 182	1282	Chlorophthalmidae
1237	Atlantic butterfish, 129	1283	Chlorophthalmus agassizi, 152
1238	Atlantic cod, 33	1284	Parasudis truculenta, 150
1239	Atlantic hagfish, 108	1285	Chlorophthalmus agassizi, 152
1240	Atlantic halibut, 54		
1241	Atlantic herring, 72		

1286	Citharichthys arctifrons, 111	1332	Faux-trigle armé, 78
1287	Clupea harengus, 72	1333	Flétan de l'Atlantique, 54
1288	Clupeidae	1334	Flétan noir, 110
1289	Alosa pseudoharengus, 113	1335	Foetorepus agassizii, 175
1290	Alosa sapidissima, 112	1336	Fourbeard rockling, 117
1291	Clupea harengus, 72	1337	Fourline snakeblenny, 173
1292	Coryphaenoides rupestris, 162	1338	Fourspot flounder, 148
1293	Cotte polaire, 157		
1294	Cottidae	1339	Gadidae
1295	Artediellus atlanticus, 130	1340	Gadus morhua, 33
1296	Artediellus uncinatus, 120	1341	Melanogrammus aeglefinus, 36
1297	Icelus spatula, 158	1342	Microgadus tomcod, 142
1298	Myoxocephalus aenaeus, 156	1343	Pollachius virens, 48
1299	Myoxocephalus octodecemspinosis, 75 ¹⁴	1344	Gadus morhua, 33
1300	Myoxocephalus scorpius, 155	1345	Gaspareau, 113
1301	Triglops murrayi, 78	1346	Gelatinous snailfish, 164
1302	Cottunculus microps, 157	1347	Geryonidae
1303	Crabe des neiges, 138	1348	Chaceon quinquedens, 186
1304	Crabe Hyas coarctatus, 136	1349	Glyptocephalus cynoglossus, 60
1305	Crabe lyre araignée, 139	1350	Goberge, 48
1306	Crabe rouge, 186	1351	Grande argentine, 119
1307	Crabe épineux du nord, 137	1352	Grande raie, 131
1308	Crevette nordique, 133	1353	Great spider crab, 139
1309	Cryptacanthodes maculatus, 174	1354	Greater argentine, 119
1310	Cryptacanthodidae	1355	Greenland halibut, 110
1311	Cryptacanthodes maculatus, 174	1356	Grenadier de roche, 162
1312	Cunner, 147	1357	Grenadier du Grand Banc, 122
1313	Cusk, 109	1358	Grenadier-scie, 161
1314	Cyclopteridae	1359	Grubby, 156
1315	Cyclopterus lumpus, 123	1360	Gulf Stream flounder, 111
1316	Eumicrotremus spinosus, 124		
1317	Cyclopterus lumpus, 123	1361	Haches d'argent, 181
1318	Cynoglossidae	1362	Haddock, 36
1319	Symphurus diomedeanus, 183	1363	Hameçon atlantique, 130
1320	Daubed shanny, 127	1364	Hameçon neigeux, 120
1321	Dibranchus atlanticus, 182	1365	Hareng de l'Atlantique, 72
1322	Dipturus laevis, 131	1366	Hatchetfishes, 181
1323	Doryteuthis pealeii, 185	1367	Helicolenus dactylopterus, 118
1324	Dragon-boa, 154	1368	Hemitripteridae
1325	Dragonnet tacheté, 175	1369	Hemitripterus americanus, 81
1326	Enchelyopus cimbrius, 117	1370	Hemitripterus americanus, 81
1327	Encornet rouge nordique, 105	1371	Hippoglossina oblonga, 148
1328	Etmopteridae	1372	Hippoglossoides platessoides, 57
1329	Centroscyllium fabricii, 184	1373	Hippoglossus hippoglossus, 54
1330	Eumesogrammus praecisus, 173	1374	Homard américain, 140
1331	Eumicrotremus spinosus, 124	1375	Homarus americanus, 140
		1376	Hyas araneus, 139
		1377	Hyas coarctatus, 136

1378	Hémitriptère atlantique, 81	1423	Loquette d'Amérique, 90
1379	<i>Icelus spatula</i> , 158	1424	Lotidae
1380	lcèle spatulée, 158	1425	<i>Brosme brosme</i> , 109
1381	<i>Illex illecebrosus</i> , 105	1426	<i>Enchelyopus cimbrius</i> , 117
1382	Jonah crab, 134	1427	Loup atlantique, 69
1383	Labridae	1428	Loup tacheté, 144
1384	<i>Tautogolabrus adspersus</i> , 147	1429	Loup à tête large, 145
1385	Lamproie marine, 141	1430	<i>Lumpenus lampretaeformis</i> , 126
1386	Langue fil noir, 183	1431	Lumpfish, 123
1387	Lanternfishes, 151	1432	Lussion blanc, 179
1388	Lançon, 125	1433	<i>Lycenchelys verrillii</i> , 167
1389	<i>Leptagonus decagonus</i> , 121	1434	Lycode arctique, 176
1390	<i>Leptoclinus maculatus</i> , 127	1435	Lycode du Labrador, 169, 170
1391	<i>Leucoraja erinacea</i> , 132	1436	Lycode à carreaux, 128
1392	<i>Leucoraja ocellata</i> , 99	1437	Lycode à tête longue, 167
1393	Limace atlantique, 163	1438	<i>Lycodes lavalaei</i> , 170
1394	Limace gélatineuse, 164	1439	<i>Lycodes reticulatus</i> , 176
1395	Limace marbée, 165	1440	<i>Lycodes terraenovae</i> , 169
1396	<i>Limanda ferruginea</i> , 63	1441	<i>Lycodes vahlii</i> , 128
1397	Limande à queue jaune, 63	1442	Macrouridae
1398	Limande-plie rouge, 66	1443	<i>Coryphaenoides rupestris</i> , 162
1399	Liparidae	1444	<i>Nezumia bairdii</i> , 122
1400	<i>Careproctus reinhardtii</i> , 166	1445	<i>Trachyrincus murrayi</i> , 161
1401	<i>Liparis atlanticus</i> , 163	1446	Malacoraja senta, 96
1402	<i>Liparis fabricii</i> , 164	1447	Mallotus villosus, 114
1403	<i>Liparis gibbus</i> , 165	1448	Malthe atlantique, 182
1404	<i>Liparis atlanticus</i> , 163	1449	Maquereau commun, 115
1405	<i>Liparis fabricii</i> , 164	1450	Marlin-spike grenadier, 122
1406	<i>Liparis gibbus</i> , 165	1451	Maurolicus muelleri, 153
1407	<i>Lithodes maja</i> , 137	1452	<i>Melanogrammus aeglefinus</i> , 36
1408	Lithodidae	1453	<i>Melanostigma atlanticum</i> , 177
1409	<i>Lithodes maja</i> , 137	1454	Merlu argenté, 45
1410	Little skate, 132	1455	Merlu argenté du large, 143
1411	Loliginidae	1456	Merlucciidae
1412	<i>Doryteuthis pealeii</i> , 185	1457	<i>Merluccius albidus</i> , 143
1413	Lompe, 123	1458	<i>Merluccius bilinearis</i> , 45
1414	Lompénie tachetée, 127	1459	Merluccius albidus, 143
1415	Lompénie-serpent, 126	1460	Merluccius bilinearis, 45
1416	Longfin hake, 116	1461	Merluche blanche, 39
1417	Longfin inshore squid, 185	1462	Merluche à longues nageoires, 116
1418	Longhorn sculpin, 75	1463	Merluche écureuil, 42
1419	Longnose greeneye, 150	1464	<i>Microgadus tomcod</i> , 142
1420	Lophiidae	1465	Molasse atlantique, 177
1421	<i>Lophius americanus</i> , 87	1466	Monkfish, 87
1422	<i>Lophius americanus</i> , 87	1467	Morue franche, 33
		1468	Motelle à quatre barbillons, 117
		1469	Moustache sculpin, 78

1470	Myctophidae, 151	1515	Petite poule de mer atlantique, 124
1471	Myctophidae, 151	1516	Petromyzon marinus, 141
1472	Myoxocephalus aenaeus, 156	1517	Petromyzontidae
1473	Myoxocephalus octodecemspinulosus, 75	1518	Petromyzon marinus, 141
1474	Myoxocephalus scorpius, 155	1519	Pholidae
1475	Myxine du nord, 108	1520	Pholis gunnellus, 171
1476	Myxine glutinosa, 108	1521	Pholis gunnellus, 171
1477	Myxinidae	1522	Phycidae
1478	Myxine glutinosa, 108	1523	Phycis chesteri, 116
1479	Nemichthyidae	1524	Urophycis chuss, 42
1480	Nemichthys scolopaceus, 168	1525	Urophycis tenuis, 39
1481	Nemichthys scolopaceus, 168	1526	Phycis chesteri, 116
1482	Nephropidae	1527	Picked dogfish, 102
1483	Homarus americanus, 140	1528	Pleuronectidae
1484	Newfoundland eelpout, 169, 170	1529	Glyptocephalus cynoglossus, 60
1485	Nezumia bairdii, 122	1530	Hippoglossoides platessoides, 57
1486	Northern prawn, 133	1531	Hippoglossus hippoglossus, 54
1487	Northern shortfin squid, 105	1532	Limanda ferruginea, 63
1488	Northern wolffish, 145	1533	Pseudopleuronectes americanus, 66
1489	Ocean pout, 90	1534	Reinhardtius hippoglossoides, 110
1490	Oeil-vert à long nez, 150	1535	Plie canadienne, 57
1491	Offshore silver hake, 143	1536	Plie du Gulf Stream, 111
1492	Ogcocephalidae	1537	Plie grise, 60
1493	Dibranchus atlanticus, 182	1538	Poisson-alligator arctique, 159
1494	Ommastrephidae	1539	Poisson-alligator atlantique, 84
1495	Illex illecebrosus, 105	1540	Poissons-alligator, 160
1496	Oregoniidae	1541	Poissons-lanternes, 151
1497	Chionoecetes opilio, 138	1542	Polar sculpin, 157
1498	Hyas araneus, 139	1543	Pollachius virens, 48
1499	Hyas coarctatus, 136	1544	Pollock, 48
1500	Osmeridae	1545	Poulamon atlantique, 142
1501	Mallotus villosus, 114	1546	Pseudopleuronectes americanus, 66
1502	Osmerus mordax, 146	1547	Psychrolutidae
1503	Osmerus mordax, 146	1548	Cottunculus microps, 157
1504	Pandalidae	1549	Quatre-lignes atlantique, 173
1505	Pandalus borealis, 133	1550	Queen crab, 138
1506	Pandalus borealis, 133	1551	Radiated shanny, 172
1507	Paralepididae	1552	Raie hérisson, 132
1508	Arctozenus risso, 179	1553	Raie lisse, 96
1509	Paralichthyidae	1554	Raie tachetée, 99
1510	Citharichthys arctifrons, 111	1555	Raie épineuse, 93
1511	Hippoglossina oblonga, 148	1556	Rainbow smelt, 146
1512	Parasudis triculenta, 150	1557	Rajidae
1513	Peprilus triacanthus, 129	1558	Amblyraja radiata, 93
1514	Petite limace de mer, 166	1559	Dipturus laevis, 131
		1560	Leucoraja erinacea, 132

1561	Leucoraja ocellata, 99	1608	Leptoclinus maculatus, 127
1562	Malacoraja senta, 96	1609	Lumpenus lampretaeformis, 126
1563	Red deepsea crab, 186	1610	Ulvaria subbifurcata, 172
1564	Red hake, 42	1611	Stomias boa, 154
1565	Reinhardtius hippoglossoides, 110	1612	Stomiidae
1566	Rock gunnel, 171	1613	Stomias boa, 154
1567	Roughnose grenadier, 161	1614	Stromateidae
1568	Roundnose grenadier, 162	1615	Peprilus triacanthus, 129
1569	Saint Pierre argenté, 178	1616	Stromaté fossette, 129
1570	Sand lance, 125	1617	Syphurus diomedeanus, 183
1571	Scomber scombrus, 115	1618	Sébaste chèvre, 118
1572	Scomberesocidae	1619	Sébastes de l'Atlantique, 51
1573	Scomberesox saurus, 180	1620	Tanche-tautogue, 147
1574	Scomberesox saurus, 180	1621	Tautogolabrus adspersus, 147
1575	Scombridae	1622	Terrassier tacheté, 174
1576	Scomber scombrus, 115	1623	Thorny skate, 93
1577	Scophthalmidae	1624	Tourteau jona, 134
1578	Scophthalmus aquosus, 149	1625	Tourteau poïnclos, 135
1579	Scophthalmus aquosus, 149	1626	Trachyrincus murrayi, 161
1580	Sea lamprey, 141	1627	Triglops murrayi, 78
1581	Sea raven, 81	1628	Turbot de sable, 149
1582	Sea tadpole, 166		
1583	Sebastes, 51	1629	Ulcina olrikii, 159
1584	Sebastidae	1630	Ulvaire deux-lignes, 172
1585	Helicolenus dactylopterus, 118	1631	Ulvaria subbifurcata, 172
1586	Sebastes, 51	1632	Urophycis chuss, 42
1587	Shorthorn sculpin, 155	1633	Urophycis tenuis, 39
1588	Shortnose greeneye, 152		Vahl's eelpout, 128
1589	Sigouine de roche, 171	1634	Variegated snailfish, 165
1590	Silver hake, 45	1635	
1591	Silvery John dory, 178	1636	White barracudina, 179
1592	Silvery lightfish, 153	1637	White hake, 39
1593	Slender snipe eel, 168	1638	Windowpane flounder, 149
1594	Smooth skate, 96	1639	Winter flounder, 66
1595	Snakeblenny, 126	1640	Winter skate, 99
1596	Spatulate sculpin, 158	1641	Witch flounder, 60
1597	Spotfin dragonet, 175	1642	Wolf eelpout, 167
1598	Spotted wolffish, 144	1643	Wrymouth, 174
1599	Spottedfin tonguefish, 183		
1600	Squalidae	1644	Yellowtail flounder, 63
1601	Squalus acanthias, 102		
1602	Squalus acanthias, 102	1645	Zeidae
1603	Sternopychidae, 181	1646	Zenopsis conchifer, 178
1604	Maurolicus muelleri, 153	1647	Zenopsis conchifer, 178
1605	Sternopychidae, 181	1648	Zoarces americanus, 90
1606	Stichaeidae	1649	Zoarcidae
1607	Eumesogrammus praecisus, 173	1650	Lycenchelys verrillii, 167

- 1651 Lycodes lavalaei, 170
 - 1652 Lycodes reticulatus, 176
 - 1653 Lycodes terraenovae, 169
 - 1654 Lycodes vahlii, 128
 - 1655 Melanostigma atlanticum, 177
 - 1656 Zoarces americanus, 90
-
- 1657 Éperlan arc-en-ciel, 146
 - 1658 Éperlan du large, 152