

Marine Fish and Invertebrate Atlas: Summarizing Geographic Distribution and Population Indices in the Scotian Shelf and Bay of Fundy (1970-2020)

Daniel Ricard, Jamie Emberley, Catalina Gomez and Catriona Regnier-McKellar

Science Branch
Gulf Region
Fisheries and Oceans Canada
Moncton, New Brunswick, E1C 5K4, Canada

Science Branch
Maritimes Region
Fisheries and Oceans Canada
Dartmouth, Nova Scotia, B2Y 4A2, Canada

2021

Canadian Technical Report of
Fisheries and Aquatic Sciences ####



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Canada

Canadian Technical Report of Fisheries and Aquatic Sciences

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Rapport technique canadien des sciences halieutiques et aquatiques

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Canadian Technical Report of
Fisheries and Aquatic Sciences nnn

2021

6 MARINE FISH AND INVERTEBRATE ATLAS: SUMMARIZING GEOGRAPHIC DISTRIBUTION,
7 POPULATION INDICES AND ENVIRONMENTAL PREFERENCES IN THE SCOTIAN SHELF
8 AND BAY OF FUNDY (1970-2020)

by

10 Daniel Ricard¹ Jamie Emberley² Catalina Gomez² Catriona Regnier-McKellar²

¹Science Branch
Gulf Region
Fisheries and Oceans Canada
Moncton, New Brunswick, E1C 5K4, Canada
²Science Branch
Maritimes Region
Fisheries and Oceans Canada
Dartmouth, Nova Scotia, B2Y 4A2, Canada

¹⁹ © Her Majesty the Queen in Right of Canada, 2021
²⁰ Cat. No. Fs97-6/nnnE-PDF ISBN ISSN 1488-5379

²¹ Correct citation for this publication:

²² Ricard, D., Emberley, J., Gomez, C. and Regnier-McKellar, C. 2021. Marine Fish and Invertebrate Atlas: Summarizing Geographic Distribution, Population Indices and Environmental Preferences in the Scotian Shelf and Bay of Fundy (1970-2020). Can. Tech. Rep. Fish. Aquat. Sci. nnn: x + 188 p.

CONTENTS

27	ABSTRACT	ix
28	RÉSUMÉ	x
29	1 Introduction	1
30	2 Methods	2
31	2.1 Survey Description	2
32	2.2 Sampling Design	3
33	2.3 Taxonomic Levels	4
34	2.4 Analyses	13
35	2.4.1 Geographic distribution of catches	13
36	2.4.2 Abundance and biomass indices	13
37	2.4.3 Distribution indices	13
38	2.4.4 Length frequencies	13
39	2.4.5 Length-weight relationship and condition factor	14
40	2.4.6 Depth, temperature and salinity distribution of catches	14
41	2.4.7 Density-dependent habitat selection	14
42	3 Results	15
43	3.1 Description of Figures	15
44	3.1.1 Type A	15
45	3.1.2 Type B	15
46	3.1.3 Type C.	15
47	3.1.4 Type D.	16
48	3.1.5 Type E.	16
49	3.1.6 Type F.	16
50	3.2 Summary of successful tows by year and stratum	16

51	4 Discussion	20
52	4.1 Diversity of approaches used for mapping fish and invertebrates in the Scotian	
53	Shelf bioregion	20
54	4.2 Interpreting spatial results for marine spatial planning purposes	22
55	5 Acknowledgements	23
56	6 References	24
57	7 Appendix	28
58	7.1 Atlantic cod (Morue franche) - species code 10 (category LF)	29
59	7.2 Haddock (Aiglefin) - species code 11 (category LF)	32
60	7.3 White hake (Merluche blanche) - species code 12 (category LF)	35
61	7.4 Red hake (Merluche écureuil) - species code 13 (category LF)	38
62	7.5 Silver hake (Merlu argenté) - species code 14 (category LF)	41
63	7.6 Pollock (Goberge) - species code 16 (category LF)	44
64	7.7 Atlantic redfishes (Sébastes de l'Atlantique) - species code 23 (category LF)	47
65	7.8 Atlantic halibut (Flétan de l'Atlantique) - species code 30 (category LF)	50
66	7.9 American plaice (Plie canadienne) - species code 40 (category LF)	53
67	7.10 Witch flounder (Plie grise) - species code 41 (category LF)	56
68	7.11 Yellowtail flounder (Limande à queue jaune) - species code 42 (category LF)	59
69	7.12 Winter flounder (Limande-plie rouge) - species code 43 (category LF)	62
70	7.13 Atlantic wolffish (Loup atlantique) - species code 50 (category LF)	65
71	7.14 Atlantic herring (Hareng de l'Atlantique) - species code 60 (category LF)	68
72	7.15 Longhorn sculpin (Chabosseau à dix-huit épines) - species code 300 (category LF)	71
73	7.16 Moustache sculpin (Faux-trigle armé) - species code 304 (category LF)	74
74	7.17 Sea raven (Hémithriptère atlantique) - species code 320 (category LF)	77
75	7.18 Alligatorfish (Poisson-alligator atlantique) - species code 340 (category LF)	80
76	7.19 Monkfish (Baudroie d'Amérique) - species code 400 (category LF)	83

77	7.20 Ocean pout (Loquette d'Amérique) - species code 640 (category LF)	86
78	7.21 Thorny skate (Raie épineuse) - species code 201 (category LF)	89
79	7.22 Smooth skate (Raie lisse) - species code 202 (category LF)	92
80	7.23 Winter skate (Raie tachetée) - species code 204 (category LF)	95
81	7.24 Picked dogfish (Aiguillat commun) - species code 220 (category LF)	98
82	7.25 Northern shortfin squid (Encornet rouge nordique) - species code 4511 (category LF)	101
84	7.26 Atlantic hagfish (Myxine du nord) - species code 241 (category LI)	104
85	7.27 Cusk (Brosme) - species code 15 (category LI)	105
86	7.28 Greenland halibut (Flétan noir) - species code 31 (category LI)	106
87	7.29 Gulf Stream flounder (Plie du Gulf Stream) - species code 44 (category LI)	107
88	7.30 American shad (Alose savoureuse) - species code 61 (category LI)	108
89	7.31 Alewife (Gaspareau) - species code 62 (category LI)	109
90	7.32 Capelin (Capelan) - species code 64 (category LI)	110
91	7.33 Atlantic mackerel (Maquereau commun) - species code 70 (category LI)	111
92	7.34 Longfin hake (Merluche à longues nageoires) - species code 112 (category LI)	112
93	7.35 Fourbeard rockling (Motelle à quatre barbillons) - species code 114 (category LI)	113
94	7.36 Blackbelly rosefish (Sébaste chèvre) - species code 123 (category LI)	114
95	7.37 Greater argentine (Grande argentine) - species code 160 (category LI)	115
96	7.38 Arctic hookear sculpin (Hameçon neigeux) - species code 306 (category LI)	116
97	7.39 Atlantic poacher (Agone atlantique) - species code 350 (category LI)	117
98	7.40 Marlin-spike grenadier (Grenadier du Grand Banc) - species code 410 (category LI)	118
99	7.41 Lumpfish (Lompe) - species code 501 (category LI)	119
100	7.42 Atlantic spiny lumpsucker (Petite poule de mer atlantique) - species code 502 (category LI)	120
102	7.43 Sand lance (Lançon) - species code 610 (category LI)	121
103	7.44 Snakeblenny (Lompénie-serpent) - species code 622 (category LI)	122
104	7.45 Daubed shanny (Lompénie tachetée) - species code 623 (category LI)	123

105	7.46 Vahl's eelpout (Lycode à carreaux) - species code 647 (category LI)	124
106	7.47 Atlantic butterfish (Stromaté fossette) - species code 701 (category LI)	125
107	7.48 Atlantic hookear sculpin (Hameçon atlantique) - species code 880 (category LI) . . .	126
108	7.49 Barndoor skate (Grande raie) - species code 200 (category LI)	127
109	7.50 Little skate (Raie hérisson) - species code 203 (category LI)	128
110	7.51 Northern prawn (Crevette nordique) - species code 2211 (category SF)	129
111	7.52 Jonah crab (Tourteau jona) - species code 2511 (category SF)	130
112	7.53 Atlantic rock crab (Tourteau poïnclos) - species code 2513 (category SF)	131
113	7.54 Arctic lyre crab (Crabe Hyas coarctatus) - species code 2521 (category SF)	132
114	7.55 Atlantic king crab (Crabe épineux du nord) - species code 2523 (category SF) . . .	133
115	7.56 Queen crab (Crabe des neiges) - species code 2526 (category SF)	134
116	7.57 Great spider crab (Crabe lyre araignée) - species code 2527 (category SF)	135
117	7.58 American lobster (Homard américain) - species code 2550 (category SF)	136
118	7.59 Sea lamprey (Lamproie marine) - species code 240 (category LR)	137
119	7.60 Atlantic tomcod (Poulamon atlantique) - species code 17 (category LR)	138
120	7.61 Offshore silver hake (Merlu argenté du large) - species code 19 (category LR) . . .	139
121	7.62 Spotted wolffish (Loup tacheté) - species code 51 (category LR)	140
122	7.63 Northern wolffish (Loup à tête large) - species code 52 (category LR)	141
123	7.64 Rainbow smelt (Éperlan arc-en-ciel) - species code 63 (category LR)	142
124	7.65 Cunner (Tanche-tautogue) - species code 122 (category LR)	143
125	7.66 Fourspot flounder (Cardeau à quatre ocelles) - species code 142 (category LR) . .	144
126	7.67 Windowpane flounder (Turbot de sable) - species code 143 (category LR)	145
127	7.68 Longnose greeneye (Oeil-vert à long nez) - species code 149 (category LR)	146
128	7.69 Lanternfishes (Poissons-lanternes) - species code 150 (category LR)	147
129	7.70 Shortnose greeneye (Éperlan du large) - species code 156 (category LR)	148
130	7.71 Silvery lightfish (Brossé améthyste) - species code 158 (category LR)	149
131	7.72 Boa dragonfish (Dragon-boa) - species code 159 (category LR)	150

132	7.73 Shorthorn sculpin (Chabosseau à épines courtes) - species code 301 (category LR)	151
133	7.74 Grubby (Chabosseau bronzé) - species code 303 (category LR)	152
134	7.75 Polar sculpin (Cotte polaire) - species code 307 (category LR)	153
135	7.76 Spatulate sculpin (Icèle spatulée) - species code 314 (category LR)	154
136	7.77 Arctic alligatorfish (Poisson-alligator arctique) - species code 341 (category LR)	155
137	7.78 Alligatorfishes (Poissons-alligator) - species code 351 (category LR)	156
138	7.79 Roughnose grenadier (Grenadier-scie) - species code 412 (category LR)	157
139	7.80 Roundnose grenadier (Grenadier de roche) - species code 414 (category LR)	158
140	7.81 Atlantic seasnail (Limace atlantique) - species code 503 (category LR)	159
141	7.82 Gelatinous snailfish (Limace gélatineuse) - species code 505 (category LR)	160
142	7.83 Variegated snailfish (Limace marbée) - species code 512 (category LR)	161
143	7.84 Sea tadpole (Petite limace de mer) - species code 520 (category LR)	162
144	7.85 Wolf eelpout (Lycode à tête longue) - species code 603 (category LR)	163
145	7.86 Slender snipe eel (Avocette ruban) - species code 604 (category LR)	164
146	7.87 Newfoundland eelpout (Lycode du Labrador) - species code 619 (category LR)	165
147	7.88 Newfoundland eelpout (Lycode du Labrador) - species code 620 (category LR)	166
148	7.89 Rock gunnel (Sigouine de roche) - species code 621 (category LR)	167
149	7.90 Radiated shanny (Ulvaire deux-lignes) - species code 625 (category LR)	168
150	7.91 Fourline snakeblenny (Quatre-lignes atlantique) - species code 626 (category LR)	169
151	7.92 Wrymouth (Terrassier tacheté) - species code 630 (category LR)	170
152	7.93 Spotfin dragonet (Dragonnet tacheté) - species code 637 (category LR)	171
153	7.94 Arctic eelpout (Lycode arctique) - species code 641 (category LR)	172
154	7.95 Atlantic soft pout (Molasse atlantique) - species code 646 (category LR)	173
155	7.96 Silvery John dory (Saint Pierre argenté) - species code 704 (category LR)	174
156	7.97 White barracudina (Lussion blanc) - species code 712 (category LR)	175
157	7.98 Atlantic saury (Balaou atlantique) - species code 720 (category LR)	176
158	7.99 Hatchetfishes (Haches d'argent) - species code 741 (category LR)	177

159	7.100	Atlantic batfish (Malthe atlantique) - species code 742 (category LR)	178
160	7.101	Spottedfin tonguefish (Langue fil noir) - species code 816 (category LR)	179
161	7.102	Black dogfish (Aiguillat noir) - species code 221 (category LR)	180
162	7.103	Longfin inshore squid (Calmar totam) - species code 4512 (category LR)	181
163	7.104	Red deepsea crab (Crabe rouge) - species code 2532 (category SR)	182

164	INDEX	183
-----	--------------	------------

ABSTRACT

166 Ricard, D., Emberley, J., Gomez, C. and Regnier-McKellar, C. 2021. Marine Fish and Invertebrate
167 Atlas: Summarizing Geographic Distribution, Population Indices and Environmental
168 Preferences in the Scotian Shelf and Bay of Fundy (1970-2020). Can. Tech. Rep. Fish. Aquat.
169 Sci. nnn: x + 188 p.

170 The summer groundfish research vessel survey on the Scotian Shelf and in the Bay of
171 Fundy started in 1970 and was designed to measure the distribution and abundance of
172 major commercial fish species. Over time, additional information on non-commercial species
173 was collected, and allowed considerable insight into ecosystem function and structure, as
174 documented in many primary publications whose analyses used the survey data. The same
175 groundfish survey database has also been used to produce species status reports, atlases of
176 species distribution and remains an essential source of information for stock assessments in the
177 Maritimes Region of Fisheries and Oceans Canada. This report builds on previous work and
178 former atlases by updating a comprehensive suite of indices to assess population status and
179 environmental preferences of 104 species. For each species, trends in geographic distribution
180 and biomass or abundance were plotted. The spatial extent of distribution was plotted over
181 time to gauge how the area occupied has changed. The relationship between abundance or
182 biomass and spatial extent reflected whether the species distribution expands when abundance
183 or biomass increases. Length frequencies over time depicted any changes in mean size. The
184 plots of condition over time revealed whether individual fish are fatter or thinner than their long
185 term mean. Depth, temperature and salinity preferences were estimated to gauge the range
186 of suitable environmental parameters for each species. Finally, for each stratum, the slope
187 describing how local density varies with regional abundance was estimated.

RÉSUMÉ

189 Ricard, D., Emberley, J., Gomez, C. and Regnier-McKellar, C. 2021. Marine Fish and Invertebrate
190 Atlas: Summarizing Geographic Distribution, Population Indices and Environmental
191 Preferences in the Scotian Shelf and Bay of Fundy (1970-2020). Can. Tech. Rep. Fish. Aquat.
192 Sci. nnn: x + 188 p.

193 Le relevé estival par navires de recherche sur le poisson de fond sur le plateau néo-écossais
194 et dans la baie de Fundy a débuté en 1970 et visait à mesurer la répartition et l'abondance
195 des principales espèces de poissons commerciales. Au fil du temps, des informations
196 supplémentaires sur les espèces non commerciales ont été recueillies et ont permis de mieux
197 comprendre la fonction et la structure de l'écosystème, comme le montrent de nombreuses
198 publications primaires dont les analyses ont utilisé les données des relevés. La même base
199 de données sur les relevés du poisson de fond a également été utilisée pour produire des
200 rapports sur la situation des espèces, des atlas de la répartition des espèces et demeure une
201 source essentielle d'information pour les évaluations des stocks dans la région des Maritimes
202 de Pêches et Océans Canada. Ce rapport s'appuie sur des travaux antérieurs et d'anciens
203 atlas en mettant à jour une série complète d'indices pour évaluer l'état de la population et les
204 préférences environnementales de 104 espèces. Pour chaque espèce, les tendances de la
205 répartition géographique et de la biomasse ou de l'abondance ont été tracées. L'étendue spatiale
206 de la distribution a été tracée au fil du temps pour évaluer comment la zone occupée a changé.
207 La relation entre l'abondance ou la biomasse et l'étendue spatiale indique si la répartition des
208 espèces augmente lorsque l'abondance ou la biomasse augmente. Les fréquences de longueur
209 au fil du temps représentaient tout changement dans la taille moyenne. Les graphiques de l'état
210 au fil du temps ont révélé si les poissons individuels sont plus gros ou plus minces que leur
211 moyenne à long terme. Les préférences en matière de profondeur, de température et de salinité
212 ont été estimées pour évaluer la gamme de paramètres environnementaux appropriés pour
213 chaque espèce. Enfin, pour chaque strate, la pente décrivant comment la densité locale varie
214 avec l'abondance régionale a été estimée.

1 Introduction

216 The summer (July-August) groundfish research vessel survey on the Scotian Shelf and in the
217 Bay of Fundy was started in 1970 by Fisheries and Oceans Canada Maritimes Region. The
218 survey was originally designed to measure the distribution and abundance of major commercial
219 fish species. Over time, information on non-commercial species was also collected. The
220 groundfish survey database storing the information collected during the annual survey provides
221 the main source of fisheries-independent information for marine species in the region. This
222 information is routinely used to support stock assessments, to produce species status reports
223 and has been previously used to publish atlases of species distribution.

224 The current document is an update of an earlier report (Ricard and Shackell 2013) that built on
225 former atlases by updating a comprehensive suite of derived indices for 104 species to assess
226 population status and environmental preferences. The information collected during the survey is
227 stored in a relational database management system archived at Fisheries and Oceans Canada
228 Maritimes Region which contains detailed information about the sampling locations and the
229 associated catch. Tow-level survey data is also publicly available from the Ocean Biogeographic
230 Information System (DFO 2016) and (DFO 2021). The present atlas follows on the work done
231 by Fisheries and Oceans colleagues from the northern Gulf of St. Lawrence (Bourdages and
232 Ouellet 2012), southern Gulf of St. Lawrence (Benoît et al. 2003) and on earlier work in the
233 Scotian Shelf (Simon and Comeau 1994; Horsman and Shackell 2009).

234 To facilitate updates and foster collaboration on the analyses of the survey data, the computer
235 code necessary to extract the data, to perform the analyses presented herein, and to reproduce
236 and update the current document is made available in a git repository (Ricard and Gomez 2021).

237 The survey area covers three major Northwest Atlantic Fisheries Organization (NAFO) zones
238 that divide the shelf into the colder east 4V and 4W (strata 440-466) and warmer west 4X (strata
239 470-495). Temporal trends are plotted by NAFO regions for several species. For each species,
240 trends in geographic distribution and biomass or abundance are plotted. Some caution is
241 required in interpreting the results obtained for several taxa due to low sample size as explained
242 later in the text. The spatial extent of distribution is plotted over time to gauge how the area
243 occupied has changed. The relationship between biomass and spatial extent reflects whether the
244 species distribution expands when biomass increases. For each strata, the slope describing how
245 local density varies with regional abundance was estimated (Myers and Stokes 1989). These
246 slopes were then plotted against a habitat suitability index to identify important strata for each
247 species. Then, length frequencies over time depicted any changes in mean size. The plots of
248 condition over time revealed whether individual fish are fatter or thinner than their long term
249 mean. Finally, depth, temperature and salinity preferences were estimated to gauge the range
250 of environmental parameters (Perry and Smith 1994). A full ecological interpretation of trends
251 is beyond the scope of this report. Other documents stemming from peer-reviewed scientific
252 processes under the auspices of the [Canadian Science Advisory Secretariat](#) (CSAS) provide
253 further descriptions of spatio-temporal trends in different indicators and put the information
254 collected during the summer groundfish research vessel survey in a more focused context (see
255 for example Clark and Emberley (2011)).

2 Methods

257 2.1 Survey Description

258 The survey is conducted annually in July-August and covers the Scotian Shelf and the Bay of
 259 Fundy (Figure 1). It normally involves two separate two-week trips on board an offshore fisheries
 260 vessel from the Canadian Coast Guard.

261 A number of changes in fishing gear type and vessels used occurred since the onset of sampling
 262 activities (Clark and Emberley 2011).




Figure 1. Map of the Scotian Shelf and Bay of Fundy.

263 **2.2 Sampling Design**

264 The summer survey covers divisions 4V, 4W and 4X of the Northwest Atlantic Fisheries
265 Organization (NAFO) which includes the Scotian Shelf and the Bay of Fundy. The eastern limit of
266 the survey is the Laurentian Channel and the western limit is the Fundian Channel (Figure 1).

267 The survey follows a stratified random design (Doubleday and Rivard 1981; Lohr 1999)
268 (Figure 2). The number of tows conducted in each stratum is approximately proportional to its
269 surface area.




Figure 2. Map of the Summer survey strata.

270 The basic sampling unit of the survey is a 30-minute fishing tow conducted at a speed of 3.5
271 knots. This yields a distance towed of 1.75 nautical miles.

272 After each tow the catch is sorted by species and weighed. Each fish caught is then measured,
273 and further sampling of individual fish weight, maturity status and age are performed for different
274 length classes. When catches exceed 300 individuals, a random sub-sample is used to obtain
275 the length and weight measurements.

276 The location of representative tows appears in Figure 3.




Figure 3. Map of the Summer survey tows.

277 **2.3 Taxonomic Levels**

278 Fish species caught during the surveys are identified by trained scientific personnel and their
279 scientific name is determined. An internal species code used in the relational database is
280 reported for each species (Losier and Waite 1989).

281 By its nature as a bottom trawl, the fishing gear used in the survey catches certain species
282 better than others. To ensure that meaningful ecological information can be extracted from
283 catch samples, we report the catch records for the subset of species that are caught reliably
284 by the gear. To appear in this atlas, a species must have had a minimum of 10 observations over
285 the duration of the survey activities. While both catch abundance and weight are recorded, the
286 weight of species that appear at low abundances is often recorded as zero in the earlier parts of
287 the survey when scales of appropriate precision were not available.

288 We divided the species caught into five categories based on 1) their taxonomic classification,
289 2) the number of recorded observations, and 3) their period of valid identification (Table 1).
290 Category "LF", for "long frequent", was assigned to species that have more than 1000 records
291 since 1970 and have been consistently identified since the onset of the survey. Category
292 "LI", for "long intermediate", was assigned to species that had between 1000 and 200 catch
293 records. Rare and elusive species (those with less than 200 catch records over the duration
294 of the survey) are also reported but to a lower level of analytical details (Category "LR", for
295 "long rare"). Category "SF", for "short frequent", was assigned to invertebrate species that were
296 consistently sampled only since 1999 (Tremblay M. J. 2007). And category "SR", for "short rare"
297 for invertebrate species consistently sampled only since 1999 and with less than 200 catch
298 records. The list of taxa covered in this document is presented in phylogenetic order (Nelson J.
299 S. et al. 2004) in Table 2. To ensure concordance with authoritative taxonomic information, the
300 AphiaID from the World Register of Marine Species is also provided in Table 2 (Appeltans et al.
301 2012).

Category	Name	Description
L	long - consistently identified since the onset of the survey in 1970	
LF	long frequent	species that have more than 1000 catch records
LI	long intermediate	species that had between 1000 and 200 catch records
LR	long rare	species with less than 200 catch records
S	short - invertebrate species that were consistently sampled only since 1999	
SF	short frequent	species with more than 200 catch records
SR	short rare	species with less than 200 catch records

Table 1. Taxonomic levels

Table 2. List of species included in the Atlas. The species reported here are listed in phylogenetic order as per Page L. M. et al. (2013). For each taxonomic order and class, each species is listed in the table, its taxonomic family and scientific name is provided, along with its French and English common names, the species code used in the survey database, its AphiaID and a link to the World Registry of Marine Species, its number of catch records in the survey database and its classification category as defined in section 2.3.

Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Myxini							
<i>Myxiniformes</i>							
Myxinidae	<i>Myxine glutinosa</i>	Atlantic hagfish	Myxine du nord	241	101170	804	LI
Petromyzonti							
<i>Petromyzontiformes</i>							
Petromyzontidae	<i>Petromyzon marinus</i>	Sea lamprey	Lamproie marine	240	101174	16	LR
Actinopterygii							
<i>Gadiformes</i>							
Gadidae	<i>Gadus morhua</i>	Atlantic cod	Morue franche	10	126436	5451	LF
	<i>Melanogrammus aeglefinus</i>	Haddock	Aiglefin	11	126437	5827	LF
Phycidae	<i>Urophycis tenuis</i>	White hake	Merluche blanche	12	126504	3524	LF
	<i>Urophycis chuss</i>	Red hake	Merluche écureuil	13	126503	2195	LF
Merlucciidae	<i>Merluccius bilinearis</i>	Silver hake	Merlu argenté	14	158962	4936	LF
Lotidae	<i>Brosme brosme</i>	Cusk	Brosme	15	126447	688	LI
Gadidae	<i>Pollachius virens</i>	Pollock	Goberge	16	126441	2787	LF
	<i>Microgadus tomcod</i>	Atlantic tomcod	Poulamon atlantique	17	158928	44	LR
Merlucciidae	<i>Merluccius albidus</i>	Offshore silver hake	Merlu argenté du large	19	158748	161	LR
<i>Scorpaeniformes</i>							
Sebastidae	<i>Sebastes</i>	Atlantic redfishes	Sébastes de l'Atlantique	23	126175	4152	LF
<i>Pleuronectiformes</i>							
Pleuronectidae	<i>Hippoglossus hippoglossus</i>	Atlantic halibut	Flétan de l'Atlantique	30	127138	1634	LF

Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Pleuronectidae	<i>Reinhardtius hippoglossoides</i>	Greenland halibut	Flétan noir	31	127144	736	LI
	<i>Hippoglossoides platessoides</i>	American plaice	Plie canadienne	40	127137	6023	LF
	<i>Glyptocephalus cynoglossus</i>	Witch flounder	Plie grise	41	127136	4301	LF
	<i>Limanda ferruginea</i>	Yellowtail flounder	Limande à queue jaune	42	158879	3233	LF
	<i>Pseudopleuronectes americanus</i>	Winter flounder	Limande-plie rouge	43	158885	1632	LF
Paralichthyidae	<i>Citharichthys arctifrons</i>	Gulf Stream flounder	Plie du Gulf Stream	44	158791	382	LI
<i>Perciformes</i>							
Anarhichadidae	<i>Anarhichas lupus</i>	Atlantic wolffish	Loup atlantique	50	126758	1572	LF
	<i>Anarhichas minor</i>	Spotted wolffish	Loup tacheté	51	126759	20	LR
	<i>Anarhichas denticulatus</i>	Northern wolffish	Loup à tête large	52	126757	17	LR
<i>Clupeiformes</i>							
Clupeidae	<i>Clupea harengus</i>	Atlantic herring	Hareng de l'Atlantique	60	126417	3487	LF
	<i>Alosa sapidissima</i>	American shad	Alose savoureuse	61	158670	468	LI
	<i>Alosa pseudoharengus</i>	Alewife	Gaspareau	62	158669	977	LI
<i>Osmeriformes</i>							
Osmeridae	<i>Osmerus mordax</i>	Rainbow smelt	Éperlan arc-en-ciel	63	126737	59	LR
	<i>Mallotus villosus</i>	Capelin	Capelan	64	126735	540	LI
<i>Perciformes</i>							
Scombridae	<i>Scomber scombrus</i>	Atlantic mackerel	Maquereau commun	70	127023	696	LI
<i>Gadiformes</i>							

Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Phycidae	<i>Phycis chesteri</i>	Longfin hake	Merluche à longues nageoires	112	158988	784	LI
Lotidae	<i>Enchelyopus cimbricus</i>	Fourbeard rockling	Motelle à quatre barbillons	114	126450	693	LI
<i>Perciformes</i>							
Labridae	<i>Tautogolabrus adspersus</i>	Cunner	Tanche-tautogue	122	159785	82	LR
<i>Scorpaeniformes</i>							
Sebastidae	<i>Helicolenus dactylopterus</i>	Blackbelly rosefish	Sébaste chèvre	123	127251	610	LI
<i>Pleuronectiformes</i>							
Paralichthyidae	<i>Hippoglossina oblonga</i>	Fourspot flounder	Cardeau à quatre ocelles	142	158833	76	LR
Scophthalmidae	<i>Scophthalmus aquosus</i>	Windowpane flounder	Turbot de sable	143	158907	115	LR
<i>Aulopiformes</i>							
Chlorophthalmidae	<i>Parasudis truculenta</i>	Longnose greeneye	Oeil-vert à long nez	149	158868	45	LR
<i>Myctophiformes</i>							
Myctophidae	<i>Myctophidae</i>	Lanternfishes	Poissons-lanternes	150	125498	160	LR
<i>Aulopiformes</i>							
Chlorophthalmidae	<i>Chlorophthalmus agassizi</i>	Shortnose greeneye	Éperlan du large	156	126336	78	LR
<i>Stomiiformes</i>							
Sternopychidae	<i>Maurolicus muelleri</i>	Silvery lightfish	Brossé améthyste	158	127312	52	LR
Stomiidae	<i>Stomias boa</i>	Boa dragonfish	Dragon-boa	159	127374	20	LR
<i>Argentiniformes</i>							
Argentinidae	<i>Argentina silus</i>	Greater argentine	Grande argentine	160	126715	963	LI
<i>Scorpaeniformes</i>							
Cottidae	<i>Myoxocephalus octodecemspinosus</i>	Longhorn sculpin	Chabosseau à dix-huit épines	300	159520	3292	LF

Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
	<i>Myoxocephalus scorpius</i>	Shorthorn sculpin	Chabosseau à épines courtes	301	127203	131	LR
	<i>Myoxocephalus aenaeus</i>	Grubby	Chabosseau bronzé	303	159519	40	LR
	<i>Triglops murrayi</i>	Moustache sculpin	Faux-trigle armé	304	127205	1182	LF
	<i>Arctediellus uncinatus</i>	Arctic hookear sculpin	Hameçon neigeux	306	127195	306	LI
Psychrolutidae	<i>Cottunculus microps</i>	Polar sculpin	Cotte polaire	307	127235	29	LR
Cottidae	<i>Icelus spatula</i>	Spatulate sculpin	Icele spatulée	314	127200	40	LR
Hemitripteridae	<i>Hemitripterus americanus</i>	Sea raven	Hémithriptère atlantique	320	159518	2126	LF
Agonidae	<i>Aspidophoroides monopterygius</i>	Alligatorfish	Poisson-alligator atlantique	340	159459	1029	LF
	<i>Ulcina olrikii</i>	Arctic alligatorfish	Poisson-alligator arctique	341	274356	13	LR
	<i>Leptagonus decagonus</i>	Atlantic poacher	Agone atlantique	350	127191	266	LI
	<i>Agonidae</i>	Alligatorfishes	Poissons-alligator	351	125588	43	LR
<i>Lophiiformes</i>							
Lophiidae	<i>Lophius americanus</i>	Monkfish	Baudroie d'Amérique	400	159184	1970	LF
<i>Gadiformes</i>							
Macrouridae	<i>Nezumia bairdii</i>	Marlin-spike grenadier	Grenadier du Grand Banc	410	183289	529	LI
	<i>Trachyrincus murrayi</i>	Roughnose grenadier	Grenadier-scie	412	126481	18	LR
	<i>Coryphaenoides rupestris</i>	Roundnose grenadier	Grenadier de roche	414	158960	17	LR
<i>Scorpaeniformes</i>							
Cyclopteridae	<i>Cyclopterus lumpus</i>	Lumpfish	Lompe	501	127214	216	LI

Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
	<i>Eumicrotremus spinosus</i>	Atlantic spiny lump sucker	Petite poule de mer atlantique	502	127217	226	LI
Liparidae	<i>Liparis atlanticus</i>	Atlantic seasnail	Limace atlantique	503	159524	34	LR
	<i>Liparis fabricii</i>	Gelatinous snailfish	Limace gélatineuse	505	127218	27	LR
	<i>Liparis gibbus</i>	Variegated snailfish	Limace marbée	512	159526	41	LR
	<i>Careproctus reinhardtii</i>	Sea tadpole	Petite limace de mer	520	127212	18	LR
<i>Perciformes</i>							
Zoarcidae	<i>Lycenchelys verrillii</i>	Wolf eelpout	Lycode à tête longue	603	159258	40	LR
<i>Anguilliformes</i>							
Nemichthyidae	<i>Nemichthys scolopaceus</i>	Slender snipe eel	Avocette ruban	604	126306	28	LR
<i>Perciformes</i>							
Ammodytidae	<i>Ammodytes dubius</i>	Sand lance	Lançon	610	151520	1283	LI
Zoarcidae	<i>Lycodes terraenovae</i>	Newfoundland eelpout	Lycode du Labrador	619	127117	64	LR
	<i>Lycodes lavalaei</i>	Newfoundland eelpout	Lycode du Labrador	620	127107	72	LR
Pholidae	<i>Pholis gunnellus</i>	Rock gunnel	Sigouine de roche	621	126996	21	LR
Stichaeidae	<i>Lumpenus lampretaeformis</i>	Snakeblenny	Lompénie-serpent	622	154675	423	LI
	<i>Leptoclinus maculatus</i>	Daubed shanny	Lompénie tachetée	623	127072	443	LI
	<i>Ulvaria subbifurcata</i>	Radiated shanny	Ulvaire deux-lignes	625	159821	145	LR
	<i>Eumesogrammus praecisus</i>	Fourline snakeblenny	Quatre-lignes atlantique	626	159817	40	LR
Cryptacanthodidae	<i>Cryptacanthodes maculatus</i>	Wrymouth	Terrassier tacheté	630	159675	120	LR
Callionymidae	<i>Foetorepus agassizii</i>	Spotfin dragonet	Dragonnet tacheté	637	276339	20	LR

Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Zoarcidae	<i>Zoarces americanus</i>	Ocean pout	Loquette d'Amérique	640	159267	1478	LF
	<i>Lycodes reticulatus</i>	Arctic eelpout	Lycode arctique	641	127112	70	LR
	<i>Melanostigma atlanticum</i>	Atlantic soft pout	Molasse atlantique	646	127120	43	LR
	<i>Lycodes vahlii</i>	Vahl's eelpout	Lycode à carreaux	647	127118	565	LI
Stromateidae	<i>Peprilus triacanthus</i>	Atlantic butterfish	Stromaté fossette	701	159828	487	LI
<i>Zeiformes</i>							
Zeidae	<i>Zenopsis conchifer</i>	Silvery John dory	Saint Pierre argenté	704	127426	39	LR
<i>Aulopiformes</i>							
Paralepididae	<i>Arctozenus risso</i>	White barracudina	Lussion blanc	712	126352	196	LR
<i>Beloniformes</i>							
Scomberesocidae	<i>Scomberesox saurus</i>	Atlantic saury	Balaou atlantique	720	126392	37	LR
<i>Stomiiformes</i>							
Sternopychidae	<i>Sternopychidae</i>	Hatchetfishes	Haches d'argent	741	125603	21	LR
<i>Lophiiformes</i>							
Ogcocephalidae	<i>Dibranchus atlanticus</i>	Atlantic batfish	Malthe atlantique	742	126558	18	LR
<i>Pleuronectiformes</i>							
Cynoglossidae	<i>Symphurus diomedeanus</i>	Spottedfin tonguefish	Langue fil noir	816	159358	24	LR
<i>Scorpaeniformes</i>							
Cottidae	<i>Artediellus atlanticus</i>	Atlantic hookear sculpin	Hameçon atlantique	880	127193	258	LI
<i>Elasmobranchii</i>							
<i>Rajiformes</i>							
Rajidae	<i>Dipturus laevis</i>	Barndoor skate	Grande raie	200	158548	246	LI
	<i>Amblyraja radiata</i>	Thorny skate	Raie épineuse	201	105865	3937	LF
	<i>Malacoraja senta</i>	Smooth skate	Raie lisse	202	158554	1773	LF

Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category	
	<i>Leucoraja erinacea</i>	Little skate	Raie hérisson	203	158551	712	LI	
	<i>Leucoraja ocellata</i>	Winter skate	Raie tachetée	204	158553	1180	LF	
Squaliformes	Squalidae	<i>Squalus acanthias</i>	Picked dogfish	Aiguillat commun	220	105923	1985	LF
	Etmopteridae	<i>Centroscyllium fabricii</i>	Black dogfish	Aiguillat noir	221	105906	31	LR
Cephalopoda								
<i>Oegopsida</i>	Ommastrephidae	<i>Illex illecebrosus</i>	Northern shortfin squid	Encornet rouge nordique	4511	153087	4836	LF
<i>Myopsida</i>	Loliginidae	<i>Doryteuthis pealeii</i>	Longfin inshore squid	Calmar totam	4512	574541	96	LR
Malacostraca								
<i>Decapoda</i>	Pandalidae	<i>Pandalus borealis</i>	Northern prawn	Crevette nordique	2211	107649	718	SF
	Cancridae	<i>Cancer borealis</i>	Jonah crab	Tourteau jona	2511	158056	1387	SF
	Oregoniidae	<i>Hyas coarctatus</i>	Arctic lyre crab	Crabe Hyas coarctatus	2521	107323	711	SF
	Lithodidae	<i>Lithodes maja</i>	Atlantic king crab	Crabe épineux du nord	2523	107205	531	SF
	Oregoniidae	<i>Chionoecetes opilio</i>	Queen crab	Crabe des neiges	2526	107315	1546	SF
		<i>Hyas araneus</i>	Great spider crab	Crabe lyre araignée	2527	107322	625	SF
	Geryonidae	<i>Chaceon quinquedens</i>	Red deepsea crab	Crabe rouge	2532	158407	33	SR
	Nephropidae	<i>Homarus americanus</i>	American lobster	Homard américain	2550	156134	1623	SF

302 **2.4 Analyses**

303 The Oracle relational database where all data are stored was accessible from the Bedford
304 Institute of Oceanography in Dartmouth, Nova Scotia. Structured Query Language (SQL) is
305 used to extract the data from the production server and to create the data products used in
306 all subsequent analyses. Catch records classified as "valid" (i.e. a representative tow without
307 damage to the net) are used in the current analyses. To make the available samples comparable,
308 catch number and weight for each species was standardized for the distance towed.

309 All data processing and analyses were conducted using the R software (R Core Team 2020)
310 using packages gstat (Pebesma 2004), PBSmapping (Schnute et al. 2019), RODBC (Ripley
311 and Lapsley 2019), spatstat (Baddeley 2015), maptools (Bivand and Lewin-Koh 2020), rgeos
312 (Bivand and Rundel 2020), classInt(Bivand 2020), RColorBrewer(Neuwirth 2014), MASS (Ripley
313 et al. 2020), worms (Holstein 2018), and tidyverse (Wickham 2019). The present document is
314 rendered as a Technical Report using the csasdown R package developed and maintained by
315 Fisheries and Oceans Canada scientists (Anderson et al. In press).

316 **2.4.1 Geographic distribution of catches**

317 Spatial interpolation of catch biomass (kg/tow) or abundance (number/tow) was done using a
318 weighting inversely proportional to the distance, using function "idw" of the spatstat R package
319 (Baddeley 2015).

320 **2.4.2 Abundance and biomass indices**

321 For each species, stratified random estimates of catch abundance and biomass (Smith 1996)
322 were computed for each year. Yearly estimates of the standard error were also computed.

323 **2.4.3 Distribution indices**

324 For each Category L, I and S fish species, the minimum area required to account for 75% and
325 95% of the total biomass or abundance were computed (D75% and D95%). These measures of
326 distributions were computed for each year by using the Lorenz curve of mean stratum-level catch
327 estimates and the area of occupied strata (Swain and Sinclair 1994; Swain and Morin 1996).

328 **2.4.4 Length frequencies**

329 The length frequency distribution of catch is tabulated for each seven-year period (1970-2009),
330 and last ten-year period (2010-2020).

331 **2.4.5 Length-weight relationship and condition factor**

332 The relationship between the weight and the length of fish was estimated using the following
333 non-linear isometric relationship:

$$W = \alpha L^\beta$$

334 where W is the total weight (g), L is the length (cm), and, α and β are the parameters to be
335 estimated.

336 Average fish condition (C) was computed as:

$$C = \frac{W}{\alpha L^\beta}$$

337

338 **2.4.6 Depth, temperature and salinity distribution of catches**

339 For each category L species, We followed the methods developed by (Perry and Smith 1994)
340 and generated cumulative frequency distributions of depth, temperature and salinity of survey
341 catches.

342 **2.4.7 Density-dependent habitat selection**

343 We followed the methods of (Myers and Stokes 1989) to evaluate how fish abundance in each
344 stratum varied with overall temporal fluctuations of population abundance.

345 For each category L species, we fitted a model of the relationship between stratum-level density
346 and overall abundance (the yearly stratified random estimate of abundance, defined above).
347 To properly use the observations of zero catch while accounting for the logarithmic distribution
348 of catch abundance, we implemented the model as a generalised linear using a log link and a
349 Poisson error distribution:

$$Y_{h,i} = \alpha_{h,i} Y_i^{\beta_{h,i}}$$

350 where, $y_{h,i}$ is the average abundance of stratum h in year i , and $\alpha_{h,i}$ and $\beta_{h,i}$ are the fitted
351 parameters. The estimated parameter $\beta_{h,i}$ is referred to as the “slope parameter” and indicates
352 whether stratum-level density is positively ($\beta_{h,i} <= 0$), negatively ($\beta_{h,i} >= 0$) or negligibly
353 ($\beta_{h,i} \approx 0$) related to population abundance.

354 To estimate the suitability of each stratum, the median abundance observed during the years
355 that are in the top 25% of yearly estimates is used. We combine the slope parameter estimates
356 from the above model with the median abundance to identify strata that have consistently high
357 abundance and whose local density is weakly related to fluctuation in population abundance
358 ($\beta_{h,i} \approx 0$). Preferred strata are identified for each category L species.

359

3 Results

360 The plots generated for each species are presented in the Appendix.

361 3.1 Description of Figures

362 3.1.1 Type A

363 For Category L and S species:

364 Spatial distribution of catch-per unit of effort, (CPUE, kilograms per tow) in July-August for the
365 Bay of Fundy and Scotian Shelf in five-year periods. Spatial interpolation between tows was
366 done using Inverse Distance Weight (IDW). The probability of occurrence (proportion of tows with
367 catch records for a given species) was also reported for each five-year period.

368 For Category LR and SR:

369 Location of tows with catch over the period 1970-2012 (Type LR) or the period 1999-2012 (Type
370 SR). Location of tows with catch over the period 1970-2012 (Type LR) or the period 1999-2012
371 (Type SR).

372 3.1.2 Type B

373 For Category L, S and I species:

374 Stratified random estimate of CPUE (left panel), distribution indices (D75% and D95%, the
375 minimum area containing 75% and 95% of biomass, middle panel), and distribution vs. weight
376 per tow (right panel). The stratified random mean is plotted as a solid line with the 95%
377 confidence region indicated by the solid grey line. The overall mean is plotted as a grey
378 horizontal line and the overall mean plus or minus 50% of the standard deviation appear as
379 horizontal dashed lines. In all three panels, the early years appear in blue and the last years
380 appear in red. The predictions from a loess estimator are overlaid on the distribution indices
381 (middle panel). The Pearson correlation coefficient between D75% and biomass, and its
382 statistical significance, are also reported in the right panel.

383 3.1.3 Type C.

384 Length frequency distribution for NAFO divisions 4X and 4VW. A smoothed length frequency
385 distribution is shown for each 7-year periods covered by the surveys.

386 **3.1.4 Type D.**

387 Average fish condition for all fish lengths (black dots and black line), large fish (thick gray line),
388 and small fish (thin gray line). Fish condition is presented for NAFO divisions 4VW (right panel)
389 and 4X (left panel).

390 **3.1.5 Type E.**

391 Cumulative frequency distributions of depth, temperature and salinity at all sampled locations
392 (thick solid line) and at fishing locations with catch records (thin dashed line). The depth,
393 temperature and salinity associated with 5%, 25%, 50%, 75% and 95% of the cumulative catch is
394 shown in tabular fashion on the bottom right panel.

395 **3.1.6 Type F.**

396 Slopes estimates from the density-dependent habitat selection model (y axis) plotted versus
397 the median abundance during the top 25% of years. The red box indicates strata of particular
398 importance for a species by identifying slopes that are within a standard error from zero and that
399 are within the top 25% of median abundance. Each stratum is identified on the plot by the last
400 two digits of its number.

401 **3.2 Summary of successful tows by year and stratum**

402 There is something weird going on here, there are 2 tows with NAs for stratum, (HAM1980042
403 set 62 and HAM1982072 set 13).

<!-- Number of tows by stratum-year -->

Table 3. Number of representative tows conducted in each stratum during the period 1970 to 1991.

Stratum	NAFO Div.	Area (km2)	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991
440	4VN	3173.016	4	2	2	3	3	3	3	3	3	3	3	3	3	3	3	4	5	5	6	4	4	4
441	4VN	3434.000	4	2	2	3	3	3	1	3	3	3	3	3	3	3	3	5	5	4	4	6	5	5
442	4VN	4934.658	3	2	2	2	3	3	2	3	3	3	3	3	3	3	3	3	5	6	7	5	5	5
443	4VSW	4526.012	4	2	4	4	8	3	1	2	4	4	4	3	5	4	4	4	6	6	5	2	4	2
444	4VSW	13478.450	3	2	5	4	6	4	6	7	4	4	4	5	5	6	4	4	6	6	3	6	7	8
445	4VSW	3512.982	5	2	5	4	5	5	1	3	4	4	4	5	5	3	4	5	6	4	4	4	4	4
446	4VSW	1686.094	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	4	3	3	3	3	3	3
447	4VSW	5549.344	4	2	6	5	7	4	4	3	4	4	5	4	4	4	4	4	5	7	6	6	8	7
448	4VSW	4975.866	5	2	5	4	5	4	4	4	4	4	4	6	4	4	4	5	5	5	5	9	6	6
449	4VSW	494.496	2	2	2	2	3	2	2	2	1	2	2	2	1	2	2	2	2	2	2	2	2	2
450	4VSW	1315.222	2	2	3	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
451	4VSW	504.798	1	2	2	2	2	2	2	2	2	3	2	2	3	2	2	2	2	2	2	2	2	2
452	4VSW	1184.730	2	2	2	2	2	2	2	2	2	2	2	4	2	2	2	2	2	3	2	2	2	3
453	4VSW	889.406	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	2	2	2	3
454	4VSW	1713.566	3	2	3	3	3	3	3	3	3	3	3	2	3	3	3	3	3	3	2	2	2	3
455	4VSW	7286.948	7	6	7	6	7	6	6	7	7	7	7	7	7	7	7	8	8	7	7	12	10	10
456	4VSW	3279.470	5	4	6	5	5	6	4	6	6	6	6	6	7	6	6	6	6	7	6	6	10	7
457	4VSW	2784.974	2	2	2	2	3	2	2	2	2	2	3	2	2	2	2	2	2	4	2	2	4	2
458	4VSW	2259.572	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	5	5	3	3	9	8
459	4VSW	10810.232	3	2	4	4	4	4	4	4	4	4	4	4	3	4	4	6	6	5	6	5	5	5
460	4VSW	4615.296	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2	4	3	3	3	3
461	4VSW	3962.836	3	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3	3	2	1	2
462	4VSW	7266.344	3	3	4	3	4	4	4	4	4	4	4	4	4	4	4	4	6	5	4	4	5	5
463	4VSW	1037.068	2	2	2	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2	3	2
464	4VSW	4453.898	4	3	5	3	3	6	5	5	5	5	5	5	5	4	5	5	5	7	6	5	5	9
465	4VSW	8183.222	6	5	5	4	5	4	5	5	5	5	5	7	6	5	5	5	5	8	8	8	12	9
466	4VSW	776.084	2	2	3	2	3	3	3	3	3	3	3	2	3	3	3	3	3	3	2	2	3	2
470	4X	3159.280	1	2	2	2	3	2	2	2	2	2	2	2	2	2	2	2	2	2	3	3	2	2
471	4X	3447.736	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
472	4X	4289.066	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2	2	4	4	6	4
473	4X	910.010	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3	2
474	4X	552.874	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	0	2	2	2	2	2	2
475	4X	535.704	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
476	4X	5075.452	2	2	2	2	2	2	3	2	2	2	1	2	2	2	2	2	2	2	4	4	4	4
477	4X	4230.688	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2	5	4	4	5	5
478	4X	800.122	2	2	3	2	3	3	3	3	2	3	3	3	3	3	3	3	3	2	2	2	2	
480	4X	2249.270	4	4	4	3	3	3	4	4	3	4	3	3	4	4	4	4	4	4	4	4	8	
481	4X	6438.750	5	3	4	4	4	3	4	4	5	4	3	4	4	4	4	4	4	6	7	6	8	
482	4X	3578.228	2	1	2	2	2	2	3	2	2	3	2	2	2	2	2	2	2	3	3	3	3	
483	4X	1826.888	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
484	4X	7774.576	2	2	3	3	3	3	3	3	2	3	3	3	4	3	3	3	4	4	4	4	3	
485	4X	5432.588	2	2	2	3	3	3	3	3	3	2	3	4	3	3	3	3	6	7	6	2	3	
490	4X	2063.834	2	2	2	2	2	3	3	3	3	2	3	3	3	3	3	3	3	4	4	4	4	
491	4X	2359.158	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	4	4	4	3	
492	4X	3729.324	3	2	4	3	3	3	3	3	3	3	3	3	3	3	3	3	3	4	4	4	3	
493	4X	1830.322	1	2	3	3	3	3	3	3	3	3	2	3	3	3	3	3	3	3	3	3	3	
494	4X	1431.978	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
495	4X	2005.456	2	2	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2	
		171809.888	134	110	146	134	153	143	135	144	141	147	145	150	150	146	143	152	171	188	177	170	213	189

Table 4. Number of representative tows conducted in each stratum during the period 1992 to 2013.

Stratum	NAFO Div.	Area (km2)	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
440	4VN	3173.016	4	3	4	4	4	4	4	4	6	4	4	4	4	4	4	4	3	4	4	5	4	4
441	4VN	3434.000	5	5	5	5	5	5	6	7	6	6	7	6	7	6	6	5	6	6	7	6	6	6
442	4VN	4934.658	6	5	6	6	6	6	7	6	6	5	6	7	5	5	5	5	5	6	5	6	6	6
443	4VSW	4526.012	4	3	3	4	4	5	5	4	5	4	5	5	5	5	5	5	4	4	4	4	5	5
444	4VSW	13478.450	8	9	6	8	8	7	8	8	9	10	9	9	9	8	10	8	6	9	11	13	9	8
445	4VSW	3512.982	4	5	7	4	4	4	3	3	6	5	5	5	5	6	5	4	3	3	3	4	3	3
446	4VSW	1686.094	3	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	2	3	3	3	4	3
447	4VSW	5549.344	7	7	7	7	6	7	7	6	7	7	7	7	7	7	6	6	4	6	6	8	6	7
448	4VSW	4975.866	6	7	7	7	6	7	6	7	8	8	8	8	7	8	8	6	5	7	7	10	8	8
449	4VSW	494.496	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2	2	4	2	2
450	4VSW	1315.222	3	3	3	3	3	3	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
451	4VSW	504.798	2	2	2	2	4	2	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2
452	4VSW	1184.730	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
453	4VSW	889.406	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3	1	2	2	1	3
454	4VSW	1713.566	2	2	2	2	3	2	2	2	2	2	2	2	2	3	2	2	2	2	2	4	2	2
455	4VSW	7286.948	10	9	10	10	10	13	8	11	11	11	11	11	8	12	11	7	5	8	8	10	10	11
456	4VSW	3279.470	7	8	8	8	8	8	8	6	8	10	8	8	8	8	8	6	2	7	7	9	8	8
457	4VSW	2784.974	2	2	2	2	2	2	2	1	4	2	2	2	2	2	2	2	2	2	2	4	2	2
458	4VSW	2259.572	8	8	8	8	7	8	5	6	10	8	7	8	8	10	8	5	2	7	6	9	8	6
459	4VSW	10810.232	6	4	6	6	4	5	5	6	6	8	6	6	6	6	6	5	3	6	6	7	6	6
460	4VSW	4615.296	3	3	3	3	3	3	3	3	3	3	3	3	3	4	3	3	2	3	3	4	4	3
461	4VSW	3962.836	2	2	2	2	2	2	2	2	2	2	2	2	2	2	4	2	2	2	2	3	3	2
462	4VSW	7266.344	4	4	4	4	4	4	4	4	4	4	4	4	4	5	4	4	3	4	4	4	6	4
463	4VSW	1037.068	2	2	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2	3	2	2
464	4VSW	4453.898	7	7	7	7	7	4	7	7	7	7	7	7	5	8	7	6	4	5	6	7	7	7
465	4VSW	8183.222	10	10	10	10	10	10	9	10	10	10	10	10	10	10	10	7	8	7	8	10	10	10
466	4VSW	776.084	2	2	2	3	2	2	3	2	2	2	2	2	2	2	2	2	1	3	2	2	2	2
470	4X	3159.280	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3
471	4X	3447.736	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3
472	4X	4289.066	4	4	4	4	4	3	4	4	4	4	4	4	4	4	4	4	3	4	3	4	6	4
473	4X	910.010	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
474	4X	552.874	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
475	4X	535.704	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
476	4X	5075.452	4	4	4	4	4	4	4	4	4	4	4	4	5	4	4	4	4	4	4	4	4	4
477	4X	4230.688	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	4	5
478	4X	800.122	2	2	2	3	3	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2
480	4X	2249.270	8	8	8	8	8	8	8	8	7	8	8	8	7	9	8	6	8	8	8	7	8	
481	4X	6438.750	9	9	9	9	7	9	9	9	8	9	8	9	9	6	12	9	7	8	8	8	10	9
482	4X	3578.228	3	3	3	3	3	3	3	3	3	3	3	3	3	2	4	3	3	3	3	4	3	3
483	4X	1826.888	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3	2	2
484	4X	7774.576	3	3	3	3	3	3	3	3	3	3	3	3	3	4	3	3	4	3	3	5	5	5
485	4X	5432.588	3	3	3	3	3	3	3	3	3	4	3	5	5	3	2	5	4	5	5	6	5	5
490	4X	2063.834	4	4	4	5	4	4	4	3	4	4	4	4	6	4	3	3	3	4	3	3	4	2
491	4X	2359.158	3	3	3	3	3	3	3	3	3	3	3	3	3	5	3	3	4	3	4	4	4	4
492	4X	3729.324	3	3	3	2	3	3	3	3	3	3	3	3	5	2	3	4	4	4	4	4	6	4
493	4X	1830.322	3	3	3	3	2	3	3	2	3	3	4	5	2	4	4	3	3	4	3	4	4	4
494	4X	1431.978	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3	3	4	4	4	4
495	4X	2005.456	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	5	3	3	4	3	4	4
171809.888			193	190	195	195	191	193	186	191	213	201	208	216	188	222	209	177	165	196	196	243	210	208

Table 5. Number of representative tows conducted in each stratum during the period 2014 to 2020 and for the whole 1970 to 2020 period.

Stratum	NAFO Div.	Area (km2)	2014	2015	2016	2017	2018	2019	2020	Total
440	4VN	3173.016	4	4	4	4	0	5	4	190
441	4VN	3434.000	6	6	6	6	0	7	4	238
442	4VN	4934.658	6	6	6	6	0	6	5	240
443	4VSW	4526.012	3	7	4	5	0	9	4	214
444	4VSW	13478.450	9	9	11	10	0	6	8	352
445	4VSW	3512.982	3	4	4	4	0	6	3	215
446	4VSW	1686.094	3	2	3	2	0	3	2	145
447	4VSW	5549.344	7	7	7	7	0	6	5	291
448	4VSW	4975.866	8	7	6	6	0	7	4	299
449	4VSW	494.496	2	2	2	2	0	2	2	100
450	4VSW	1315.222	3	3	3	2	0	3	2	144
451	4VSW	504.798	2	2	2	2	0	2	2	104
452	4VSW	1184.730	1	4	3	3	0	3	3	110
453	4VSW	889.406	3	2	2	1	0	2	2	116
454	4VSW	1713.566	2	2	2	2	0	3	2	121
455	4VSW	7286.948	11	9	9	8	0	9	6	429
456	4VSW	3279.470	6	5	6	6	0	6	4	331
457	4VSW	2784.974	2	3	3	3	0	3	2	113
458	4VSW	2259.572	4	5	5	5	0	6	3	269
459	4VSW	10810.232	6	7	7	6	0	9	7	262
460	4VSW	4615.296	3	5	5	5	3	6	5	151
461	4VSW	3962.836	2	3	3	3	2	3	3	113
462	4VSW	7266.344	5	5	5	5	0	5	5	212
463	4VSW	1037.068	2	3	2	2	0	2	2	107
464	4VSW	4453.898	7	6	6	4	0	6	4	288
465	4VSW	8183.222	10	10	9	7	3	10	7	397
466	4VSW	776.084	2	2	2	3	0	3	2	118
470	4X	3159.280	2	3	3	3	4	3	2	112
471	4X	3447.736	2	3	3	3	4	4	3	110
472	4X	4289.066	4	4	4	4	4	4	4	172
473	4X	910.010	2	2	2	2	2	2	2	104
474	4X	552.874	2	2	2	2	2	2	2	100
475	4X	535.704	2	2	2	2	2	2	2	103
476	4X	5075.452	4	5	5	5	5	5	5	177
477	4X	4230.688	6	5	5	4	4	6	4	204
478	4X	800.122	2	2	2	3	2	2	2	119
480	4X	2249.270	6	7	7	7	5	7	5	306
481	4X	6438.750	9	8	10	9	6	9	6	350
482	4X	3578.228	3	3	4	4	3	4	3	141
483	4X	1826.888	2	2	3	3	2	3	2	105
484	4X	7774.576	4	6	5	7	7	7	7	186
485	4X	5432.588	5	6	6	6	4	6	5	196
490	4X	2063.834	3	4	4	4	3	4	3	173
491	4X	2359.158	4	4	4	4	3	4	3	168
492	4X	3729.324	4	3	4	4	3	4	4	171
493	4X	1830.322	3	3	4	6	3	3	3	159
494	4X	1431.978	3	4	4	3	2	4	3	128
495	4X	2005.456	2	4	4	4	3	4	3	127
		171809.888	196	212	214	208	81	227	175	9080

405 A total of 9080 representative tows were conducted for the period spanning from 1970 to 2020.

406

4 Discussion

407 This report builds on previous work and former atlases by updating a comprehensive suite of
408 indices to give a snapshot of population status and environmental preferences of 104 fish and
409 invertebrate species. The current document is not meant to replace stock assessments, species-
410 specific analyses of abundance, biomass and distribution, or any targeted attempts to integrate
411 information about species or group of species from the wide and disparate sources of data about
412 marine organisms in the area covered by the DFO Maritimes summer trawl survey. It is rather
413 meant to provide a reproducible set of tools to extract and visualize the information collected
414 in the summer groundfish research vessel survey. It is hoped that this document can provide a
415 stepping stone to conduct other ecological analyses using the trawl survey data and increase
416 reproducibility and transparency of ecological information collected annually.

417 **4.1 Diversity of approaches used for mapping fish and invertebrates in the Scotian Shelf
418 bioregion**

419 Different methods have been applied in the Northwest Atlantic, and specifically on the Scotian
420 Shelf bioregion, to map fish and invertebrate species distribution. The present report, for
421 example, builds upon the atlas of important habitat developed to map the persistence of relatively
422 high biomass for key fish species using the summer groundfish research vessel survey (Horsman
423 and Shackell 2009). Important habitat was obtained by interpolating observed weight per each
424 species using the IDW, and calculating areas with relatively persistent high biomass for periods
425 representing different fishery management eras. To compliment information from this atlas,
426 including additional representations of biomass and diversity, a similar IDW interpolation mapping
427 procedure was followed by Smith et al. (2015), Ward-Paige and Bundy (2015), and Bundy et al.
428 (2017). The summer groundfish research vessel survey is typically conducted during the month
429 of July. However, from the fall of 1978 through to the spring of 1985, DFO also conducted spring
430 and fall surveys using the same sampling design. This unique seasonal data was used to map
431 the seasonal spatial distribution of key demersal and other fish species using IDW interpolation
432 on the Scotian Shelf from the spring, summer and fall between 1978 and 1985 (Smith et al.
433 2015). Following recommendations provided by Kenchington and Kenchington (2017), the spatial
434 distribution of three indicators of biodiversity for fish and invertebrates were mapped using IDW
435 interpolation to identify areas with persistently high values across fishery management eras,
436 and compared with areas of persistently high abundance for selected species (Ward-Paige and
437 Bundy 2015). This analysis revealed a lack of consistent relationships between areas of persist
438 high diversity and persistent high biomass, suggesting that both can be used as independent
439 and important spatial indicators of the system (Ward-Paige and Bundy 2015). Groupings of
440 fishes and invertebrates based on size, habitat and feeding guild, were also mapped using
441 IDW interpolations to identify hotspots of functional group diversity (Bundy et al. 2017). This
442 analysis revealed a spatially and temporally variable distribution of functional diversity across
443 the Scotian Shelf with notable areas of high and low diversity (Bundy et al. 2017). Top quintiles
444 of each functional group using the IDW approach were used as representative layers for fish

445 and invertebrates in the MPA Network design in the Scotian Shelf Bioregion (Serdynska et al. In
446 press). IDW interpolation methods have also been used to map the distribution of individual
447 species such as sea cucumbers (*Cucumaria frondosa*) in the Scotian Shelf bioregion (Shackell et
448 al. 2013a), and sea scallop (*Placopecten magellanicus*) in Georges and Browns Bank (Shackell
449 et al. 2013b).

450 Species Distribution Modelling (SDM), instead of IDW, can also be used to evaluate spatio-
451 temporal dynamics by predicting and understanding past, present and future distribution
452 of species using environmental predictors (Robinson et al. 2017). A variety of modelling
453 approaches are being implemented in Maritimes Region to map and predict fish and invertebrate
454 species distribution by incorporating environmental predictors to account for seasonal and
455 temporal variability. For example, a stock assessment of snow crab (*Chionoecetes opilio*) on
456 the Scotian Shelf used data from the snow crab survey from 2005 to 2018 to map spatial data
457 products for this stock, including annual predicted interpolations of potential habitat using
458 Generalized Additive Models (GAM) and several environmental covariates including depth,
459 curvature, slope, species composition, and annual temperature (Zisserson et al. 2019). Sea
460 scallop predicted habitat using Maximum Entropy (MaxEnt) models were computed in the
461 German Bank using data compiled via benthic habitat mapping and seafloor geotechnical
462 surveys in 2006, 2009, and 2010 (Brown et al. 2012). Predictions in the Scotian Shelf bioregion
463 and the Northeast United States using datasets from DFO and the National Oceanic and
464 Atmospheric Administration from 1993 to 2012 also predicted sea scallop habitat at a wider scale
465 based on three scenarios of seasonal temperature and salinity climatologies (NOAA) (Lowen
466 et al. 2019). Offshore American lobster stock assessments (*Homarus americanus*) used data
467 from the RV, DFO Georges Bank, and National Marine Fisheries Service (NMFS) Northeast
468 Fisheries Science Center (NEFSC) bottom trawl surveys (1970 to 2015) to predict species
469 distribution using boosted regression trees and several environmental predictors (bathymetry,
470 slope, curvature, and annual temperature interpolations) (Cook et al. 2017). Information on
471 the potential for recovery of cusk (*Brosme brosme*) used data from the bottom longline Halibut
472 industry survey and Cusk absences in the Summer groundfish research vessel survey from
473 1998-2013 to predict suitable habitat using GAM, MaxEnt, and random forest models and
474 several physical environmental variables (e.g. complexity, benthic current stress and complexity,
475 temperature, salinity, primary production, chlorophyll, suspended matter) (Harris et al. 2018).
476 Atlantic halibut (*Hippoglossus hippoglossus*) assessments using Summer groundfish research
477 vessel survey and NOAA survey data from 2001 to 2013 predicted juvenile habitat using MaxEnt
478 model and environmental predictors (bathymetry, slope, bottom temperature) (French et al.
479 2018). Persistent areas of high Atlantic halibut juvenile abundance were predicted using data
480 from 27 bottom trawl surveys combined (NMFS and DFO) from 1978 to 2013 and applying
481 Bayesian hierarchical spatiotemporal models with two environmental predictors (depth and
482 temperature) (Boudreau et al. 2017).

483 These examples of mapping efforts in Maritimes Region showcase the diversity of approaches
484 relevant to a variety of important research questions and management applications. Approaches,
485 methods, datasets, and environmental predictors are selected based on individual project
486 research questions, and considerations for each species, communities or stock. This allows
487 research groups to maintain innovation and keep up with emerging methods and technologies to
488 improve assessments, predictions, and ultimately, science advice. The diversity of approaches
489 also leads to complexity when looking across studies as each data compilation and predictive
490 method carries its own independent assumptions and can lead to different spatial outputs.

491 **4.2 Interpreting spatial results for marine spatial planning purposes**

492 Fisheries and Oceans Canada is leading a marine spatial planning process that brings together
493 relevant authorities and stakeholders to better coordinate how we use and manage marine
494 spaces to achieve ecological, economic and social objectives. Operationalizing marine spatial
495 planning includes a series of steps, including the process of analyzing existing conditions
496 by collecting and mapping information about ecological, environmental and oceanographic
497 conditions (Ehler and Douvere 2009; Agardy et al. 2011). Mapping the distribution of species
498 is critical for the implementation of spatial management and as a first step in marine spatial
499 planning processes. Species distribution have supported the identification of important sites for
500 a given species or areas of high richness and diversity, which in turn can be used to inform siting
501 decisions of new activities such as Marine Protected Areas (MPA), aquaculture sites or wind
502 turbines. In the Scotian Shelf bioregion, mapping species distributions has been used to highlight
503 areas of high biological diversity to support the identification of Ecologically or Biologically
504 Significant Areas [Ricard and Shackell (2013); Ward-Paige and Bundy (2015)], to distinguish
505 important and persistent habitat of significant species and functional groups to support MPA and
506 conservation planning (Horsman and Shackell 2009; Smith et al. 2015; Ward-Paige and Bundy
507 2015; Bundy et al. 2017), to identify important habitat for Species at Risk (Harris et al. 2018) and
508 to highlight reserves for data-poor invertebrate fisheries (Shackell et al. 2013a). Mapping species
509 distribution has also been used to illustrate multi-decadal scale projections of changes in species
510 distribution in the context of climate change and adaption (Stanley et al. 2018; W. et al. 2019).

511 In support of the marine spatial planning process, a public web-based atlas with relevant
512 geospatial information is being developed to support decision-making. This Atlantic Canada-
513 wide compilation of data and information will be a web-based, public platform with interactive
514 maps of ocean ecosystems, human uses and management areas. This atlas cannot host the
515 vast diversity of products and mapping approaches available in Maritimes Region. Consequently,
516 we recommend that data products presented in this report should not be used for the atlas until
517 an evaluation of the spatial information available and used in the past, is conducted.

518 This diverse portfolio of approaches and applications is not unique to the Maritimes Region. A
519 recent review of global distribution modelling efforts recommended the adoption of a consistent
520 framework that integrates multi-model approaches and a clear expression of errors and
521 uncertainties (Robinson et al. 2017). In this context, Pacific Region has developed two initiatives
522 to enable consistency and frequent publication, reproducibility, and transparency. One initiative
523 developed a fully automated reproducible report to give a synthesis of data availability, population
524 trends, fishing trends, growth and maturity patterns for 113 groundfish species in British
525 Columbia to support stock assessment (Anderson et al. 2019). The second initiative developed
526 a SDM framework that was applied to twelve species on Canada's Pacific coast as part of the
527 Regional Response Plan (Nephin et al. 2019). The Maritimes and Gulf region, through this and
528 past reports, are also using similar reproducible approaches to facilitate annual updates and
529 transparency (Ricard and Shackell 2013; Ricard and Gomez 2021).

530 Recognizing the diversity of approaches for mapping fish and invertebrates in the Scotian Shelf
531 bioregion, we recommend the development of a regional community of practice to compare
532 and evaluate approaches for mapping, interpolating and/or modelling fish and invertebrates
533 so future publications and advice related to spatial outputs can lead to more comparable work
534 and consistent science advice to support processes such as marine spatial planning. At the

535 international level, guidelines and standards related to appropriate variables and methods
536 for mapping and modeling species and communities of deep-sea habitats were proposed to
537 encourage the production of publications that will lead to more comparable work (Kenchington et
538 al. 2019). Similar general guidance for group practice approach mapping would be a worthwhile
539 product in Maritimes Region. Until then, we proposed the use of the Open Data record (DFO
540 2021) for the version 1.0 of the public web-based atlas.

541

5 Acknowledgements

542 We thank all the dedicated personnel involved in running trawl surveys in the Maritimes Region.
543 We thank the numerous colleagues in Maritimes Region that have shared information and advice
544 in support of this report.

6 References

- 546 Agardy, T., Notarbartolo di Sciara, G., and Christie, P. 2011. Addressing the shortcomings of
 548 marine protected areas through large scale marine spatial planning. *Marine Policy* 35(2):
 549 226–232.
- 550 Anderson, S.C., Grandin, C., Edwards, A.M., Grinnell, M.H., Ricard, D., and Haigh, R. (In
 551 press). Csasdown: Reproducible CSAS Reports with Bookdown.
- 552 Anderson, S.C., Keppel, E.A., and Edwards, A.M. 2019. *A reproducible data synopsis for*
 553 *over 100 species of British Columbia groundfish*. DFO Can. Sci. Advis. Sec. Res. Doc.
 554 2019/041. vii + 321 p.
- 555 Appeltans, W., Bouchet, P., Boxshall, G.A., De Broyer, C., Voogd, N.J. de, Gordon, D.P.,
 556 Hoeksema, B.W., Horton, T., Kennedy, M., J., M., Poore, G.C.B., Read, G., Stöhr,
 557 S., Walter, T.C., and Costello, M.J. (*Editors*). 2012. World register of marine species.
 558 Accessed at <http://www.marinespecies.org> on 2020-12-1.
- 559 Baddeley, R., A. 2015. Spatial point patterns: Methodology and applications
 560 with R. Chapman; Hall/CRC Press, London [http://www.crcpress.com/
 561 Spatial-Point-Patterns-Methodology-and-Applications-with-R/
 562 Baddeley-Rubak-Turner/9781482210200/](http://www.crcpress.com/Spatial-Point-Patterns-Methodology-and-Applications-with-R-Baddeley-Rubak-Turner/9781482210200/).
- 563 Benoît, H.P., Abgrall, M.-J., and Swain, D.P. 2003. An assessment of the general status of
 564 marine and diadromous fish species in the southern Gulf of St. Lawrence based on annual
 565 bottom trawl surveys (1971-2002). *Can. Tech. Rep. Fish. Aquat. Sci.* 2472: iv + 183 p.
- 566 Bivand, R. 2020. classInt: Choose univariate class intervals. R package version 0.4-3 <https://CRAN.R-project.org/package=classInt>.
- 567 Bivand, R., and Lewin-Koh, N. 2020. Maptools: Tools for handling spatial objects. R package
 568 version 1.0-2 <https://CRAN.R-project.org/package=maptools>.
- 569 Bivand, R., and Rundel, C. 2020. Rgeos: Interface to geometry engine - open source
 ('GEOS'). R package version 0.5-5 <http://CRAN.R-project.org/package=rgeos>.
- 570 Boudreau, S.A., Shackell, N.L., Carson, S., and Heyer C. E., den. 2017. *Connectivity,*
 571 *persistence, and loss of high abundance areas of a recovering marine fish population*
 572 *in the Northwest Atlantic Ocean*. *Ecol. Evol.* 7: 9739–9749.
- 573 Bourdages, H., and Ouellet, J.-F. 2012. *Geographic distribution and abundance indices of*
 574 *marine fish in the northern Gulf of St. Lawrence (1990-2009)*. *Can. Tech. Rep. Fish.*
 575 *Aquat. Sci.* 2963: vi + 171 p.
- 576 Brown, C., Sameoto, J., and Smith, S. 2012. *Multiple methods, maps, and management*
 577 *applications: Purpose made seafloor maps in support of ocean management*. *Journal*
 578 *of Sea research* 72: 113.
- 579 Bundy, A., Will, E., Serdynska, A., Cook, A., and Ward-Paige, C.A. 2017. *Defining and*
 580 *mapping functional groups for fishes and invertebrates in the Scotian Shelf bioregion*.
 581 *Can. Tech. Rep. Fish. Aquat. Sci.* 3186: iv + 49 p.

- 584 Clark, D.W., and Emberley, J. 2011. Update of the 2010 summer scotian shelf and bay of
585 fundy research vessel survey. Can. Tech. Rep. Fish. Aquat. Sci.: 1238: ix + 98 p.
- 586 Cook, A.M., Cassista Da-Ros, M., and Denton, C. 2017. [Framework assessment of the](#)
587 [offshore American Lobster in Lobster Fishing Area \(LFA\) 41](#). ICES Journal of Marine
588 Science 2017/065 viii + 186 p.
- 589 DFO. 2016. DFO maritimes research vessel trawl surveys invertebrate observations. Version
590 7 in OBIS canada digital collections. Bedford Institute of Oceanography, Dartmouth, NS,
591 Canada, Published by OBIS, Digital 2016.
- 592 DFO. 2021. Maritimes research vessel surveys [dataset]. Retrieved from
593 <https://open.canada.ca/data/en/dataset/8ddcaeeaa-b806-4958-a79f-ba9ab645f53b>.
- 594 Doubleday, W.G., and Rivard, D. 1981. Bottom trawl surveys. Can. Spec. Publ. Fish. Aquat.
595 Sci: 58: 237 pp.
- 596 Ehler, C., and Douvere, F. 2009. Marine spatial planning: A step-by-step approach.
- 597 French, K., Shackell, N., and Heyer, N. den. 2018. [Information on the potential for recovery of](#)
598 [cusk in canadian waters](#). Fish. Bull. 116: 107–121.
- 599 Harris, L.E., Greenlaw, M., McCurdy, D., and MacDonald, D. 2018. [Information on the](#)
600 [potential for recovery of cusk in canadian waters](#). DFO Can. Sci. Advis. Sec.Res. Doc.
601 2018/002. vi + 62 p.
- 602 Holstein, J. 2018. Worms: Retriving aphia information from world register of marine species.
603 R package version 0.2.2 <https://CRAN.R-project.org/package=worms>.
- 604 Horsman, T.L., and Shackell, N.L. 2009. [Atlas of important habitat for key fish species of the](#)
605 [Scotian Shelf, Canada](#). Can. Tech. Rep. Fish. Aquat. Sci. 2835: viii + 82 p.
- 606 Kenchington, E., Callery, O., Davidson, F., Grehan, A., Morato, T., Appiott, J., Davis, A.,
607 Dunstan, P., Du Preez, C., Finney, J., González-Irusta, J.M., Howell, K., Knudby, A.,
608 Lacharité, M., Lee, J., Murillo, F.J., Beazley, L., Roberts, J.M., Roberts, M., Rooper, C.,
609 Rowden, A., Rubidge, E., Stanley, R., Stirling, D., Tanaka, K.R., Vanhatalo, J., Weigel, B.,
610 Woolley, S., and Yesson, C. 2019. [Use of species distribution modeling in the deep sea](#).
611 Can. Tech. Rep. Fish. Aquat. Sci. 3296: ix + 76 p.
- 612 Kenchington, T.J., and Kenchington, E.L.R. 2017. [Biodiversity metrics for use in the](#)
613 [ecosystem approach to oceans management](#). Can. Tech. Rep. Fish. Aquat. Sci. 3186:
614 iv + 49 p.
- 615 Lohr, S. 1999. Sampling: Design and analysis. Pacific Grove, CA: Brooks/Cole Publishing
616 Company.
- 617 Losier, R.J., and Waite, L.E. 1989. Systematic listing of scientific and/or common names of
618 invertebrates and marine plants and their respective codes used by marine fish division,
619 Fisheries and Oceans, Scotia-Fundy Region. Canadian Data Report of Fisheries and
620 Aquatic Sciences (721).
- 621 Lowen, B., Hart, D., Stanley, R., Lehnert, S., Bradbury, I., and C., D. 2019. [Assessing effects](#)
622 [of genetic, environmental, and biotic gradients in species distribution modelling](#). ICES

- Journal of Marine Science 76(6): 1762–1775.

Myers, R.A., and Stokes, K. 1989. Density-dependent habitat utilization of groundfish and the improvement of research surveys. (D:15). International Council for the Exploration of the Sea Council Meeting.

Nelson J. S., Crossman E. J., Espinosa-Perez H., Findley L. T., Gilbert C. R., Lea R. N., and D., W.J. 2004. Common and scientific names of fishes from the united states canada and mexico. Number 29 in Special Publication. American Fisheries Society, Bethesda, Maryland, sixth edition.

Nepchin, J., Gregr, E.J., St. Germain, C., Fields, C., and Finney, J.L. and. 2019. [Development of a Species Distribution Modelling framework and its application to twelve species on Canada's Pacific coast](#). DFO Can. Sci. Advis. Sec. Res. Doc. 2020/004. xii + 107 p.

Neuwirth, E. 2014. RColorBrewer:ColorBrewer palettes. R package version 1.1-2 <https://CRAN.R-project.org/package=RColorBrewer>.

Page L. M., Espinosa-Pérez H., Findley L. T., Gilbert C. R., Lea R. N., Mandrak N. E., and M.R.L., and S., N.J. 2013. Common and scientific names of fishes from the United States, Canada, and Mexico. American Fisheries Society.

Pebesma, E. 2004. Multivariable geostatistics in S: The gstat package. In Computers and Geosciences.

Perry, R.I., and Smith, S.J. 1994. Identifying habitat associations of marine fishes using survey data: An application to the northwest atlantic. Canadian Journal of Fisheries and Aquatic Sciences (51(3)): 589–602.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Ricard, D., and Gomez, C. 2021. Maritimes-SUMMER-atlas. <https://github.com/dfo-gulf-science/Maritimes-SUMMER-Atlas>; GitHub.

Ricard, D., and Shackell, N.L. 2013. [Population status \(abundance/biomass, geographic extent, body size and condition\), important habitat, depth, temperature and salinity of marine fish and invertebrates on the Scotian Shelf and Bay of Fundy \(1970-2012\)](#). Can. Tech. Rep. Fish. Aquat. Sci. 3012: viii + 180 p.

Ripley, B., and Lapsley, M. 2019. RODBC: ODBC database access. R package version 1.3-16 <http://CRAN.R-project.org/package=RODBC>.

Ripley, B., Venables, B., Bates, D., Hornik, K., Gebhardt, A., and Firth, D. 2020. Modern applied statistics with s. R package version 7.3-53 <https://cran.r-project.org/web/packages/MASS/index.html>.

Robinson, N.M., Nelson, W.A., Costello, M.J., Sutherland, J.E., and Lundquist, C.J. 2017. Systematic review of marine-based species distribution models (SDMs) with recommendations for best practice. Front. Mar. Sci (4): 421.

Schnute, J.T., Boers, N., and Haigh, R. 2019. PBSmapping: Mapping fisheries data and spatial analysis tools. R package version 2.72.1 <https://cran.r-project.org/web/packages/PBSmapping/index.html>.

- 662 packages/PBSmapping/index.html.
- 663 Shackell, N., Brickman, D., and Frank, K. 2013b. [Georges Bank 'a' and Browns Bank 'North'](#)
664 [Scallop stock assessment](#). Aquatic Conserv: Mar. Freshw. Ecosyst. 23: 723–731.
- 665 Shackell, N., Brickman, D., and Frank, K. 2013a. [Reserve site selection for data-poor](#)
666 [invertebrate fisheries using patch scale and dispersal dynamics: A case study of sea](#)
667 [cucumber](#). Aquatic Conserv: Mar. Freshw. Ecosyst. 23: 723–731.
- 668 Simon, J.E., and Comeau, P.A. 1994. [Summer distribution and abundance trends of species](#)
669 [caught on the Scotian Shelf from 1970-92, by the research vessel groundfish survey](#). Can.
670 Tech. Rep. Fish. Aquat. Sci. 1953.
- 671 Smith, C.D., Serdynska, A.R., King, M.C., and Shackell, N.L. 2015. [Spring, summer and fall](#)
672 [distribution of common demersal fishes on the scotian shelf between 1978 and 1985](#). Can.
673 Tech. Rep. Fish. Aquat. Sci. 3068: vi + 38 p.
- 674 Smith, S.J. 1996. Assessment of groundfish stocks based on bottom trawl survey results.
675 NAFO Scientific Council Studies 28: 25–53.
- 676 Stanley, R.E., DiBacco, C., Lowen, B., Beiko, R., Jeffery, N., Wyngaarden, M., Bentzen, P.,
677 Brickman, D., Benestan, L., Bernatchez, L., Johnson, C., Snelgrove, P., Wang, Z., and
678 Wringe, I., B. Bradbury. 2018. A climate-associated multispecies cryptic cline in the
679 northwest Atlantic. Science Advances 4(3): 1–7.
- 680 Swain, D.P., and Morin, R. 1996. Relationships between geographic distribution and
681 abundance of American plaice (*Hippoglossoides platessoides*) in the southern Gulf of
682 St. Lawrence. Canadian Journal of Fisheries and Aquatic Sciences 53(1): 106–119.
- 683 Swain, D.P., and Sinclair, A.F. 1994. Fish distribution and catchability: What is the appropriate
684 measure of distribution? Canadian Journal of Fisheries and Aquatic Sciences 51(5):
685 1046–1054.
- 686 Tremblay M. J., B.R., Black G. A. P. 2007. The distribution of common decapod crustaceans
687 and other invertebrates recorded in annual ecosystem surveys of the scotian shelf 1999-
688 2006, by the research vessel groundfish survey. Can. Tech. Rep. Fish. Aquat. Sci. 74.
- 689 W., G.B.J., Shackell, N.L., Ferguson, K., P., G., Cogswell, A., Brickman, D., Wang, Z., Cook,
690 A., Brennan, C.E., and Saba, V.S. 2019. Climate change vulnerability of American lobster
691 fishing communities in Atlantic Canada. Frontiers in Marine Science 6: 579.
- 692 Ward-Paige, C.A., and Bundy, A. 2015. [Mapping biodiversity on the Scotian Shelf and in the](#)
693 [Bay of fundy](#). Can. Tech. Rep. Fish. Aquat. Sci. 3068: vi + 38 p.
- 694 Wickham, H. 2019. Tidyverse: Easily install and load the 'tidyverse'. R package version 1.3.0
695 <https://cran.r-project.org/web/packages/tidyverse/index.html>.
- 696 Zisserson, B.M., Cameron, B.J., Glass, A.C., and Choi, J.S. 2019. [assessment of scotian shelf](#)
697 [snow crab in 2017](#). DFO Can. Sci. Advis. Sec. Res. Doc. 2018/051. ix + 147 p.

7 Appendix

699

7.1 Atlantic cod (*Morue franche*) - species code 10 (category LF)

700

Scientific name: [Gadus morhua](#)

701




Figure 7.1A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic cod.

702




Figure 7.1B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic cod.




Figure 7.1C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic cod.






Figure 7.1D. Average fish condition in NAFO units 4X and 4VW for Atlantic cod.



Freq	Depth	Temp	Sal
F5	26	1.2	31.00
F25	43	3.5	32.47
F50	60	6.0	33.27
F75	82	8.1	34.40
F95	126	10.0	35.03

Figure 7.1E. Catch distribution by depth, temperature and salinity of Atlantic cod.




Figure 7.1F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic cod.

707

7.2 Haddock (Aiglefin) - species code 11 (category LF)

708

Scientific name: [Melanogrammus aeglefinus](#)

709




Figure 7.2A. Inverse distance weighted distribution of catch biomass (kg/tow) for Haddock.

710




Figure 7.2B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Haddock.




Figure 7.2C. Length frequency distribution in NAFO units 4X and 4VW for Haddock.






Figure 7.2D. Average fish condition in NAFO units 4X and 4VW for Haddock.



713



Freq	Depth	Temp	Sal
F5	27	1.1	31.00
F25	49	3.0	32.45
F50	66	5.5	33.14
F75	87	7.9	34.36
F95	127	10.0	35.03

Figure 7.2E. Catch distribution by depth, temperature and salinity of Haddock.




Figure 7.2F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Haddock.

715

7.3 White hake (Merluche blanche) - species code 12 (category LF)

716

Scientific name: [Urophycis tenuis](#)

717




Figure 7.3A. Inverse distance weighted distribution of catch biomass (kg/tow) for White hake.

718




Figure 7.3B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of White hake.




Figure 7.3C. Length frequency distribution in NAFO units 4X and 4VW for White hake.






Figure 7.3D. Average fish condition in NAFO units 4X and 4VW for White hake.



721



Freq	Depth	Temp	Sal
F5	40	1.1	31.00
F25	70	3.0	32.46
F50	89	5.5	33.20
F75	115	7.9	34.39
F95	163	10.0	35.04

Figure 7.3E. Catch distribution by depth, temperature and salinity of White hake.

722




Figure 7.3F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for White hake.

723

7.4 Red hake (Merluche écureuil) - species code 13 (category LF)

724

Scientific name: [Urophycis chuss](#)

725




Figure 7.4A. Inverse distance weighted distribution of catch biomass (kg/tow) for Red hake.

726




Figure 7.4B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Red hake.




Figure 7.4C. Length frequency distribution in NAFO units 4X and 4VW for Red hake.






Figure 7.4D. Average fish condition in NAFO units 4X and 4VW for Red hake.



729



Freq	Depth	Temp	Sal
F5	35	1.1	31.00
F25	62	2.9	32.43
F50	82	5.4	33.12
F75	103	7.9	34.32
F95	130	10.0	35.08

Figure 7.4E. Catch distribution by depth, temperature and salinity of Red hake.

730




Figure 7.4F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Red hake.

731

7.5 Silver hake (*Merlu argenté*) - species code 14 (category LF)

732

Scientific name: [Merluccius bilinearis](#)

733




Figure 7.5A. Inverse distance weighted distribution of catch biomass (kg/tow) for Silver hake.

734




Figure 7.5B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Silver hake.




Figure 7.5C. Length frequency distribution in NAFO units 4X and 4VW for Silver hake.






Figure 7.5D. Average fish condition in NAFO units 4X and 4VW for Silver hake.




737



Freq	Depth	Temp	Sal
F5	28	1.2	31.00
F25	55	3.1	32.46
F50	77	5.5	33.20
F75	104	8.0	34.37
F95	137	10.0	35.07

Figure 7.5E. Catch distribution by depth, temperature and salinity of Silver hake.



738

Figure 7.5F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Silver hake.

739

7.6 Pollock (Goberge) - species code 16 (category LF)

740

Scientific name: [Pollachius virens](#)

741




Figure 7.6A. Inverse distance weighted distribution of catch biomass (kg/tow) for Pollock.

742




Figure 7.6B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Pollock.




Figure 7.6C. Length frequency distribution in NAFO units 4X and 4VW for Pollock.






Figure 7.6D. Average fish condition in NAFO units 4X and 4VW for Pollock.



745



Freq	Depth	Temp	Sal
F5	37	1.1	31.00
F25	60	2.8	32.45
F50	82	5.3	33.14
F75	108	7.7	34.33
F95	137	10.0	35.03

Figure 7.6E. Catch distribution by depth, temperature and salinity of Pollock.

746




Figure 7.6F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Pollock.

747

7.7 Atlantic redfishes (Sébastes de l'Atlantique) - species code 23 (category LF)

748

Scientific name: [Sebastes](#)

749




Figure 7.7A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic redfishes.

750




Figure 7.7B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic redfishes.




Figure 7.7C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic redfishes.






Figure 7.7D. Average fish condition in NAFO units 4X and 4VW for Atlantic redfishes.



753



Freq	Depth	Temp	Sal
F5	47	1.2	31.00
F25	68	3.4	32.48
F50	86	5.9	33.29
F75	114	8.1	34.41
F95	166	10.0	35.05

Figure 7.7E. Catch distribution by depth, temperature and salinity of Atlantic redfishes.




Figure 7.7F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic redfishes.

755

7.8 Atlantic halibut (Flétan de l'Atlantique) - species code 30 (category LF)

756

Scientific name: [Hippoglossus hippoglossus](#)

757




Figure 7.8A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic halibut.

758




Figure 7.8B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic halibut.




Figure 7.8C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic halibut.






Figure 7.8D. Average fish condition in NAFO units 4X and 4VW for Atlantic halibut.



761



Freq	Depth	Temp	Sal
F5	31	1.3	31.00
F25	49	3.2	32.45
F50	60	5.8	33.16
F75	75	8.3	34.34
F95	122	10.0	35.08

Figure 7.8E. Catch distribution by depth, temperature and salinity of Atlantic halibut.

762




Figure 7.8F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic halibut.

763

7.9 American plaice (*Ple canadienne*) - species code 40 (category LF)

764

Scientific name: [Hippoglossoides platessoides](#)

765




Figure 7.9A. Inverse distance weighted distribution of catch biomass (kg/tow) for American plaice.

766




Figure 7.9B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of American plaice.




Figure 7.9C. Length frequency distribution in NAFO units 4X and 4VW for American plaice.






Figure 7.9D. Average fish condition in NAFO units 4X and 4VW for American plaice.



769



Freq	Depth	Temp	Sal
F5	29	1.3	31.00
F25	46	3.7	32.48
F50	67	6.1	33.27
F75	87	8.1	34.41
F95	133	10.0	35.05

Figure 7.9E. Catch distribution by depth, temperature and salinity of American plaice.




Figure 7.9F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for American plaice.

771

7.10 Witch flounder (*Ple grise*) - species code 41 (category LF)

772

Scientific name: [Glyptocephalus cynoglossus](#)

773




Figure 7.10A. Inverse distance weighted distribution of catch biomass (kg/tow) for Witch flounder.

774




Figure 7.10B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Witch flounder.




Figure 7.10C. Length frequency distribution in NAFO units 4X and 4VW for Witch flounder.






Figure 7.10D. Average fish condition in NAFO units 4X and 4VW for Witch flounder.




777



Freq	Depth	Temp	Sal
F5	32	1.3	31.00
F25	57	3.5	32.49
F50	77	6.1	33.30
F75	102	8.2	34.45
F95	152	10.0	35.06

Figure 7.10E. Catch distribution by depth, temperature and salinity of Witch flounder.



778

Figure 7.10F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Witch flounder.

779

7.11 Yellowtail flounder (Limande à queue jaune) - species code 42 (category LF)

780

Scientific name: [Limanda ferruginea](#)

781




Figure 7.11A. Inverse distance weighted distribution of catch biomass (kg/tow) for Yellowtail flounder.

782




Figure 7.11B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Yellowtail flounder.




Figure 7.11C. Length frequency distribution in NAFO units 4X and 4VW for Yellowtail flounder.






Figure 7.11D. Average fish condition in NAFO units 4X and 4VW for Yellowtail flounder.



785



Freq	Depth	Temp	Sal
F5	22	1.4	31.00
F25	32	3.7	32.52
F50	43	6.3	33.25
F75	56	8.2	34.41
F95	81	10.0	35.06

Figure 7.11E. Catch distribution by depth, temperature and salinity of Yellowtail flounder.

786




Figure 7.11F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Yellowtail flounder.

787

7.12 Winter flounder (Limande-plie rouge) - species code 43 (category LF)

788

Scientific name: [Pseudopleuronectes americanus](#)

789




Figure 7.12A. Inverse distance weighted distribution of catch biomass (kg/tow) for Winter flounder.

790




Figure 7.12B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Winter flounder.




Figure 7.12C. Length frequency distribution in NAFO units 4X and 4VW for Winter flounder.






Figure 7.12D. Average fish condition in NAFO units 4X and 4VW for Winter flounder.



793



Freq	Depth	Temp	Sal
F5	22	1.1	31.00
F25	31	3.0	32.48
F50	42	5.5	33.17
F75	54	8.3	34.47
F95	84	10.0	35.10

Figure 7.12E. Catch distribution by depth, temperature and salinity of Winter flounder.

794




Figure 7.12F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Winter flounder.

795

7.13 Atlantic wolffish (*Loup atlantique*) - species code 50 (category LF)

796

Scientific name: [Anarhichas lupus](#)

797




Figure 7.13A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic wolffish.

798




Figure 7.13B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic wolffish.




Figure 7.13C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic wolffish.




Figure 7.13D. Average fish condition in NAFO units 4X and 4VW for Atlantic wolffish.





Figure 7.13E. Catch distribution by depth, temperature and salinity of Atlantic wolffish.




Figure 7.13F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic wolffish.

803 **7.14 Atlantic herring (Hareng de l'Atlantique) - species code 60 (category LF)**

804 Scientific name: [Clupea harengus](#)

805




Figure 7.14A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic herring.

806




Figure 7.14B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic herring.




Figure 7.14C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic herring.






Figure 7.14D. Average fish condition in NAFO units 4X and 4VW for Atlantic herring.



809



Freq	Depth	Temp	Sal
F5	32	1.2	31.00
F25	54	3.2	32.45
F50	74	5.6	33.22
F75	95	8.3	34.38
F95	132	10.0	35.10

Figure 7.14E. Catch distribution by depth, temperature and salinity of Atlantic herring.




Figure 7.14F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic herring.

811 **7.15 Longhorn sculpin (Chaboisseau à dix-huit épines) - species code 300 (category**
 812 **LF)**

813 Scientific name: [Myoxocephalus octodecemspiniferus](#)

814




Figure 7.15A. Inverse distance weighted distribution of catch biomass (kg/tow) for Longhorn sculpin.

815




Figure 7.15B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Longhorn sculpin.




Figure 7.15C. Length frequency distribution in NAFO units 4X and 4VW for Longhorn sculpin.






Figure 7.15D. Average fish condition in NAFO units 4X and 4VW for Longhorn sculpin.



818



Freq	Depth	Temp	Sal
F5	23	1.2	31.00
F25	33	3.3	32.51
F50	48	5.8	33.29
F75	64	8.1	34.38
F95	96	10.0	35.05

Figure 7.15E. Catch distribution by depth, temperature and salinity of Longhorn sculpin.

819




Figure 7.15F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Longhorn sculpin.

820

7.16 Moustache sculpin (Faux-trigle armé) - species code 304 (category LF)

821

Scientific name: [Triglops murrayi](#)

822




Figure 7.16A. Inverse distance weighted distribution of catch biomass (kg/tow) for Moustache sculpin.

823




Figure 7.16B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Moustache sculpin.




Figure 7.16C. Length frequency distribution in NAFO units 4X and 4VW for Moustache sculpin.






Figure 7.16D. Average fish condition in NAFO units 4X and 4VW for Moustache sculpin.



826



Freq	Depth	Temp	Sal
F5	30	1.5	31.00
F25	40	4.0	32.48
F50	50	6.5	33.31
F75	63	8.2	34.39
F95	88	10.0	35.06

Figure 7.16E. Catch distribution by depth, temperature and salinity of Moustache sculpin.

827




Figure 7.16F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Moustache sculpin.

828

7.17 Sea raven (Hémithriptère atlantique) - species code 320 (category LF)

829

Scientific name: [Hemitripterus americanus](#)

830




Figure 7.17A. Inverse distance weighted distribution of catch biomass (kg/tow) for Sea raven.

831




Figure 7.17B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Sea raven.




Figure 7.17C. Length frequency distribution in NAFO units 4X and 4VW for Sea raven.






Figure 7.17D. Average fish condition in NAFO units 4X and 4VW for Sea raven.




834



Freq	Depth	Temp	Sal
F5	22	1.1	31.00
F25	32	3.1	32.43
F50	45	5.4	33.15
F75	61	8.0	34.35
F95	89	10.0	35.05

Figure 7.17E. Catch distribution by depth, temperature and salinity of Sea raven.



835

Figure 7.17F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Sea raven.

836 **7.18 Alligatorfish (Poisson-alligator atlantique) - species code 340 (category LF)**

837 Scientific name: [Aspidophoroides monopterygius](#)

838




Figure 7.18A. Inverse distance weighted distribution of catch biomass (kg/tow) for Alligatorfish.

839




Figure 7.18B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Alligatorfish.




Figure 7.18C. Length frequency distribution in NAFO units 4X and 4VW for Alligatorfish.






Figure 7.18D. Average fish condition in NAFO units 4X and 4VW for Alligatorfish.



842



Freq	Depth	Temp	Sal
F5	32	1.5	31.00
F25	44	3.7	32.53
F50	57	6.1	33.28
F75	72	8.2	34.45
F95	102	10.0	35.10

Figure 7.18E. Catch distribution by depth, temperature and salinity of Alligatorfish.

843




Figure 7.18F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Alligatorfish.

844

7.19 Monkfish (Baudroie d'Amérique) - species code 400 (category LF)

845

Scientific name: [Lophius americanus](#)

846




Figure 7.19A. Inverse distance weighted distribution of catch biomass (kg/tow) for Monkfish.

847




Figure 7.19B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Monkfish.




Figure 7.19C. Length frequency distribution in NAFO units 4X and 4VW for Monkfish.




Figure 7.19D. Average fish condition in NAFO units 4X and 4VW for Monkfish.




Figure 7.19E. Catch distribution by depth, temperature and salinity of Monkfish.




Figure 7.19F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Monkfish.

852

7.20 Ocean pout (Loquette d'Amérique) - species code 640 (category LF)

853

Scientific name: [Zoarces americanus](#)

854




Figure 7.20A. Inverse distance weighted distribution of catch biomass (kg/tow) for Ocean pout.

855




Figure 7.20B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Ocean pout.




Figure 7.20C. Length frequency distribution in NAFO units 4X and 4VW for Ocean pout.






Figure 7.20D. Average fish condition in NAFO units 4X and 4VW for Ocean pout.



858



Freq	Depth	Temp	Sal
F5	31	1.0	31.00
F25	52	2.8	32.46
F50	69	5.1	33.22
F75	85	7.7	34.34
F95	116	10.0	35.03

Figure 7.20E. Catch distribution by depth, temperature and salinity of Ocean pout.

859




Figure 7.20F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Ocean pout.

860 **7.21 Thorny skate (Raie épineuse) - species code 201 (category LF)**

861 Scientific name: [Amblyraja radiata](#)

862




Figure 7.21A. Inverse distance weighted distribution of catch biomass (kg/tow) for Thorny skate.

863




Figure 7.21B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Thorny skate.




Figure 7.21C. Length frequency distribution in NAFO units 4X and 4VW for Thorny skate.






Figure 7.21D. Average fish condition in NAFO units 4X and 4VW for Thorny skate.




866



Freq	Depth	Temp	Sal
F5	29	1.3	31.00
F25	45	3.7	32.50
F50	67	6.2	33.30
F75	91	8.1	34.40
F95	148	9.9	35.03

Figure 7.21E. Catch distribution by depth, temperature and salinity of Thorny skate.



867

Figure 7.21F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Thorny skate.

868

7.22 Smooth skate (Raie lisse) - species code 202 (category LF)

869

Scientific name: [Malacoraja senta](#)

870




Figure 7.22A. Inverse distance weighted distribution of catch biomass (kg/tow) for Smooth skate.

871




Figure 7.22B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Smooth skate.




Figure 7.22C. Length frequency distribution in NAFO units 4X and 4VW for Smooth skate.






Figure 7.22D. Average fish condition in NAFO units 4X and 4VW for Smooth skate.



874



Freq	Depth	Temp	Sal
F5	33	1.2	31.00
F25	59	3.5	32.47
F50	80	5.9	33.23
F75	110	8.1	34.45
F95	171	10.0	35.06

Figure 7.22E. Catch distribution by depth, temperature and salinity of Smooth skate.




Figure 7.22F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Smooth skate.

876

7.23 Winter skate (Raie tachetée) - species code 204 (category LF)

877

Scientific name: [Leucoraja ocellata](#)

878




Figure 7.23A. Inverse distance weighted distribution of catch biomass (kg/tow) for Winter skate.

879




Figure 7.23B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Winter skate.




Figure 7.23C. Length frequency distribution in NAFO units 4X and 4VW for Winter skate.






Figure 7.23D. Average fish condition in NAFO units 4X and 4VW for Winter skate.



882



Freq	Depth	Temp	Sal
F5	21	1.1	31.00
F25	32	3.5	32.44
F50	45	5.9	33.19
F75	71	8.3	34.42
F95	140	10.0	35.03

Figure 7.23E. Catch distribution by depth, temperature and salinity of Winter skate.




Figure 7.23F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Winter skate.

884 **7.24 Picked dogfish (Aiguillat commun) - species code 220 (category LF)**

885 Scientific name: [Squalus acanthias](#)

886




Figure 7.24A. Inverse distance weighted distribution of catch biomass (kg/tow) for Picked dogfish.

887




Figure 7.24B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Picked dogfish.




Figure 7.24C. Length frequency distribution in NAFO units 4X and 4VW for Picked dogfish.






Figure 7.24D. Average fish condition in NAFO units 4X and 4VW for Picked dogfish.



890



Freq	Depth	Temp	Sal
F5	35	1.1	31.00
F25	60	2.8	32.47
F50	83	5.2	33.28
F75	108	7.7	34.37
F95	139	10.0	35.02

Figure 7.24E. Catch distribution by depth, temperature and salinity of Picked dogfish.




Figure 7.24F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Picked dogfish.

892 **7.25 Northern shortfin squid (*Encornet rouge nordique*) - species code 4511 (category
893 LF)**

894 Scientific name: [Illex illecebrosus](#)

895




Figure 7.25A. Inverse distance weighted distribution of catch biomass (kg/tow) for Northern shortfin squid.

896




Figure 7.25B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Northern shortfin squid.




Figure 7.25C. Length frequency distribution in NAFO units 4X and 4VW for Northern shortfin squid.






Figure 7.25D. Average fish condition in NAFO units 4X and 4VW for Northern shortfin squid.



899



Freq	Depth	Temp	Sal
F5	31	1.3	31.00
F25	53	3.5	32.48
F50	71	5.9	33.28
F75	93	8.1	34.39
F95	139	10.0	35.05

Figure 7.25E. Catch distribution by depth, temperature and salinity of Northern shortfin squid.




Figure 7.25F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Northern shortfin squid.

901 **7.26 Atlantic hagfish (*Myxine du nord*) - species code 241 (category LI)**

902 Scientific name: [Myxine glutinosa](#)




Figure 7.26A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic hagfish.




Figure 7.26B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic hagfish.

905 **7.27 Cusk (Brosme) - species code 15 (category LI)**

906 Scientific name: [Brosme brosme](#)

907




Figure 7.27A. Inverse distance weighted distribution of catch biomass (kg/tow) for Cusk.

908




Figure 7.27B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Cusk.

909 **7.28 Greenland halibut (Flétan noir) - species code 31 (category LI)**

910 Scientific name: *Reinhardtius hippoglossoides*




Figure 7.28A. Inverse distance weighted distribution of catch biomass (kg/tow) for Greenland halibut.




Figure 7.28B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Greenland halibut.

913

7.29 Gulf Stream flounder (Plie du Gulf Stream) - species code 44 (category LI)

914

Scientific name: [Citharichthys arctifrons](#)

915




Figure 7.29A. Inverse distance weighted distribution of catch biomass (kg/tow) for Gulf Stream flounder.

916




Figure 7.29B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Gulf Stream flounder.

917

7.30 American shad (*Alose savoureuse*) - species code 61 (category LI)

918

Scientific name: [Alosa sapidissima](#)

919




Figure 7.30A. Inverse distance weighted distribution of catch biomass (kg/tow) for American shad.

920




Figure 7.30B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of American shad.

921

7.31 Alewife (Gaspareau) - species code 62 (category LI)

922

Scientific name: *Alosa pseudoharengus*

923




Figure 7.31A. Inverse distance weighted distribution of catch biomass (kg/tow) for Alewife.

924




Figure 7.31B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Alewife.

925

7.32 Capelin (Capelan) - species code 64 (category LI)

926

Scientific name: [Mallotus villosus](#)

927




Figure 7.32A. Inverse distance weighted distribution of catch biomass (kg/tow) for Capelin.

928




Figure 7.32B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Capelin.

929 **7.33 Atlantic mackerel (*Maquereau commun*) - species code 70 (category LI)**

930 Scientific name: *Scomber scombrus*




Figure 7.33A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic mackerel.




Figure 7.33B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic mackerel.

933

7.34 Longfin hake (Merluche à longues nageoires) - species code 112 (category LI)

934

Scientific name: [Phycis chesteri](#)

935




Figure 7.34A. Inverse distance weighted distribution of catch biomass (kg/tow) for Longfin hake.

936




Figure 7.34B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Longfin hake.

937

7.35 Fourbeard rockling (Motelle à quatre barbillons) - species code 114 (category LI)

938

Scientific name: [Enchelyopus cimbrius](#)

939




Figure 7.35A. Inverse distance weighted distribution of catch biomass (kg/tow) for Fourbeard rockling.

940




Figure 7.35B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Fourbeard rockling.

941

7.36 Blackbelly rosefish (Sébaste chèvre) - species code 123 (category LI)

942

Scientific name: [Helicolenus dactylopterus](#)

943




Figure 7.36A. Inverse distance weighted distribution of catch biomass (kg/tow) for Blackbelly rosefish.

944




Figure 7.36B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Blackbelly rosefish.

945

7.37 Greater argentine (Grande argentine) - species code 160 (category LI)

946

Scientific name: [Argentina silus](#)

947




Figure 7.37A. Inverse distance weighted distribution of catch biomass (kg/tow) for Greater argentine.

948




Figure 7.37B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Greater argentine.

949 **7.38 Arctic hookear sculpin (*Hameçon neigeux*) - species code 306 (category LI)**

950 Scientific name: [Artediellus uncinatus](#)




Figure 7.38A. Inverse distance weighted distribution of catch biomass (kg/tow) for Arctic hookear sculpin.




Figure 7.38B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Arctic hookear sculpin.

953

7.39 Atlantic poacher (*Agone atlantique*) - species code 350 (category LI)

954

Scientific name: [Leptagonus decagonus](#)

955




Figure 7.39A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic poacher.

956




Figure 7.39B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic poacher.

957 **7.40 Marlin-spike grenadier (Grenadier du Grand Banc) - species code 410 (category**
 958 **LI)**

959 Scientific name: [Nezumia bairdii](#)




Figure 7.40A. Inverse distance weighted distribution of catch biomass (kg/tow) for Marlin-spike grenadier.




Figure 7.40B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Marlin-spike grenadier.

962

7.41 Lumpfish (Lompe) - species code 501 (category LI)

963

Scientific name: [Cyclopterus lumpus](#)

964




Figure 7.41A. Inverse distance weighted distribution of catch biomass (kg/tow) for Lumpfish.

965




Figure 7.41B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Lumpfish.

966 **7.42 Atlantic spiny lumpsucker (Petite poule de mer atlantique) - species code 502**
 967 (**category LI**)

968 Scientific name: [Eumicrotremus spinosus](#)




Figure 7.42A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic spiny lumpsucker.




Figure 7.42B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic spiny lumpsucker.

971

7.43 Sand lance (Lançon) - species code 610 (category LI)

972

Scientific name: [Ammodytes dubius](#)

973




Figure 7.43A. Inverse distance weighted distribution of catch biomass (kg/tow) for Sand lance.

974




Figure 7.43B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Sand lance.

975

7.44 Snakeblenny (Lompénie-serpent) - species code 622 (category LI)

976

Scientific name: [Lumpenus lampretaeformis](#)

977




Figure 7.44A. Inverse distance weighted distribution of catch biomass (kg/tow) for Snakeblenny.

978




Figure 7.44B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Snakeblenny.

979

7.45 Daubed shanny (Lompénie tachetée) - species code 623 (category LI)

980

Scientific name: [Leptoclinus maculatus](#)

981




Figure 7.45A. Inverse distance weighted distribution of catch biomass (kg/tow) for Daubed shanny.

982




Figure 7.45B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Daubed shanny.

983

7.46 Vahl's eelpout (*Lycodes vahlii*) - species code 647 (category LI)

984

Scientific name: [Lycodes vahlii](#)

985




Figure 7.46A. Inverse distance weighted distribution of catch biomass (kg/tow) for Vahl's eelpout.

986




Figure 7.46B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Vahl's eelpout.

987

7.47 Atlantic butterfish (*Stromaté fossette*) - species code 701 (category LI)

988

Scientific name: [Peprilus triacanthus](#)

989




Figure 7.47A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic butterfish.

990




Figure 7.47B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic butterfish.

991 **7.48 Atlantic hookear sculpin (*Hameçon atlantique*) - species code 880 (category LI)**

992 Scientific name: [Artediellus atlanticus](#)




Figure 7.48A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic hookear sculpin.




Figure 7.48B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic hookear sculpin.

995 **7.49 Barndoor skate (Grande raie) - species code 200 (category LI)**

996 Scientific name: *Dipturus laevis*




Figure 7.49A. Inverse distance weighted distribution of catch biomass (kg/tow) for Barndoor skate.




Figure 7.49B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Barndoor skate.


999

7.50 Little skate (Raie hérisson) - species code 203 (category LI)

1000

Scientific name: [Leucoraja erinacea](#)

1001



1002




Figure 7.50B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Little skate.

1003

7.51 Northern prawn (Crevette nordique) - species code 2211 (category SF)

1004

Scientific name: [Pandalus borealis](#)

1005




Figure 7.51A. Inverse distance weighted distribution of catch biomass (kg/tow) for Northern prawn.

1006




Figure 7.51B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Northern prawn.

1007

7.52 Jonah crab (*Tourteau jona*) - species code 2511 (category SF)

1008

Scientific name: [Cancer borealis](#)

1009




Figure 7.52A. Inverse distance weighted distribution of catch biomass (kg/tow) for Jonah crab.

1010




Figure 7.52B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Jonah crab.

1011

7.53 Atlantic rock crab (Tourteau poïnclos) - species code 2513 (category SF)

1012

Scientific name: [Cancer irroratus](#)

1013




Figure 7.53A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic rock crab.

1014




Figure 7.53B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic rock crab.

1015 **7.54 Arctic lyre crab (*Crabe Hyas coarctatus*) - species code 2521 (category SF)**

1016 Scientific name: [Hyas coarctatus](#)

1017




Figure 7.54A. Inverse distance weighted distribution of catch biomass (kg/tow) for Arctic lyre crab.

1018




Figure 7.54B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Arctic lyre crab.

1019

7.55 Atlantic king crab (Crabe épineux du nord) - species code 2523 (category SF)

1020

Scientific name: [Lithodes maja](#)

1021




Figure 7.55A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic king crab.

1022




Figure 7.55B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic king crab.

1023

7.56 Queen crab (Crabe des neiges) - species code 2526 (category SF)

1024

Scientific name: [Chionoecetes opilio](#)

1025




Figure 7.56A. Inverse distance weighted distribution of catch biomass (kg/tow) for Queen crab.

1026




Figure 7.56B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Queen crab.

1027

7.57 Great spider crab (Crabe lyre araignée) - species code 2527 (category SF)

1028

Scientific name: [Hyas araneus](#)

1029




Figure 7.57A. Inverse distance weighted distribution of catch biomass (kg/tow) for Great spider crab.

1030




Figure 7.57B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Great spider crab.

1031

7.58 American lobster (Homard américain) - species code 2550 (category SF)

1032

Scientific name: [Homarus americanus](#)

1033




Figure 7.58A. Inverse distance weighted distribution of catch biomass (kg/tow) for American lobster.

1034




Figure 7.58B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of American lobster.

1035

7.59 Sea lamprey (*Lamproie marine*) - species code 240 (category LR)

1036

Scientific name: [Petromyzon marinus](#)

1037




Figure 7.59A. Catch distribution for Sea lamprey.

1038

7.60 Atlantic tomcod (*Poulamon atlantique*) - species code 17 (category LR)

1039

Scientific name: [Micogadus tomcod](#)

1040




Figure 7.60A. Catch distribution for Atlantic tomcod.

1041

7.61 Offshore silver hake (Merlu argenté du large) - species code 19 (category LR)

1042

Scientific name: [Merluccius albidus](#)

1043




Figure 7.61A. Catch distribution for Offshore silver hake.

1044

7.62 Spotted wolffish (*Loup tacheté*) - species code 51 (category LR)

1045

Scientific name: [Anarhichas minor](#)

1046




Figure 7.62A. Catch distribution for Spotted wolffish.

1047

7.63 Northern wolffish (Loup à tête large) - species code 52 (category LR)

1048

Scientific name: [Anarhichas denticulatus](#)

1049




Figure 7.63A. Catch distribution for Northern wolffish.

1050

7.64 Rainbow smelt (Éperlan arc-en-ciel) - species code 63 (category LR)

1051

Scientific name: [Osmerus mordax](#)

1052




Figure 7.64A. Catch distribution for Rainbow smelt.

1053

7.65 Cunner (Tanche-tautogue) - species code 122 (category LR)

1054

Scientific name: [Tautogolabrus adspersus](#)

1055




Figure 7.65A. Catch distribution for Cunner.

1056

7.66 Fourspot flounder (Cardeau à quatre ocelles) - species code 142 (category LR)

1057

Scientific name: [Hippoglossina oblonga](#)

1058




Figure 7.66A. Catch distribution for Fourspot flounder.

1059

7.67 Windowpane flounder (Turbot de sable) - species code 143 (category LR)

1060

Scientific name: [Scophthalmus aquosus](#)

1061




Figure 7.67A. Catch distribution for Windowpane flounder.

1062

7.68 Longnose greeneye (Oeil-vert à long nez) - species code 149 (category LR)

1063

Scientific name: [Parasudis triculenta](#)

1064




Figure 7.68A. Catch distribution for Longnose greeneye.

1065

7.69 Lanternfishes (Poissons-lanternes) - species code 150 (category LR)

1066

Scientific name: [Myctophidae](#)

1067




Figure 7.69A. Catch distribution for Lanternfishes.

1068

7.70 Shortnose greeneye (Éperlan du large) - species code 156 (category LR)

1069

Scientific name: [Chlorophthalmus agassizi](#)

1070




Figure 7.70A. Catch distribution for Shortnose greeneye.

1071

7.71 Silvery lightfish (Brossé améthyste) - species code 158 (category LR)

1072

Scientific name: [Maurolicus muelleri](#)

1073




Figure 7.71A. Catch distribution for Silvery lightfish.

1074

7.72 Boa dragonfish (Dragon-boa) - species code 159 (category LR)

1075

Scientific name: [Stomias boa](#)

1076




Figure 7.72A. Catch distribution for Boa dragonfish.

1077 **7.73 Shorthorn sculpin (Chabosseau à épines courtes) - species code 301 (category**
1078 **LR)**

1079 Scientific name: [Myoxocephalus scorpius](#)




Figure 7.73A. Catch distribution for Shorthorn sculpin.

1081

7.74 Grubby (Chabosseau bronzé) - species code 303 (category LR)

1082

Scientific name: [Myoxocephalus aenaeus](#)

1083




Figure 7.74A. Catch distribution for Grubby.

1084

7.75 Polar sculpin (Cotte polaire) - species code 307 (category LR)

1085

Scientific name: [Cottunculus microps](#)

1086




Figure 7.75A. Catch distribution for Polar sculpin.

1087

7.76 Spatulate sculpin (Icèle spatulée) - species code 314 (category LR)

1088

Scientific name: *Icelus spatula*

1089




Figure 7.76A. Catch distribution for Spatulate sculpin.

1090

7.77 Arctic alligatorfish (Poisson-alligator arctique) - species code 341 (category LR)

1091

Scientific name: [Ulcina olrikii](#)

1092




Figure 7.77A. Catch distribution for Arctic alligatorfish.

1093

7.78 Alligatorfishes (Poissons-alligator) - species code 351 (category LR)

1094

Scientific name: [Agonidae](#)

1095




Figure 7.78A. Catch distribution for Alligatorfishes.

1096

7.79 Roughnose grenadier (Grenadier-scie) - species code 412 (category LR)

1097

Scientific name: [Trachyrincus murrayi](#)

1098




Figure 7.79A. Catch distribution for Roughnose grenadier.

1099

7.80 Roundnose grenadier (Grenadier de roche) - species code 414 (category LR)

1100

Scientific name: [Coryphaenoides rupestris](#)

1101




Figure 7.80A. Catch distribution for Roundnose grenadier.

1102

7.81 Atlantic seasnail (*Limace atlantique*) - species code 503 (category LR)

1103

Scientific name: [Liparis atlanticus](#)

1104




Figure 7.81A. Catch distribution for Atlantic seasnail.


1105

7.82 Gelatinous snailfish (*Limace gélatineuse*) - species code 505 (category LR)

1106

Scientific name: [Liparis fabricii](#)

1107




1108

7.83 Variegated snailfish (*Limace marbée*) - species code 512 (category LR)

1109

Scientific name: [Liparis gibbus](#)

1110



1111

Figure 7.83A. Catch distribution for Variegated snailfish.

1111 7.84 Sea tadpole (Petite limace de mer) - species code 520 (category LR)

1112 Scientific name: [Careproctus reinhardtii](#)




Figure 7.84A. Catch distribution for Sea tadpole.

1114

7.85 Wolf eelpout (*Lycodes à tête longue*) - species code 603 (category LR)

1115

Scientific name: [Lycenchelys verrillii](#)

1116




Figure 7.85A. Catch distribution for Wolf eelpout.

1117 7.86 Slender snipe eel (*Avocette ruban*) - species code 604 (category LR)

1118 Scientific name: [Nemichthys scolopaceus](#)




Figure 7.86A. Catch distribution for Slender snipe eel.

1120 **7.87 Newfoundland eelpout (Lycodes du Labrador) - species code 619 (category LR)**

1121 Scientific name: [Lycodes terraenovae](#)




Figure 7.87A. Catch distribution for Newfoundland eelpout.

1123 7.88 Newfoundland eelpout (*Lycodes* du Labrador) - species code 620 (category LR)

1124 Scientific name: [Lycodes lavalaei](#)




Figure 7.88A. Catch distribution for Newfoundland eelpout.

1126

7.89 Rock gunnel (Sigouine de roche) - species code 621 (category LR)

1127

Scientific name: [Pholis gunnellus](#)

1128




Figure 7.89A. Catch distribution for Rock gunnel.

1129 **7.90 Radiated shanny (*Ulvaire deux-lignes*) - species code 625 (category LR)**

1130 Scientific name: [Ulvaria subbifurcata](#)




Figure 7.90A. Catch distribution for Radiated shanny.

1132

7.91 Fourline snakeblenny (Quatre-lignes atlantique) - species code 626 (category LR)

1133

Scientific name: [Eumesogrammus praecisus](#)

1134




Figure 7.91A. Catch distribution for Fourline snakeblenny.

1135

7.92 Wrymouth (Terrassier tacheté) - species code 630 (category LR)

1136

Scientific name: [Cryptacanthodes maculatus](#)

1137




Figure 7.92A. Catch distribution for Wrymouth.

1138

7.93 Spotfin dragonet (Dragonnet tacheté) - species code 637 (category LR)

1139

Scientific name: [Foetorepus agassizii](#)

1140




Figure 7.93A. Catch distribution for Spotfin dragonet.

1141

7.94 Arctic eelpout (*Lycodes arctique*) - species code 641 (category LR)

1142

Scientific name: [Lycodes reticulatus](#)

1143




Figure 7.94A. Catch distribution for Arctic eelpout.

1144

7.95 Atlantic soft pout (*Molasse atlantique*) - species code 646 (category LR)

1145

Scientific name: [Melanostigma atlanticum](#)

1146




Figure 7.95A. Catch distribution for Atlantic soft pout.

1147

7.96 Silvery John dory (Saint Pierre argenté) - species code 704 (category LR)

1148

Scientific name: [Zenopsis conchifer](#)

1149




Figure 7.96A. Catch distribution for Silvery John dory.

1150

7.97 White barracudina (*Lussion blanc*) - species code 712 (category LR)

1151

Scientific name: [Arctozenus risso](#)

1152




Figure 7.97A. Catch distribution for White barracudina.

1153

7.98 Atlantic saury (*Balaou atlantique*) - species code 720 (category LR)

1154

Scientific name: [Scomberesox saurus](#)

1155




Figure 7.98A. Catch distribution for Atlantic saury.

1156

7.99 Hatchetfishes (Haches d'argent) - species code 741 (category LR)

1157

Scientific name: [Sternopychidae](#)

1158




Figure 7.99A. Catch distribution for Hatchetfishes.

1159

7.100 Atlantic batfish (*Malthe atlantique*) - species code 742 (category LR)

1160

Scientific name: [Dibranchus atlanticus](#)

1161




Figure 7.100A. Catch distribution for Atlantic batfish.

1162

7.101 Spottedfin tonguefish (Langue fil noir) - species code 816 (category LR)

1163

Scientific name: [Symphurus diomedeanus](#)

1164




Figure 7.101A. Catch distribution for Spottedfin tonguefish.

1165

7.102 Black dogfish (Aiguillat noir) - species code 221 (category LR)

1166

Scientific name: [Centroscyllium fabricii](#)

1167




Figure 7.102A. Catch distribution for Black dogfish.

1168

7.103 Longfin inshore squid (*Calmar totam*) - species code 4512 (category LR)

1169

Scientific name: [Doryteuthis pealeii](#)

1170




Figure 7.103A. Catch distribution for Longfin inshore squid.

1171

7.104 Red deepsea crab (Crabe rouge) - species code 2532 (category SR)

1172

Scientific name: [Chaceon quinquedens](#)

1173




Figure 7.104A. Catch distribution for Red deepsea crab.

INDEX

1174	Agone atlantique, 117	1220	Atlantic hookear sculpin, 126
1175	Agonidae, 156	1221	Atlantic king crab, 133
1176	Agonidae, 156	1222	Atlantic mackerel, 111
1177	Aspidophoroides monopterygius, 80	1223	Atlantic poacher, 117
1178	Leptagonus decagonus, 117	1224	Atlantic redfishes, 47
1179	Ulcina olrikii, 155	1225	Atlantic rock crab, 131
1180	Aiglefin, 32	1226	Atlantic saury, 176
1181	Aiguillat commun, 98	1227	Atlantic seasnail, 159
1182	Aiguillat noir, 180	1228	Atlantic soft pout, 173
1183	Alewife, 109	1229	Atlantic spiny lumpsucker, 120
1184	Alligatorfish, 80	1230	Atlantic tomcod, 138
1185	Alligatorfishes, 156	1231	Atlantic wolffish, 65
1186	Alosa pseudoharengus, 109	1232	Avocette ruban, 164
1187	Alosa sapidissima, 108	1233	Balaou atlantique, 176
1188	Alose savoureuse, 108	1234	Barndoor skate, 127
1189	Amblyraja radiata, 89	1235	Baudroie d'Amérique, 83
1190	American lobster, 136	1236	Black dogfish, 180
1191	American plaice, 53	1237	Blackbelly rosefish, 114
1192	American shad, 108	1238	Boa dragonfish, 150
1193	Ammodytes dubius, 121	1239	Brosme, 105
1194	Ammodytidae	1240	Brosme brosme, 105
1195	Ammodytes dubius, 121	1241	Brossé améthyste, 149
1196	Anarhichadidae	1242	Callionymidae
1197	Anarhichas denticulatus, 141	1243	Foetorepus agassizii, 171
1198	Anarhichas lupus, 65	1244	Calmar totam, 181
1199	Anarhichas minor, 140	1245	Cancer borealis, 130
1200	Anarhichas denticulatus, 141	1246	Cancer irroratus, 131
1201	Anarhichas lupus, 65	1247	Cancridae
1202	Anarhichas minor, 140	1248	Cancer borealis, 130
1203	Arctic alligatorfish, 155	1249	Cancer irroratus, 131
1204	Arctic eelpout, 172	1250	Capelan, 110
1205	Arctic hookear sculpin, 116	1251	Capelin, 110
1206	Arctic lyre crab, 132	1252	Cardeau à quatre ocelles, 144
1207	Arctozenus risso, 175	1253	Careproctus reinhardti, 162
1208	Argentina silus, 115	1254	Centroscyllium fabricii, 180
1209	Argentinidae	1255	Chabosseau bronzé, 152
1210	Argentina silus, 115	1256	Chabosseau à dix-huit épines, 71
1211	Artediellus atlanticus, 126	1257	Chabosseau à épines courtes, 151
1212	Artediellus uncinatus, 116	1258	Chaceon quinquedens, 182
1213	Aspidophoroides monopterygius, 80	1259	Chionoecetes opilio, 134
1214	Atlantic batfish, 178	1260	Chlorophthalmidae
1215	Atlantic butterfish, 125	1261	Chlorophthalmus agassizi, 148
1216	Atlantic cod, 29	1262	Parasudis truculenta, 146
1217	Atlantic hagfish, 104	1263	Chlorophthalmus agassizi, 148
1218	Atlantic halibut, 50		
1219	Atlantic herring, 68		

1264	Citharichthys arctifrons, 107	1310	Faux-trigle armé, 74
1265	Clupea harengus, 68	1311	Flétan de l'Atlantique, 50
1266	Clupeidae	1312	Flétan noir, 106
1267	Alosa pseudoharengus, 109	1313	Foetorepus agassizii, 171
1268	Alosa sapidissima, 108	1314	Fourbeard rockling, 113
1269	Clupea harengus, 68	1315	Fourline snakeblenny, 169
1270	Coryphaenoides rupestris, 158	1316	Fourspot flounder, 144
1271	Cotte polaire, 153		
1272	Cottidae	1317	Gadidae
1273	Artediellus atlanticus, 126	1318	Gadus morhua, 29
1274	Artediellus uncinatus, 116	1319	Melanogrammus aeglefinus, 32
1275	Icelus spatula, 154	1320	Microgadus tomcod, 138
1276	Myoxocephalus aenaeus, 152	1321	Pollachius virens, 44
1277	Myoxocephalus octodecemspinosis, 79 ²	1322	Gadus morhua, 29
1278	Myoxocephalus scorpius, 151	1323	Gaspareau, 109
1279	Triglops murrayi, 74	1324	Gelatinous snailfish, 160
1280	Cottunculus microps, 153	1325	Geryonidae
1281	Crabe des neiges, 134	1326	Chaceon quinquedens, 182
1282	Crabe Hyas coarctatus, 132	1327	Glyptocephalus cynoglossus, 56
1283	Crabe lyre araignée, 135	1328	Goberge, 44
1284	Crabe rouge, 182	1329	Grande argentine, 115
1285	Crabe épineux du nord, 133	1330	Grande raie, 127
1286	Crevette nordique, 129	1331	Great spider crab, 135
1287	Cryptacanthodes maculatus, 170	1332	Greater argentine, 115
1288	Cryptacanthodidae	1333	Greenland halibut, 106
1289	Cryptacanthodes maculatus, 170	1334	Grenadier de roche, 158
1290	Cunner, 143	1335	Grenadier du Grand Banc, 118
1291	Cusk, 105	1336	Grenadier-scie, 157
1292	Cyclopteridae	1337	Grubby, 152
1293	Cyclopterus lumpus, 119	1338	Gulf Stream flounder, 107
1294	Eumicrotremus spinosus, 120		
1295	Cyclopterus lumpus, 119	1339	Haches d'argent, 177
1296	Cynoglossidae	1340	Haddock, 32
1297	Symphurus diomedeanus, 179	1341	Hameçon atlantique, 126
1298	Daubed shanny, 123	1342	Hameçon neigeux, 116
1299	Dibranchus atlanticus, 178	1343	Hareng de l'Atlantique, 68
1300	Dipturus laevis, 127	1344	Hatchetfishes, 177
1301	Doryteuthis pealeii, 181	1345	Helicolenus dactylopterus, 114
1302	Dragon-boa, 150	1346	Hemipteridae
1303	Dragonnet tacheté, 171	1347	Hemitripterus americanus, 77
1304	Enchelyopus cimbrius, 113	1348	Hemitripterus americanus, 77
1305	Encornet rouge nordique, 101	1349	Hippoglossina oblonga, 144
1306	Etmopteridae	1350	Hippoglossoides platessoides, 53
1307	Centroscyllium fabricii, 180	1351	Hippoglossus hippoglossus, 50
1308	Eumesogrammus praecisus, 169	1352	Homard américain, 136
1309	Eumicrotremus spinosus, 120	1353	Homarus americanus, 136
		1354	Hyas araneus, 135
		1355	Hyas coarctatus, 132

1356	Hémitriptère atlantique, 77	1401	Loquette d'Amérique, 86
1357	<i>Icelus spatula</i> , 154	1402	Lotidae
1358	lcèle spatulée, 154	1403	<i>Brosme brosme</i> , 105
1359	<i>Illex illecebrosus</i> , 101	1404	<i>Enchelyopus cimbrius</i> , 113
1360	Jonah crab, 130	1405	Loup atlantique, 65
1361	Labridae	1406	Loup tacheté, 140
1362	<i>Tautogolabrus adspersus</i> , 143	1407	Loup à tête large, 141
1363	Lamproie marine, 137	1408	<i>Lumpenus lampretaeformis</i> , 122
1364	Langue fil noir, 179	1409	Lumpfish, 119
1365	Lanternfishes, 147	1410	Lussion blanc, 175
1366	Lançon, 121	1411	<i>Lycenchelys verrillii</i> , 163
1367	<i>Leptagonus decagonus</i> , 117	1412	Lycode arctique, 172
1368	<i>Leptoclinus maculatus</i> , 123	1413	Lycode du Labrador, 165, 166
1369	<i>Leucoraja erinacea</i> , 128	1414	Lycode à carreaux, 124
1370	<i>Leucoraja ocellata</i> , 95	1415	Lycode à tête longue, 163
1371	Limace atlantique, 159	1416	<i>Lycodes lavalaei</i> , 166
1372	Limace gélatineuse, 160	1417	<i>Lycodes reticulatus</i> , 172
1373	Limace marbée, 161	1418	<i>Lycodes terraenovae</i> , 165
1374	<i>Limanda ferruginea</i> , 59	1419	<i>Lycodes vahlii</i> , 124
1375	Limande à queue jaune, 59	1420	Macrouridae
1376	Limande-plie rouge, 62	1421	<i>Coryphaenoides rupestris</i> , 158
1377	Liparidae	1422	<i>Nezumia bairdii</i> , 118
1378	<i>Careproctus reinhardtii</i> , 162	1423	<i>Trachyrincus murrayi</i> , 157
1379	<i>Liparis atlanticus</i> , 159	1424	Malacoraja senta, 92
1380	<i>Liparis fabricii</i> , 160	1425	Mallotus villosus, 110
1381	<i>Liparis gibbus</i> , 161	1426	Malthe atlantique, 178
1382	<i>Liparis atlanticus</i> , 159	1427	Maquereau commun, 111
1383	<i>Liparis fabricii</i> , 160	1428	Marlin-spike grenadier, 118
1384	<i>Liparis gibbus</i> , 161	1429	<i>Maurolicus muelleri</i> , 149
1385	Lithodes maja, 133	1430	<i>Melanogrammus aeglefinus</i> , 32
1386	Lithodidae	1431	<i>Melanostigma atlanticum</i> , 173
1387	<i>Lithodes maja</i> , 133	1432	Merlu argenté, 41
1388	Little skate, 128	1433	Merlu argenté du large, 139
1389	Loliginidae	1434	Merlucciidae
1390	<i>Doryteuthis pealeii</i> , 181	1435	<i>Merluccius albidus</i> , 139
1391	Lompe, 119	1436	<i>Merluccius bilinearis</i> , 41
1392	Lompénie tachetée, 123	1437	Merluccius albidus, 139
1393	Lompénie-serpent, 122	1438	Merluccius bilinearis, 41
1394	Longfin hake, 112	1439	Merluche blanche, 35
1395	Longfin inshore squid, 181	1440	Merluche à longues nageoires, 112
1396	Longhorn sculpin, 71	1441	Merluche écureuil, 38
1397	Longnose greeneye, 146	1442	<i>Microgadus tomcod</i> , 138
1398	Lophiidae	1443	Molasse atlantique, 173
1399	<i>Lophius americanus</i> , 83	1444	Monkfish, 83
1400	Lophius americanus, 83	1445	Morue franche, 29
		1446	Motelle à quatre barbillons, 113
		1447	Moustache sculpin, 74

1448	Myctophidae, 147	1493	Petite poule de mer atlantique, 120
1449	Myctophidae, 147	1494	Petromyzon marinus, 137
1450	Myoxocephalus aenaeus, 152	1495	Petromyzontidae
1451	Myoxocephalus octodecemspinulosus, 71	1496	Petromyzon marinus, 137
1452	Myoxocephalus scorpius, 151	1497	Pholidae
1453	Myxine du nord, 104	1498	Pholis gunnellus, 167
1454	Myxine glutinosa, 104	1499	Pholis gunnellus, 167
1455	Myxinidae	1500	Phycidae
1456	Myxine glutinosa, 104	1501	Phycis chesteri, 112
1457	Nemichthyidae	1502	Urophycis chuss, 38
1458	Nemichthys scolopaceus, 164	1503	Urophycis tenuis, 35
1459	Nemichthys scolopaceus, 164	1504	Phycis chesteri, 112
1460	Nephropidae	1505	Picked dogfish, 98
1461	Homarus americanus, 136	1506	Pleuronectidae
1462	Newfoundland eelpout, 165, 166	1507	Glyptocephalus cynoglossus, 56
1463	Nezumia bairdii, 118	1508	Hippoglossoides platessoides, 53
1464	Northern prawn, 129	1509	Hippoglossus hippoglossus, 50
1465	Northern shortfin squid, 101	1510	Limanda ferruginea, 59
1466	Northern wolffish, 141	1511	Pseudopleuronectes americanus, 62
1467	Ocean pout, 86	1512	Reinhardtius hippoglossoides, 106
1468	Oeil-vert à long nez, 146	1513	Plie canadienne, 53
1469	Offshore silver hake, 139	1514	Plie du Gulf Stream, 107
1470	Ogcocephalidae	1515	Plie grise, 56
1471	Dibranchus atlanticus, 178	1516	Poisson-alligator arctique, 155
1472	Ommastrephidae	1517	Poisson-alligator atlantique, 80
1473	Illex illecebrosus, 101	1518	Poissons-alligator, 156
1474	Oregoniidae	1519	Poissons-lanternes, 147
1475	Chionoecetes opilio, 134	1520	Polar sculpin, 153
1476	Hyas araneus, 135	1521	Pollachius virens, 44
1477	Hyas coarctatus, 132	1522	Pollock, 44
1478	Osmeridae	1523	Poulamon atlantique, 138
1479	Mallotus villosus, 110	1524	Pseudopleuronectes americanus, 62
1480	Osmerus mordax, 142	1525	Psychrolutidae
1481	Osmerus mordax, 142	1526	Cottunculus microps, 153
1482	Pandalidae	1527	Quatre-lignes atlantique, 169
1483	Pandalus borealis, 129	1528	Queen crab, 134
1484	Pandalus borealis, 129	1529	Radiated shanny, 168
1485	Paralepididae	1530	Raie hérisson, 128
1486	Arctozenus risso, 175	1531	Raie lisse, 92
1487	Paralichthyidae	1532	Raie tachetée, 95
1488	Citharichthys arctifrons, 107	1533	Raie épineuse, 89
1489	Hippoglossina oblonga, 144	1534	Rainbow smelt, 142
1490	Parasudis triculenta, 146	1535	Rajidae
1491	Peprilus triacanthus, 125	1536	Amblyraja radiata, 89
1492	Petite limace de mer, 162	1537	Dipturus laevis, 127
		1538	Leucoraja erinacea, 128

1539	Leucoraja ocellata, 95	1586	Leptoclinus maculatus, 123
1540	Malacoraja senta, 92	1587	Lumpenus lampretaeformis, 122
1541	Red deepsea crab, 182	1588	Ulvaria subbifurcata, 168
1542	Red hake, 38	1589	Stomias boa, 150
1543	Reinhardtius hippoglossoides, 106	1590	Stomiidae
1544	Rock gunnel, 167	1591	Stomias boa, 150
1545	Roughnose grenadier, 157	1592	Stromateidae
1546	Roundnose grenadier, 158	1593	Peprilus triacanthus, 125
1547	Saint Pierre argenté, 174	1594	Stromaté fossette, 125
1548	Sand lance, 121	1595	Syphurus diomedeanus, 179
1549	Scomber scombrus, 111	1596	Sébaste chèvre, 114
1550	Scomberesocidae	1597	Sébastes de l'Atlantique, 47
1551	Scomberesox saurus, 176	1598	Tanche-tautogue, 143
1552	Scomberesox saurus, 176	1599	Tautogolabrus adspersus, 143
1553	Scombridae	1600	Terrassier tacheté, 170
1554	Scomber scombrus, 111	1601	Thorny skate, 89
1555	Scophthalmidae	1602	Tourteau jona, 130
1556	Scophthalmus aquosus, 145	1603	Tourteau poïnclos, 131
1557	Scophthalmus aquosus, 145	1604	Trachyrincus murrayi, 157
1558	Sea lamprey, 137	1605	Triglops murrayi, 74
1559	Sea raven, 77	1606	Turbot de sable, 145
1560	Sea tadpole, 162		
1561	Sebastes, 47	1607	Ulcina olrikii, 155
1562	Sebastidae	1608	Ulvaire deux-lignes, 168
1563	Helicolenus dactylopterus, 114	1609	Ulvaria subbifurcata, 168
1564	Sebastes, 47	1610	Urophycis chuss, 38
1565	Shorthorn sculpin, 151	1611	Urophycis tenuis, 35
1566	Shortnose greeneye, 148	1612	Vahl's eelpout, 124
1567	Sigouine de roche, 167	1613	Variegated snailfish, 161
1568	Silver hake, 41		
1569	Silvery John dory, 174	1614	White barracudina, 175
1570	Silvery lightfish, 149	1615	White hake, 35
1571	Slender snipe eel, 164	1616	Windowpane flounder, 145
1572	Smooth skate, 92	1617	Winter flounder, 62
1573	Snakeblenny, 122	1618	Winter skate, 95
1574	Spatulate sculpin, 154	1619	Witch flounder, 56
1575	Spotfin dragonet, 171	1620	Wolf eelpout, 163
1576	Spotted wolffish, 140	1621	Wrymouth, 170
1577	Spottedfin tonguefish, 179		
1578	Squalidae	1622	Yellowtail flounder, 59
1579	Squalus acanthias, 98		
1580	Squalus acanthias, 98	1623	Zeidae
1581	Sternopychidae, 177	1624	Zenopsis conchifer, 174
1582	Maurolicus muelleri, 149	1625	Zenopsis conchifer, 174
1583	Sternopychidae, 177	1626	Zoarces americanus, 86
1584	Stichaeidae	1627	Zoarcidae
1585	Eumesogrammus praecisus, 169	1628	Lycenchelys verrillii, 163

- 1629 *Lycodes lavalaei*, 166
 - 1630 *Lycodes reticulatus*, 172
 - 1631 *Lycodes terraenovae*, 165
 - 1632 *Lycodes vahlii*, 124
 - 1633 *Melanostigma atlanticum*, 173
 - 1634 *Zoarces americanus*, 86
-
- 1635 Éperlan arc-en-ciel, 142
 - 1636 Éperlan du large, 148