

Marine Fish and Invertebrate Atlas: Summarizing Geographic Distribution and Population Indices in the Scotian Shelf and Bay of Fundy (1970-2020)

Daniel Ricard, Jamie Emberley, Catalina Gomez and Catriona Regnier-McKellar

Science Branch
Gulf Region
Fisheries and Oceans Canada
Moncton, New Brunswick, E1C 5K4, Canada

Science Branch
Maritimes Region
Fisheries and Oceans Canada
Dartmouth, Nova Scotia, B2Y 4A2, Canada

2021

Canadian Technical Report of
Fisheries and Aquatic Sciences ####



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Canada

Canadian Technical Report of Fisheries and Aquatic Sciences

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Rapport technique canadien des sciences halieutiques et aquatiques

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Canadian Technical Report of
Fisheries and Aquatic Sciences nnn

2021

6 MARINE FISH AND INVERTEBRATE ATLAS: SUMMARIZING GEOGRAPHIC DISTRIBUTION,
7 POPULATION INDICES AND ENVIRONMENTAL PREFERENCES IN THE SCOTIAN SHELF
8 AND BAY OF FUNDY (1970-2020)

by

10 Daniel Ricard¹ Jamie Emberley² Catalina Gomez² Catriona Regnier-McKellar²

¹Science Branch
Gulf Region
Fisheries and Oceans Canada
Moncton, New Brunswick, E1C 5K4, Canada
²Science Branch
Maritimes Region
Fisheries and Oceans Canada
Dartmouth, Nova Scotia, B2Y 4A2, Canada

¹⁹ © Her Majesty the Queen in Right of Canada, 2021
²⁰ Cat. No. Fs97-6/nnnE-PDF ISBN ISSN 1488-5379

²¹ Correct citation for this publication:

²² Ricard, D., Emberley, J., Gomez, C. and Regnier-McKellar, C. 2021. Marine Fish and Invertebrate Atlas: Summarizing Geographic Distribution, Population Indices and Environmental Preferences in the Scotian Shelf and Bay of Fundy (1970-2020). Can. Tech. Rep. Fish. Aquat. Sci. nnn: viii + 192 p.

CONTENTS

26	ABSTRACT	vii
28	RÉSUMÉ	viii
29	1 Introduction	1
30	2 Methods	2
31	2.1 Survey Description	2
32	2.2 Sampling Design	2
33	2.3 Taxonomic Levels	5
34	2.4 Analyses	16
35	2.4.1 Geographic distribution of catches	16
36	2.4.2 Abundance and biomass indices	16
37	2.4.3 Distribution indices	16
38	2.4.4 Length frequencies	16
39	2.4.5 Length-weight relationship and condition factor	17
40	2.4.6 Depth, temperature and salinity distribution of catches	17
41	2.4.7 Density-dependent habitat selection	17
42	3 Results	18
43	3.1 Description of Figures	18
44	3.1.1 Type A	18
45	3.1.2 Type B	18
46	3.1.3 Type C	18
47	3.1.4 Type D	19
48	3.1.5 Type E	19
49	3.1.6 Type F	19
50	3.2 Summary of successful tows by year and stratum	19
51	4 Discussion	24
52	4.1 Diversity of approaches used for mapping fish and invertebrates in the Scotian	
53	Shelf bioregion	24
54	4.2 Interpreting spatial results for marine spatial planning purposes	26
55	5 Acknowledgements	27
56	6 References	28
57	7 Appendix	32
58	7.1 Atlantic cod (<i>Morue franche</i>) - species code 10 (category LF)	33
59	7.2 Haddock (<i>Aiglefin</i>) - species code 11 (category LF)	36
60	7.3 White hake (<i>Merluche blanche</i>) - species code 12 (category LF)	39
61	7.4 Red hake (<i>Merluche écureuil</i>) - species code 13 (category LF)	42
62	7.5 Silver hake (<i>Merlu argenté</i>) - species code 14 (category LF)	45
63	7.6 Pollock (<i>Goberge</i>) - species code 16 (category LF)	48
64	7.7 Atlantic redfishes (<i>Sébastes de l'Atlantique</i>) - species code 23 (category LF)	51
65	7.8 Atlantic halibut (<i>Flétan de l'Atlantique</i>) - species code 30 (category LF)	54

66	7.9 American plaice (Plie canadienne) - species code 40 (category LF)	57
67	7.10 Witch flounder (Plie grise) - species code 41 (category LF)	60
68	7.11 Yellowtail flounder (Limande à queue jaune) - species code 42 (category LF)	63
69	7.12 Winter flounder (Limande-plie rouge) - species code 43 (category LF)	66
70	7.13 Atlantic Wolffish (Loup atlantique) - species code 50 (category LF)	69
71	7.14 Atlantic herring (Hareng de l'Atlantique) - species code 60 (category LF)	72
72	7.15 Longhorn sculpin (Chabosseau à dix-huit épines) - species code 300 (category LF)	75
73	7.16 Moustache sculpin (Faux-trigle armé) - species code 304 (category LF)	78
74	7.17 Sea raven (Hémithriptère atlantique) - species code 320 (category LF)	81
75	7.18 Alligatorfish (Poisson-alligator atlantique) - species code 340 (category LF)	84
76	7.19 Monkfish (Baudroie d'Amérique) - species code 400 (category LF)	87
77	7.20 Ocean pout (Loquette d'Amérique) - species code 640 (category LF)	90
78	7.21 Thorny skate (Raie épineuse) - species code 201 (category LF)	93
79	7.22 Smooth skate (Raie lisse) - species code 202 (category LF)	96
80	7.23 Winter skate (Raie tachetée) - species code 204 (category LF)	99
81	7.24 Picked dogfish (Aiguillat commun) - species code 220 (category LF)	102
82	7.25 Northern shortfin squid (Encornet rouge nordique) - species code 4511 (category LF)	105
83	7.26 Atlantic hagfish (Myxine du nord) - species code 241 (category LI)	108
84	7.27 Cusk (Brosme) - species code 15 (category LI)	109
85	7.28 Greenland halibut (Flétan noir) - species code 31 (category LI)	110
86	7.29 Gulf Stream flounder (Plie du Gulf Stream) - species code 44 (category LI)	111
87	7.30 American shad (Alose savoureuse) - species code 61 (category LI)	112
88	7.31 Alewife (Gaspareau) - species code 62 (category LI)	113
89	7.32 Capelin (Capelan) - species code 64 (category LI)	114
90	7.33 Atlantic mackerel (Maquereau commun) - species code 70 (category LI)	115
91	7.34 Longfin hake (Merluche à longues nageoires) - species code 112 (category LI)	116
92	7.35 Fourbeard rockling (Motelle à quatre barbillons) - species code 114 (category LI)	117
93	7.36 Blackbelly rosefish (Sébaste chèvre) - species code 123 (category LI)	118
94	7.37 Greater argentine (Grande argentine) - species code 160 (category LI)	119
95	7.38 Arctic hookear sculpin (Hameçon neigeux) - species code 306 (category LI)	120
96	7.39 Atlantic poacher (Agone atlantique) - species code 350 (category LI)	121
97	7.40 Marlin-spike grenadier (Grenadier du Grand Banc) - species code 410 (category LI)	122
98	7.41 Lumpfish (Lompe) - species code 501 (category LI)	123
99	7.42 Atlantic spiny lumpsucker (Petite poule de mer atlantique) - species code 502 (category LI)	124
100	7.43 Sand lance (Lançon) - species code 610 (category LI)	125
101	7.44 Snakeblenny (Lompénie-serpent) - species code 622 (category LI)	126
102	7.45 Daubed shanny (Lompénie tachetée) - species code 623 (category LI)	127
103	7.46 Vahl's eelpout (Lycoide à carreaux) - species code 647 (category LI)	128
104	7.47 Atlantic butterfish (Stromaté fossette) - species code 701 (category LI)	129
105	7.48 Atlantic hookear sculpin (Hameçon atlantique) - species code 880 (category LI)	130
106	7.49 Barndoor skate (Grande raie) - species code 200 (category LI)	131
107	7.50 Little skate (Raie hérisson) - species code 203 (category LI)	132
108	7.51 Northern prawn (Crevette nordique) - species code 2211 (category SF)	133
109	7.52 Jonah crab (Tourteau jona) - species code 2511 (category SF)	134
110	7.53 Atlantic rock crab (Tourteau poïnclos) - species code 2513 (category SF)	135

113	7.54 Arctic lyre crab (Crabe Hyas coarctatus) - species code 2521 (category SF)	136
114	7.55 Atlantic king crab (Crabe épineux du nord) - species code 2523 (category SF)	137
115	7.56 Queen crab (Crabe des neiges) - species code 2526 (category SF)	138
116	7.57 Great spider crab (Crabe lyre araignée) - species code 2527 (category SF)	139
117	7.58 American lobster (Homard américain) - species code 2550 (category SF)	140
118	7.59 Sea lamprey (Lamproie marine) - species code 240 (category LR)	141
119	7.60 Atlantic tomcod (Poulamon atlantique) - species code 17 (category LR)	142
120	7.61 Offshore silver hake (Merlu argenté du large) - species code 19 (category LR)	143
121	7.62 Spotted wolffish (Loup tacheté) - species code 51 (category LR)	144
122	7.63 Northern wolffish (Loup à tête large) - species code 52 (category LR)	145
123	7.64 Rainbow smelt (Éperlan arc-en-ciel) - species code 63 (category LR)	146
124	7.65 Cunner (Tanche-tautogue) - species code 122 (category LR)	147
125	7.66 Fourspot flounder (Cardeau à quatre ocelles) - species code 142 (category LR)	148
126	7.67 Windowpane flounder (Turbot de sable) - species code 143 (category LR)	149
127	7.68 Longnose greeneye (Oeil-vert à long nez) - species code 149 (category LR)	150
128	7.69 Lanternfishes (Poissons-lanternes) - species code 150 (category LR)	151
129	7.70 Shortnose greeneye (Éperlan du large) - species code 156 (category LR)	152
130	7.71 Silvery lightfish (Brossé améthyste) - species code 158 (category LR)	153
131	7.72 Boa dragonfish (Dragon-boa) - species code 159 (category LR)	154
132	7.73 Shorthorn sculpin (Chabosseau à épines courtes) - species code 301 (category LR)	155
133	7.74 Grubby (Chabosseau bronzé) - species code 303 (category LR)	156
134	7.75 Polar sculpin (Cotte polaire) - species code 307 (category LR)	157
135	7.76 Spatulate sculpin (Icèle spatulée) - species code 314 (category LR)	158
136	7.77 Arctic alligatorfish (Poisson-alligator arctique) - species code 341 (category LR) . .	159
137	7.78 Alligatorfishes (Poissons-alligator) - species code 351 (category LR)	160
138	7.79 Roughnose grenadier (Grenadier-scie) - species code 412 (category LR)	161
139	7.80 Roundnose grenadier (Grenadier de roche) - species code 414 (category LR)	162
140	7.81 Atlantic seasnail (Limace atlantique) - species code 503 (category LR)	163
141	7.82 Gelatinous snailfish (Limace gélantineuse) - species code 505 (category LR)	164
142	7.83 Variegated snailfish (Limace marbée) - species code 512 (category LR)	165
143	7.84 Sea tadpole (Petite limace de mer) - species code 520 (category LR)	166
144	7.85 Wolf eelpout (Lycode à tête longue) - species code 603 (category LR)	167
145	7.86 Slender snipe eel (Avocette ruban) - species code 604 (category LR)	168
146	7.87 Newfoundland eelpout (Lycode du Labrador) - species code 619 (category LR) . .	169
147	7.88 Newfoundland eelpout (Lycode du Labrador) - species code 620 (category LR) . .	170
148	7.89 Rock gunnel (Sigouine de roche) - species code 621 (category LR)	171
149	7.90 Radiated shanny (Ulvaire deux-lignes) - species code 625 (category LR)	172
150	7.91 Fourline snakeblenny (Quatre-lignes atlantique) - species code 626 (category LR) .	173
151	7.92 Wrymouth (Terrassier tacheté) - species code 630 (category LR)	174
152	7.93 Spotfin dragonet (Dragonnet tacheté) - species code 637 (category LR)	175
153	7.94 Arctic eelpout (Lycode arctique) - species code 641 (category LR)	176
154	7.95 Atlantic soft pout (Molasse atlantique) - species code 646 (category LR)	177
155	7.96 Silvery John dory (Saint Pierre argenté) - species code 704 (category LR)	178
156	7.97 White barracudina (Lussion blanc) - species code 712 (category LR)	179
157	7.98 Atlantic saury (Balaou atlantique) - species code 720 (category LR)	180
158	7.99 Hatchetfishes (Haches d'argent) - species code 741 (category LR)	181
159	7.100 Atlantic batfish (Malthe atlantique) - species code 742 (category LR)	182

160	7.101 Spottedfin tonguefish (Langue fil noir) - species code 816 (category LR)	183
161	7.102 Black dogfish (Aiguillat noir) - species code 221 (category LR)	184
162	7.103 Longfin inshore squid (Calmar totam) - species code 4512 (category LR)	185
163	7.104 Red deepsea crab (Crabe rouge) - species code 2532 (category SR)	186

164	INDEX	187
-----	--------------	------------

ABSTRACT

166 Ricard, D., Emberley, J., Gomez, C. and Regnier-McKellar, C. 2021. Marine Fish and Invertebrate
167 Atlas: Summarizing Geographic Distribution, Population Indices and Environmental
168 Preferences in the Scotian Shelf and Bay of Fundy (1970-2020). Can. Tech. Rep. Fish. Aquat.
169 Sci. nnn: viii + 192 p.

170 The summer groundfish research vessel survey on the Scotian Shelf and in the Bay of
171 Fundy started in 1970 and was designed to measure the distribution and abundance of
172 major commercial fish species. Over time, additional information on non-commercial species
173 was collected, and allowed considerable insight into ecosystem function and structure, as
174 documented in many primary publications whose analyses used the survey data. The same
175 groundfish survey database has also been used to produce species status reports, atlases of
176 species distribution and remains an essential source of information for stock assessments in the
177 Maritimes Region of Fisheries and Oceans Canada. This report builds on previous work and
178 former atlases by updating a comprehensive suite of indices to assess population status and
179 environmental preferences of 104 species. For each species, trends in geographic distribution
180 and biomass or abundance were plotted. The spatial extent of distribution was plotted over
181 time to gauge how the area occupied has changed. The relationship between abundance or
182 biomass and spatial extent reflected whether the species distribution expands when abundance
183 or biomass increases. Length frequencies over time depicted any changes in mean size. The
184 plots of condition over time revealed whether individual fish are fatter or thinner than their long
185 term mean. Depth, temperature and salinity preferences were estimated to gauge the range
186 of suitable environmental parameters for each species. Finally, for each stratum, the slope
187 describing how local density varies with regional abundance was estimated.

RÉSUMÉ

189 Ricard, D., Emberley, J., Gomez, C. and Regnier-McKellar, C. 2021. Marine Fish and Invertebrate
190 Atlas: Summarizing Geographic Distribution, Population Indices and Environmental
191 Preferences in the Scotian Shelf and Bay of Fundy (1970-2020). Can. Tech. Rep. Fish. Aquat.
192 Sci. nnn: viii + 192 p.

193 Le relevé estival par navires de recherche sur le poisson de fond sur le plateau néo-écossais
194 et dans la baie de Fundy a débuté en 1970 et visait à mesurer la répartition et l'abondance
195 des principales espèces de poissons commerciales. Au fil du temps, des informations
196 supplémentaires sur les espèces non commerciales ont été recueillies et ont permis de mieux
197 comprendre la fonction et la structure de l'écosystème, comme le montrent de nombreuses
198 publications primaires dont les analyses ont utilisé les données des relevés. La même base
199 de données sur les relevés du poisson de fond a également été utilisée pour produire des
200 rapports sur la situation des espèces, des atlas de la répartition des espèces et demeure une
201 source essentielle d'information pour les évaluations des stocks dans la région des Maritimes
202 de Pêches et Océans Canada. Ce rapport s'appuie sur des travaux antérieurs et d'anciens
203 atlas en mettant à jour une série complète d'indices pour évaluer l'état de la population et les
204 préférences environnementales de 104 espèces. Pour chaque espèce, les tendances de la
205 répartition géographique et de la biomasse ou de l'abondance ont été tracées. L'étendue spatiale
206 de la distribution a été tracée au fil du temps pour évaluer comment la zone occupée a changé.
207 La relation entre l'abondance ou la biomasse et l'étendue spatiale indique si la répartition des
208 espèces augmente lorsque l'abondance ou la biomasse augmente. Les fréquences de longueur
209 au fil du temps représentaient tout changement dans la taille moyenne. Les graphiques de l'état
210 au fil du temps ont révélé si les poissons individuels sont plus gros ou plus minces que leur
211 moyenne à long terme. Les préférences en matière de profondeur, de température et de salinité
212 ont été estimées pour évaluer la gamme de paramètres environnementaux appropriés pour
213 chaque espèce. Enfin, pour chaque strate, la pente décrivant comment la densité locale varie
214 avec l'abondance régionale a été estimée.

1 Introduction

216 The summer (July-August) groundfish research vessel survey on the Scotian Shelf and in the
217 Bay of Fundy was started in 1970 by Fisheries and Oceans Canada Maritimes Region. The
218 survey was originally designed to measure the distribution and abundance of major commercial
219 fish species. Over time, information on non-commercial species was also collected. The
220 groundfish survey database storing the information collected during the annual survey provides
221 the main source of fisheries-independent information for marine species in the region. This
222 information is routinely used to support stock assessments, to produce species status reports
223 and has been previously used to publish atlases of species distribution.

224 The current document is an update of an earlier report (Ricard and Shackell 2013) that built
225 on former atlases by updating a comprehensive suite of derived indices for 104 species to
226 assess population status and environmental preferences. The information collected during the
227 survey is stored in a relational database management system archived at Fisheries and Oceans
228 Canada Maritimes Region which contains detailed information about the sampling locations
229 and the associated catch. Tow-level survey data is also publicly available from the Ocean
230 Biogeographic Information System (DFO 2016) and from the Open data portal supported by
231 the federal government (DFO 2021). The present atlas builds upon the work done by Fisheries
232 and Oceans colleagues from the northern Gulf of St. Lawrence (Bourdages and Ouellet 2012),
233 southern Gulf of St. Lawrence (Benoît et al. 2003) and on earlier work in the Scotian Shelf
234 (Simon and Comeau 1994; Horsman and Shackell 2009).

235 To facilitate updates and foster collaboration on the analyses of the survey data, the computer
236 code necessary to extract the data, to perform the analyses presented herein, and to reproduce
237 and update the current document is made available in a git repository (Ricard and Gomez 2021).

238 The survey area covers three major Northwest Atlantic Fisheries Organization (NAFO) zones
239 that divide the Scotian Shelf into the colder east 4V and 4W (strata 440-466) and warmer
240 west 4X (strata 470-495). For each species, trends in geographic distribution and biomass
241 or abundance are plotted. Some caution is required in interpreting the results obtained
242 for several taxa due to low sample size as explained later in the text. The spatial extent of
243 distribution is plotted over time to gauge how the area occupied has changed. The relationship
244 between biomass and spatial extent reflects whether the species distribution expands when
245 biomass increases. For each strata, the slope describing how local density varies with regional
246 abundance was estimated (Myers and Stokes 1989). These slopes were then plotted against
247 a habitat suitability index to identify important strata for each species. Then, length frequencies
248 over time depicted any changes in mean size. The plots of condition over time revealed whether
249 individual fish are fatter or thinner than their long term mean. Finally, depth, temperature and
250 salinity preferences were estimated to gauge the range of environmental parameters (Perry
251 and Smith 1994). A full ecological interpretation of trends is beyond the scope of this report.
252 Other documents stemming from peer-reviewed scientific processes under the auspices of the
253 [Canadian Science Advisory Secretariat](#) (CSAS) provide further descriptions of spatio-temporal
254 trends in different indicators and put the information collected during the summer groundfish
255 research vessel survey in a more focused context (see for example Clark and Emberley (2011)).

2 Methods

257 2.1 Survey Description

258 The survey is conducted annually in July-August and covers the Scotian Shelf and the Bay of
 259 Fundy (Figure 1). It normally involves two separate two-week trips on board an offshore fisheries
 260 vessel from the Canadian Coast Guard.

261 A number of changes in fishing gear type and vessels used occurred since the onset of sampling
 262 activities (Clark and Emberley 2011). Comparative fishing experiments were conducted when
 263 those changes in survey platforms took place.




Figure 1. Map of the Scotian Shelf and Bay of Fundy where the DFO Maritimes Summer survey takes place. The bathymetry presented here is the 15 arc-second gridded data set from the General Bathymetric Chart of the Oceans ([GEBCO](#)). Geographical locations of interest and the boundaries of relevant NAFO Divisions are also shown on the map.

264 2.2 Sampling Design

265 The summer survey covers divisions 4V, 4W and 4X of the Northwest Atlantic Fisheries
 266 Organization (NAFO) which includes the Scotian Shelf and the Bay of Fundy. The eastern limit of

267 the survey is the Laurentian Channel and the western limit is the Fundian Channel (Figure 1).
 268 The survey follows a stratified random design (Doubleday and Rivard 1981; Lohr 1999)
 269 (Figure 2). The number of tows conducted in each stratum is approximately proportional to the
 270 surface area of the stratum. The area covered during the survey has remained constant since its
 271 onset, with the exception of additional deeper strata that were only sampled a few times since
 272 2000. Because the sampling of the deeper strata is opportunistic and irregular, the analyses
 273 presented herein only include strata 440 to 495 which cover NAFO Divisions 4V, 4W and 4X
 274 (Figure 2 and Table 1).
 275 The basic sampling unit of the survey is a 30-minute fishing tow conducted at a speed of 3.5
 276 knots. This yields a distance towed of 1.75 nautical miles.

Table 1. Summer survey strata details. The strata used in the analyses are presented separately for NAFO Divisions 4Vn, 4VsW and 4X. For each stratum, the depth range in fathoms and the surface area in square kilometers are reported.

Stratum	Depth range (fathom)	Area (km ²)
4Vn		
440	101-200	924
441	51-100	1000
442	0-49	1437
4VsW		
443	0-49	1318
444	51-100	3925
445	101-200	1023
446	101-200	491
447	0-49	1616
448	0-49	1449
449	51-100	144
450	51-100	383
451	101-200	147
452	101-200	345
453	101-200	259
454	51-100	499
455	0-49	2122
456	0-49	955
457	51-100	811
458	0-49	658

NAFO Div.	Stratum	Depth range (fathom)	Area (km²)
	459	0-200	3148
	460	51-100	1344
	461	101-200	1154
	462	51-100	2116
	463	0-49	302
	464	0-50	1297
	465	51-100	2383
	466	101-200	226
4X			
	470	51-100	920
	471	101-200	1004
	472	51-100	1249
	473	0-49	265
	474	0-49	161
	475	0-49	156
	476	51-100	1478
	477	51-100	1232
	478	101-200	233
	480	0-49	655
	481	51-100	1875
	482	101-200	1042
	483	101-200	532
	484	101-200	2264
	485	51-100	1582
	490	0-49	601
	491	51-100	687
	492	51-100	1086
	493	0-49	533
	494	0-49	417
	495	0-49	584

277 After each tow the catch is sorted by species and weighed. Each fish caught is then measured,
278 and further sampling of individual fish weight, maturity status and age are performed for different
279 length classes. When catches exceed 300 individuals, a random sub-sample is used to obtain
280 the length and weight measurements.

281 The location of representative tows appears in Figure 3.

282 **2.3 Taxonomic Levels**

283 Fish species caught during the surveys are identified by trained scientific personnel and their
284 scientific name is determined. An internal species code used in the relational database is
285 reported for each species (Losier and Waite 1989).


286 By its nature as a bottom trawl, the fishing gear used in the survey catches certain species
287 better than others. To ensure that meaningful ecological information can be extracted from
288 catch samples, we report the catch records for the subset of species that are caught reliably
289 by the gear. To appear in this atlas, a species must have had a minimum of 10 observations over
290 the duration of the survey activities. While both catch abundance and weight are recorded, the
291 weight of species that appear at low abundances is often recorded as zero in the earlier parts of
292 the survey when scales of appropriate precision were not available.

293 We divided the species caught into five categories based on 1) their taxonomic classification,
294 2) the number of recorded observations, and 3) their period of valid identification (Table 2).
295 Category "LF", for "long frequent", was assigned to species that have more than 1000 records
296 since 1970 and have been consistently identified since the onset of the survey. Category
297 "LI", for "long intermediate", was assigned to species that had between 1000 and 200 catch
298 records. Rare and elusive species (those with less than 200 catch records over the duration
299 of the survey) are also reported but to a lower level of analytical details (Category "LR", for
300 "long rare"). Category "SF", for "short frequent", was assigned to invertebrate species that were
301 consistently sampled only since 1999 (Tremblay M. J. 2007). And category "SR", for "short rare"
302 for invertebrate species consistently sampled only since 1999 and with less than 200 catch
303 records. The list of taxa covered in this document is presented in phylogenetic order (Nelson J.

Table 2. Taxonomic levels used to determine the analytical treatment for each species.

Category	Name	Description
L	long - consistently identified since the onset of the survey in 1970	
LF	long frequent	species that have more than 1000 catch records
LI	long intermediate	species that had between 1000 and 200 catch records
LR	long rare	species with less than 200 catch records
S	short - invertebrate species that were consistently sampled only since 1999	
SF	short frequent	species with more than 200 catch records
SR	short rare	species with less than 200 catch records

³⁰⁴ S. et al. 2004) in Table 3. To ensure concordance with authoritative taxonomic information, the
³⁰⁵ AphiaID from the World Register of Marine Species is also provided in Table 3 (Appeltans et al.
³⁰⁶ 2012).






Figure 3. Map of the Summer survey tows.

Table 3. List of species included in the Atlas. The species reported here are listed in phylogenetic order as per Page L. M. et al. (2013). For each taxonomic order and class, each species is listed in the table, its taxonomic family and scientific name is provided, along with its French and English common names, the species code used in the survey database, its AphiaID and a link to the World Registry of Marine Species, its number of catch records in the survey database and its classification category as defined in section 2.3.

	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Myxini								
<i>Myxiniformes</i>								
	Myxinidae	<i>Myxine glutinosa</i>	Atlantic hagfish	Myxine du nord	241	101170	804	LI
Petromyzonti								
<i>Petromyzontiformes</i>								
	Petromyzontidae	<i>Petromyzon marinus</i>	Sea lamprey	Lamproie marine	240	101174	16	LR
Actinopterygii								
<i>Gadiformes</i>								
	Gadidae	<i>Gadus morhua</i>	Atlantic cod	Morue franche	10	126436	5451	LF
		<i>Melanogrammus aeglefinus</i>	Haddock	Aiglefin	11	126437	5827	LF
	Phycidae	<i>Urophycis tenuis</i>	White hake	Merluche blanche	12	126504	3524	LF
		<i>Urophycis chuss</i>	Red hake	Merluche écureuil	13	126503	2195	LF
	Merlucciidae	<i>Merluccius bilinearis</i>	Silver hake	Merlu argenté	14	158962	4936	LF
	Lotidae	<i>Brosme brosme</i>	Cusk	Brosme	15	126447	688	LI
	Gadidae	<i>Pollachius virens</i>	Pollock	Goberge	16	126441	2787	LF
		<i>Microgadus tomcod</i>	Atlantic tomcod	Poulamon atlantique	17	158928	44	LR
	Merlucciidae	<i>Merluccius albibus</i>	Offshore silver hake	Merlu argenté du large	19	158748	161	LR
<i>Scorpaeniformes</i>								
	Sebastidae	<i>Sebastes</i>	Atlantic redfishes	Sébastes de l'Atlantique	23	126175	4152	LF
<i>Pleuronectiformes</i>								
	Pleuronectidae	<i>Hippoglossus hippoglossus</i>	Atlantic halibut	Flétan de l'Atlantique	30	127138	1634	LF

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Perciformes	Pleuronectidae	<i>Reinhardtius hippoglossoides</i>	<i>Reinhardtius hippoglossoides</i>	Greenland halibut	Flétan noir	31	127144	736	LI
			<i>Hippoglossoides platessoides</i>	American plaice	Plie canadienne	40	127137	6023	LF
			<i>Glyptocephalus cynoglossus</i>	Witch flounder	Plie grise	41	127136	4301	LF
			<i>Limanda ferruginea</i>	Yellowtail flounder	Limande à queue jaune	42	158879	3233	LF
		<i>Pseudopleuronectes americanus</i>	Winter flounder	Limande-plie rouge	43	158885	1632	LF	
	Paralichthyidae	<i>Citharichthys arctifrons</i>	Gulf Stream flounder	Plie du Gulf Stream	44	158791	382	LI	
Clupeiformes	Anarhichadidae	<i>Anarhichas lupus</i>	Atlantic wolffish	Loup atlantique	50	126758	1572	LF	
		<i>Anarhichas minor</i>	Spotted wolffish	Loup tacheté	51	126759	20	LR	
		<i>Anarhichas denticulatus</i>	Northern wolffish	Loup à tête large	52	126757	17	LR	
	Clupeidae	<i>Clupea harengus</i>	Atlantic herring	Hareng de l'Atlantique	60	126417	3487	LF	
Osmeriformes	Osmeridae	<i>Alosa sapidissima</i>	American shad	Alose savoureuse	61	158670	468	LI	
		<i>Alosa pseudoharengus</i>	Alewife	Gaspareau	62	158669	977	LI	
		<i>Osmerus mordax</i>	Rainbow smelt	Éperlan arc-en-ciel	63	126737	59	LR	
Perciformes	Scombridae	<i>Mallotus villosus</i>	Capelin	Capelan	64	126735	540	LI	
	Scombridae	<i>Scomber scombrus</i>	Atlantic mackerel	Maquereau commun	70	127023	696	LI	
Gadiformes									

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
<i>Perciformes</i>	Phycidae	<i>Phycis chesteri</i>	Longfin hake	Merluche à longues nageoires	112	158988	784	LI	
<i>Scorpaeniformes</i>	Lotidae	<i>Enchelyopus cimbrius</i>	Fourbeard rockling	Motelle à quatre barbillons	114	126450	693	LI	
<i>Pleuronectiformes</i>	Labridae	<i>Tautogolabrus adspersus</i>	Cunner	Tanche-tautogue	122	159785	82	LR	
<i>Aulopiformes</i>	Sebastidae	<i>Helicolenus dactylopterus</i>	Blackbelly rosefish	Sébaste chèvre	123	127251	610	LI	
<i>Myctophiformes</i>	Paralichthyidae	<i>Hippoglossina oblonga</i>	Fourspot flounder	Cardeau à quatre ocelles	142	158833	76	LR	
<i>Stomiiformes</i>	Scophthalmidae	<i>Scophthalmus aquosus</i>	Windowpane flounder	Turbot de sable	143	158907	115	LR	
<i>Argentiniiformes</i>	Chlorophthalmidae	<i>Parasudis tricrenata</i>	Longnose greeneye	Oeil-vert à long nez	149	158868	45	LR	
<i>Cottidae</i>	Myctophidae	<i>Myctophidae</i>	Lanternfishes	Poissons-lanternes	150	125498	160	LR	
<i>Aulopiformes</i>	Chlorophthalmidae	<i>Chlorophthalmus agassizi</i>	Shortnose greeneye	Éperlan du large	156	126336	78	LR	
<i>Scorpaeniformes</i>	Sternopychidae	<i>Maurolicus muelleri</i>	Silvery lightfish	Brossé améthyste	158	127312	52	LR	
<i>Argentiniiformes</i>	Stomiidae	<i>Stomias boa</i>	Boa dragonfish	Dragon-boa	159	127374	20	LR	
<i>Scorpaeniformes</i>	Argentinidae	<i>Argentina silus</i>	Greater argentine	Grande argentine	160	126715	963	LI	
<i>Cottidae</i>	Cottidae	<i>Myoxocephalus octodecemspinosis</i>	Longhorn sculpin	Chabosseau à dix-huit épines	300	159520	3292	LF	

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Perciformes	Triglidae	<i>Myoxocephalus scorpius</i>	Shorthorn sculpin	Chabot à épines courtes	301	127203	131	LR	
		<i>Myoxocephalus aenaeus</i>	Grubby	Chabot à épines courtes	303	159519	40	LR	
	Psychrolutidae	<i>Triglops murrayi</i>	Moustache sculpin	Faux-trigle armé	304	127205	1182	LF	
	Cottidae	<i>Artediellus uncinatus</i>	Arctic hookear sculpin	Hameçon neigeux	306	127195	306	LI	
	Hemitripteridae	<i>Cottunculus microps</i>	Polar sculpin	Cotte polaire	307	127235	29	LR	
Acanthomorpha	Gobiidae	<i>Icelus spatula</i>	Spatulate sculpin	Icèle spatulée	314	127200	40	LR	
	Gobiidae	<i>Hemitripterus americanus</i>	Sea raven	Hémithriptère atlantique	320	159518	2126	LF	
	Gobioidae	<i>Aspidophoroides monopterygius</i>	Alligatorfish	Poisson-alligator atlantique	340	159459	1029	LF	
		<i>Ulcina olrikii</i>	Arctic alligatorfish	Poisson-alligator arctique	341	274356	13	LR	
	Gobioidae	<i>Leptagonus decagonus</i>	Atlantic poacher	Agone atlantique	350	127191	266	LI	
	Gobioidae	<i>Agonidae</i>	Alligatorfishes	Poissons-alligator	351	125588	43	LR	
Lophiiformes	Lophiidae	<i>Lophius americanus</i>	Monkfish	Baudroie d'Amérique	400	159184	1970	LF	
Gadiformes	Macrouridae	<i>Nezumia bairdii</i>	Marlin-spike grenadier	Grenadier du Grand Banc	410	183289	529	LI	
		<i>Trachyrincus murrayi</i>	Roughnose grenadier	Grenadier-scie	412	126481	18	LR	
Scorpaeniformes	Cyclopteridae	<i>Cyclopterus lumpus</i>	Lumpfish	Lompe	501	127214	216	LI	

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
			<i>Eumicrotremus spinosus</i>	Atlantic spiny lump sucker	Petite poule de mer atlantique	502	127217	226	LI
		Liparidae	<i>Liparis atlanticus</i>	Atlantic seasnail	Limace atlantique	503	159524	34	LR
			<i>Liparis fabricii</i>	Gelatinous snailfish	Limace gélatineuse	505	127218	27	LR
			<i>Liparis gibbus</i>	Variegated snailfish	Limace marbée	512	159526	41	LR
			<i>Careproctus reinhardti</i>	Sea tadpole	Petite limace de mer	520	127212	18	LR
<hr/>									
Perciformes									
		Zoarcidae	<i>Lycenchelys verrillii</i>	Wolf eelpout	Lycode à tête longue	603	159258	40	LR
<hr/>									
Anguilliformes									
		Nemichthyidae	<i>Nemichthys scolopaceus</i>	Slender snipe eel	Avocette ruban	604	126306	28	LR
<hr/>									
Perciformes									
		Ammodytidae	<i>Ammodytes dubius</i>	Sand lance	Lançon	610	151520	1283	LI
		Zoarcidae	<i>Lycodes terraenovae</i>	Newfoundland eelpout	Lycode du Labrador	619	127117	64	LR
			<i>Lycodes lavalaei</i>	Newfoundland eelpout	Lycode du Labrador	620	127107	72	LR
		Pholidae	<i>Pholis gunnellus</i>	Rock gunnel	Sigouine de roche	621	126996	21	LR
		Stichaeidae	<i>Lumpenus lampretaeformis</i>	Snakeblenny	Lompénie-serpent	622	154675	423	LI
			<i>Leptoclinus maculatus</i>	Daubed shanny	Lompénie tachetée	623	127072	443	LI
			<i>Ulvaria subbifurcata</i>	Radiated shanny	Ulvaire deux-lignes	625	159821	145	LR
			<i>Eumesogrammus praecisus</i>	Fourline snakeblenny	Quatre-lignes atlantique	626	159817	40	LR
		Cryptacanthodidae	<i>Cryptacanthodes maculatus</i>	Wrymouth	Terrassier tacheté	630	159675	120	LR
		Callionymidae	<i>Foetorepus agassizii</i>	Spotfin dragonet	Dragonnet tacheté	637	276339	20	LR

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category
Zeiformes	Zoarcidae	Zoarcidae	<i>Zoarces americanus</i>	Ocean pout	Loquette d'Amérique	640	159267	1478	LF
			<i>Lycodes reticulatus</i>	Arctic eelpout	Lycode arctique	641	127112	70	LR
			<i>Melanostigma atlanticum</i>	Atlantic soft pout	Molasse atlantique	646	127120	43	LR
			<i>Lycodes vahlii</i>	Vahl's eelpout	Lycode à carreaux	647	127118	565	LI
	Stromateidae		<i>Peprilus triacanthus</i>	Atlantic butterfish	Stomaté fossette	701	159828	487	LI
Aulopiformes	Zeidae		<i>Zenopsis conchifer</i>	Silvery John dory	Saint Pierre argenté	704	127426	39	LR
	Paralepididae		<i>Arctozenus risso</i>	White barracudina	Lussion blanc	712	126352	196	LR
Beloniformes	Scomberesocidae	Scomberesocidae	<i>Scomberesox saurus</i>	Atlantic saury	Balaou atlantique	720	126392	37	LR
			<i>Sternopychidae</i>	Hatchetfishes	Haches d'argent	741	125603	21	LR
Lophiiformes	Ogcocephalidae	Ogcocephalidae	<i>Dibranchus atlanticus</i>	Atlantic batfish	Malthe atlantique	742	126558	18	LR
			<i>Synphurus diomedeanus</i>	Spottedfin tonguefish	Langue fil noir	816	159358	24	LR
Scorpaeniformes	Cottidae	Cottidae	<i>Artediellus atlanticus</i>	Atlantic hookear sculpin	Hameçon atlantique	880	127193	258	LI
			<i>Dipturus laevis</i>	Barndoor skate	Grande raie	200	158548	246	LI
Elasmobranchii	Rajidae	Rajidae	<i>Amblyraja radiata</i>	Thorny skate	Raie épineuse	201	105865	3937	LF
			<i>Malacoraja senta</i>	Smooth skate	Raie lisse	202	158554	1773	LF

Class	Order	Family	Scientific name	English name	French name	Species code	AphiaID	Num. records	Category	
Squaliformes			<i>Leucoraja erinacea</i>	Little skate	Raie hérisson	203	158551	712	LI	
			<i>Leucoraja ocellata</i>	Winter skate	Raie tachetée	204	158553	1180	LF	
	Squalidae	<i>Squalus acanthias</i>	Piked dogfish	Aiguillat commun	220	105923	1985	LF		
		<i>Centroscyllium fabricii</i>	Black dogfish	Aiguillat noir	221	105906	31	LR		
Cephalopoda										
<i>Oegopsida</i>										
<i>Myopsida</i>	Ommastrephidae	<i>Illex illecebrosus</i>	Northern shortfin squid	Encornet rouge nordique	4511	153087	4836	LF		
	Loliginidae	<i>Doryteuthis pealeii</i>	Longfin inshore squid	Calmar totam	4512	574541	96	LR		
Malacostraca										
<i>Decapoda</i>										
<i>Oregoniidae</i>	Pandalidae	<i>Pandalus borealis</i>	Northern prawn	Crevette nordique	2211	107649	718	SF		
	Cancridae	<i>Cancer borealis</i>	Jonah crab	Tourteau jona	2511	158056	1387	SF		
		<i>Cancer irroratus</i>	Atlantic rock crab	Tourteau poïnclos	2513	158057	788	SF		
		<i>Hyas coarctatus</i>	Arctic lyre crab	Crabe Hyas coarctatus	2521	107323	711	SF		
	Lithodidae	<i>Lithodes maja</i>	Atlantic king crab	Crabe épineux du nord	2523	107205	531	SF		
		<i>Chionoecetes opilio</i>	Queen crab	Crabe des neiges	2526	107315	1546	SF		
		<i>Hyas araneus</i>	Great spider crab	Crabe lyre araignée	2527	107322	625	SF		
	Geryonidae	<i>Chaceon quinquedens</i>	Red deepsea crab	Crabe rouge	2532	158407	33	SR		
	Nephropidae	<i>Homarus americanus</i>	American lobster	Homard américain	2550	156134	1623	SF		

307 **2.4 Analyses**

308 The Oracle relational database where all survey data are stored and archived is accessible from
309 the Bedford Institute of Oceanography in Dartmouth, Nova Scotia. Queries written in Structured
310 Query Language (SQL) are used to extract the data from the production server and to create the
311 data products used in all subsequent analyses. Catch records classified as "valid" (i.e. coming
312 from a representative tow without damage to the net) are used in the current analyses. To make
313 the available samples comparable, catch number and weight for each species was standardized
314 for the distance towed.

315 All data processing and analyses were conducted using the R software (R Core Team 2020)
316 using packages gstat (Pebesma 2004), PBSmapping (Schnute et al. 2019), RODBC (Ripley
317 and Lapsley 2019), spatstat (Baddeley 2015), maptools (Bivand and Lewin-Koh 2020), rgeos
318 (Bivand and Rundel 2020), classInt(Bivand 2020), RColorBrewer(Neuwirth 2014), MASS (Ripley
319 et al. 2020), worms (Holstein 2018), and tidyverse (Wickham 2019). The present document is
320 rendered as a Technical Report using the csasdown R package developed and maintained by
321 Fisheries and Oceans Canada scientists (Anderson et al. In press).

322 **2.4.1 Geographic distribution of catches**

323 Spatial interpolation of catch biomass (kg/tow) or abundance (number/tow) was done using a
324 weighting inversely proportional to the distance (inverse-distance weighted, IDW), using function
325 "idw" of the spatstat R package (Baddeley 2015). The IDW method was used with a power
326 parameter value of 10.

327 **2.4.2 Abundance and biomass indices**

328 For each species, stratified random estimates of catch abundance and biomass (Smith 1996)
329 were computed for each year. Yearly estimates of the standard error were also computed.

330 **2.4.3 Distribution indices**

331 For each Category L, I and S fish species, the minimum area required to account for 75% and
332 95% of the total biomass or abundance were computed (D75% and D95%). These measures of
333 distributions were computed for each year by using the Lorenz curve of mean stratum-level catch
334 estimates and the area of occupied strata (Swain and Sinclair 1994; Swain and Morin 1996).

335 **2.4.4 Length frequencies**

336 The length frequency distribution of catch (the stratified numbers-at-length) is tabulated for each
337 seven-year period (1970-2009), and last ten-year period (2010-2020).

338 **2.4.5 Length-weight relationship and condition factor**

339 The relationship between the weight and the length of fish was estimated using the following
340 non-linear isometric relationship:

$$W = \alpha L^\beta$$

341 where W is the total weight (g), L is the length (cm), and, α and β are the parameters to be
342 estimated.

343 Average fish condition (C) was computed as:

$$C = \frac{W}{\alpha L^\beta}$$

344

345 **2.4.6 Depth, temperature and salinity distribution of catches**

346 For each category L species, We followed the methods developed by (Perry and Smith 1994)
347 and generated cumulative frequency distributions of depth, temperature and salinity of survey
348 catches.

349 **2.4.7 Density-dependent habitat selection**

350 We followed the methods of (Myers and Stokes 1989) to evaluate how fish abundance in each
351 stratum varied with overall temporal fluctuations of population abundance.

352 For each category L species, we fitted a model of the relationship between stratum-level density
353 and overall abundance (the yearly stratified random estimate of abundance, defined above).
354 To properly use the observations of zero catch while accounting for the logarithmic distribution
355 of catch abundance, we implemented the model as a generalised linear using a log link and a
356 Poisson error distribution:

$$Y_{h,i} = \alpha_h Y_i^{\beta_h}$$

357 where, $y_{h,i}$ is the average abundance of stratum h in year i , and $\alpha_{h,i}$ and $\beta_{h,i}$ are the fitted
358 parameters. The estimated parameter $\beta_{h,i}$ is referred to as the “slope parameter” and indicates
359 whether stratum-level density is positively ($\beta_{h,i} <= 0$), negatively ($\beta_{h,i} >= 0$) or negligibly
360 ($\beta_{h,i} \approx 0$) related to population abundance.

361 To estimate the suitability of each stratum, the median abundance observed during the years
362 that are in the top 25% of yearly estimates is used. We combine the slope parameter estimates
363 from the above model with the median abundance to identify strata that have consistently high
364 abundance and whose local density is weakly related to fluctuation in population abundance
365 ($\beta_{h,i} \approx 0$). Preferred strata are identified for each category L species.

366

3 Results

367 The plots generated for each species are presented in the Appendix.

368 **3.1 Description of Figures**

369 **3.1.1 Type A**

370 For Category L and S species:

371 Spatial distribution of catch-per unit of effort, (CPUE, kilograms per tow) in July-August for the
372 Bay of Fundy and Scotian Shelf in five-year periods. Spatial interpolation between tows was
373 done using Inverse Distance Weight (IDW). The probability of occurrence (proportion of tows with
374 catch records for a given species) was also reported for each five-year period.

375 For Category LR and SR:

376 Location of tows with catch over the period 1970-2012 (Type LR) or the period 1999-2012 (Type
377 SR). Location of tows with catch over the period 1970-2012 (Type LR) or the period 1999-2012
378 (Type SR).

379 **3.1.2 Type B**

380 For Category L, S and I species:

381 Stratified random estimate of CPUE (left panel), distribution indices (D75% and D95%, the
382 minimum area containing 75% and 95% of biomass, middle panel), and distribution vs. weight
383 per tow (right panel). The stratified random mean is plotted as a solid line with the 95%
384 confidence region indicated by the solid grey line. The overall mean is plotted as a grey
385 horizontal line and the overall mean plus or minus 50% of the standard deviation appear as
386 horizontal dashed lines. In all three panels, the early years appear in blue and the last years
387 appear in red. The predictions from a loess estimator are overlaid on the distribution indices
388 (middle panel). The Pearson correlation coefficient between D75% and biomass, and its
389 statistical significance, are also reported in the right panel.

390 **3.1.3 Type C.**

391 Length frequency distribution for NAFO divisions 4X and 4VW. A smoothed length frequency
392 distribution is shown for each 7-year periods covered by the surveys.

393 **3.1.4 Type D.**

394 Average fish condition for all fish lengths (black dots and black line), large fish (thick gray line),
395 and small fish (thin gray line). Fish condition is presented for NAFO divisions 4VW (right panel)
396 and 4X (left panel).

397 **3.1.5 Type E.**

398 Cumulative frequency distributions of depth, temperature and salinity at all sampled locations
399 (thick solid line) and at fishing locations with catch records (thin dashed line). The depth,
400 temperature and salinity associated with 5%, 25%, 50%, 75% and 95% of the cumulative catch is
401 shown in tabular fashion on the bottom right panel.

402 **3.1.6 Type F.**

403 Slopes estimates from the density-dependent habitat selection model (y axis) plotted versus
404 the median abundance during the top 25% of years. The red box indicates strata of particular
405 importance for a species by identifying slopes that are within a standard error from zero and that
406 are within the top 25% of median abundance. Each stratum is identified on the plot by the last
407 two digits of its number.

408 **3.2 Summary of successful tows by year and stratum**

Table 4. Number of representative tows conducted in each stratum during the period 1970 to 1989.

Stratum	NAFO Div.	Area (km2)	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
440	4VN	3173.016	4	2	2	3	3	3	3	3	3	3	3	3	3	3	3	4	5	5	6	4
441	4VN	3434.000	4	2	2	3	3	3	1	3	3	3	3	3	3	3	3	5	5	4	4	4
442	4VN	4934.658	3	2	2	2	3	3	2	3	3	3	3	3	3	3	3	3	5	6	7	5
443	4VSW	4526.012	4	2	4	4	8	3	1	2	4	4	4	3	5	4	4	4	6	6	5	2
444	4VSW	13478.450	3	2	5	4	6	4	6	7	4	4	4	5	5	6	4	4	6	6	3	6
445	4VSW	3512.982	5	2	5	4	5	5	1	3	4	4	4	5	5	3	4	5	6	4	4	4
446	4VSW	1686.094	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	4	3	3	3	3
447	4VSW	5549.344	4	2	6	5	7	4	4	3	4	4	5	4	4	4	4	5	7	6	6	6
448	4VSW	4975.866	5	2	5	4	5	4	4	4	4	4	4	6	4	4	4	5	5	5	5	5
449	4VSW	494.496	2	2	2	2	3	2	2	2	1	2	2	2	1	2	2	2	2	2	2	2
450	4VSW	1315.222	2	2	3	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
451	4VSW	504.798	1	2	2	2	2	2	2	2	2	3	2	2	3	2	2	2	2	2	2	2
452	4VSW	1184.730	2	2	2	2	2	2	2	2	2	2	2	4	2	2	2	2	2	2	3	2
453	4VSW	889.406	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	2	2
454	4VSW	1713.566	3	2	3	3	3	3	3	3	3	3	3	2	3	3	3	3	3	3	2	2
455	4VSW	7286.948	7	6	7	6	7	6	6	7	7	7	7	7	7	7	7	7	8	8	7	7
456	4VSW	3279.470	5	4	6	5	5	6	4	6	6	6	6	7	6	6	6	6	6	7	6	6
457	4VSW	2784.974	2	2	2	2	3	2	2	2	2	2	2	3	2	2	2	2	2	2	4	2
458	4VSW	2259.572	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	5	5	3
459	4VSW	10810.232	3	2	4	4	4	4	4	4	4	4	4	4	3	4	4	6	6	5	6	5
460	4VSW	4615.296	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2	4	3	3
461	4VSW	3962.836	3	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3	3	2	2
462	4VSW	7266.344	3	3	4	3	4	4	4	4	4	4	4	6	4	4	4	4	6	5	4	4
463	4VSW	1037.068	2	2	2	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2
464	4VSW	4453.898	4	3	5	3	3	6	5	5	5	5	5	5	5	4	5	5	7	6	5	5
465	4VSW	8183.222	6	5	5	4	5	4	5	5	5	5	5	7	6	5	5	5	5	8	8	8
466	4VSW	776.084	2	2	3	2	3	3	3	3	3	3	3	2	3	3	3	3	3	3	2	2
470	4X	3159.280	1	2	2	2	3	2	2	2	2	2	2	2	2	2	2	2	2	2	3	3
471	4X	3447.736	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
472	4X	4289.066	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	4	4
473	4X	910.010	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
474	4X	552.874	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	0	2	2	2	2
475	4X	535.704	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
476	4X	5075.452	2	2	2	2	2	2	3	2	2	2	2	1	2	2	2	2	2	2	4	4
477	4X	4230.688	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2	5	4	4
478	4X	800.122	2	2	3	2	3	3	3	3	2	3	3	3	3	3	3	3	3	3	2	2
480	4X	2249.270	4	4	4	3	3	3	4	4	3	4	3	3	3	4	4	4	4	4	4	4
481	4X	6438.750	5	3	4	4	4	3	4	4	5	4	3	4	4	4	4	4	4	6	7	6
482	4X	3578.228	2	1	2	2	2	2	2	3	2	2	3	2	2	2	2	2	2	2	3	3
483	4X	1826.888	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2	2
484	4X	7774.576	2	2	3	3	3	3	3	3	2	3	3	3	3	4	3	3	3	4	4	4
485	4X	5432.588	2	2	2	3	3	3	3	3	3	3	3	2	3	4	3	3	3	6	7	6
490	4X	2063.834	2	2	2	2	2	3	3	3	3	3	3	2	3	3	3	3	3	3	4	4
491	4X	2359.158	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	4	4	4
492	4X	3729.324	3	2	4	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	4	4
493	4X	1830.322	1	2	3	3	3	3	3	3	3	3	3	2	3	3	3	3	3	3	3	3
494	4X	1431.978	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2

Stratum	NAFO Div.	Area (km2)	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
495	4X	2005.456 171809.888	2 134	2 110	2 146	2 134	2 153	2 143	2 135	2 144	1 141	2 147	2 145	2 150	2 150	2 146	2 143	2 152	2 171	2 188	2 177	2 170

Table 5. Number of representative tows conducted in each stratum during the period 1990 to 2009.

Stratum	NAFO Div.	Area (km2)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
440	4VN	3173.016	4	4	4	3	4	4	4	4	4	4	6	4	4	4	4	4	4	4	3	4
441	4VN	3434.000	6	5	5	5	5	5	5	5	6	7	6	6	6	7	6	6	6	5	6	
442	4VN	4934.658	5	5	6	5	6	6	6	6	6	7	6	6	5	6	6	7	5	5	5	6
443	4VSW	4526.012	4	2	4	3	3	4	4	5	5	4	5	4	5	5	5	5	4	4	5	4
444	4VSW	13478.450	7	8	8	9	6	8	8	7	8	8	9	10	9	9	9	8	10	8	6	9
445	4VSW	3512.982	4	4	4	5	7	4	4	4	3	3	6	5	5	5	5	6	5	4	3	6
446	4VSW	1686.094	3	3	3	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
447	4VSW	5549.344	8	7	7	7	7	7	6	7	7	6	7	7	7	7	7	7	6	6	4	6
448	4VSW	4975.866	9	6	6	7	7	7	6	7	6	7	8	8	8	8	7	8	8	6	5	7
449	4VSW	494.496	2	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2
450	4VSW	1315.222	3	3	3	3	3	3	3	3	2	3	3	3	3	3	3	3	3	3	3	3
451	4VSW	504.798	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
452	4VSW	1184.730	3	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2	2	2	2
453	4VSW	889.406	3	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	3	2
454	4VSW	1713.566	3	2	2	2	2	2	2	3	2	2	2	2	2	2	2	3	2	2	2	2
455	4VSW	7286.948	12	10	10	9	10	10	10	13	8	11	11	11	11	11	11	12	11	7	5	8
456	4VSW	3279.470	10	7	7	8	8	8	8	8	6	8	10	8	8	8	8	8	8	6	2	7
457	4VSW	2784.974	4	2	2	2	2	2	2	2	2	1	4	2	2	2	2	2	2	2	2	2
458	4VSW	2259.572	9	8	8	8	8	8	7	8	5	6	10	8	7	8	8	10	8	5	2	7
459	4VSW	10810.232	5	5	6	4	6	6	4	5	6	6	8	6	6	6	6	6	6	6	5	3
460	4VSW	4615.296	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	4	3	2	3
461	4VSW	3962.836	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	4	2	2	2
462	4VSW	7266.344	5	5	4	4	4	4	4	4	4	4	4	4	4	4	4	4	5	4	3	4
463	4VSW	1037.068	3	2	2	2	2	2	2	2	2	2	2	2	2	3	2	2	4	2	2	2
464	4VSW	4453.898	9	7	7	7	7	7	4	7	7	7	7	7	7	7	5	8	7	6	4	5
465	4VSW	8183.222	12	9	10	10	10	10	10	10	9	10	10	10	10	10	10	10	10	7	8	7
466	4VSW	776.084	3	2	2	2	2	2	3	2	2	3	2	2	2	2	2	2	2	1	3	2
470	4X	3159.280	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
471	4X	3447.736	2	2	2	2	2	2	2	1	2	2	2	2	2	2	2	2	2	2	2	2
472	4X	4289.066	6	4	4	4	4	4	4	3	4	4	4	4	4	4	4	4	4	4	3	4
473	4X	910.010	3	2	2	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2
474	4X	552.874	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
475	4X	535.704	2	2	2	2	2	2	2	2	2	2	2	2	2	3	2	2	2	2	2	2
476	4X	5075.452	4	4	4	4	4	4	4	4	4	4	4	4	4	4	5	4	4	4	4	4
477	4X	4230.688	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	8	5	5	5
478	4X	800.122	2	2	2	2	2	3	3	2	2	2	2	2	2	2	2	2	3	2	2	2
480	4X	2249.270	8	8	8	8	8	8	8	8	8	8	7	8	8	8	7	9	8	6	8	8
481	4X	6438.750	8	9	9	9	9	7	9	9	9	9	8	9	9	8	9	6	12	9	7	8
482	4X	3578.228	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	2	4	3	3	3

Stratum	NAFO Div.	Area (km2)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
483	4X	1826.888	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
484	4X	7774.576	3	3	3	3	3	3	3	3	3	3	3	3	4	3	3	4	4	3	3	4
485	4X	5432.588	2	3	3	3	3	3	3	3	3	3	4	3	5	5	3	2	5	4	5	5
490	4X	2063.834	4	4	4	4	4	5	4	4	4	3	4	4	4	6	4	3	3	3	4	3
491	4X	2359.158	3	3	3	3	3	3	3	3	3	3	3	3	3	5	3	3	4	3	4	4
492	4X	3729.324	3	3	3	3	3	2	3	3	3	3	3	3	3	5	2	3	4	4	4	4
493	4X	1830.322	3	3	3	3	3	3	2	3	3	2	3	3	4	5	2	4	4	3	3	4
494	4X	1431.978	2	2	2	2	2	2	2	2	2	2	2	2	2	3	4	2	2	4	3	3
495	4X	2005.456	2	2	2	2	2	2	2	2	2	2	2	2	2	2	4	2	2	5	3	3
171809.888		213	189	193	190	195	195	191	193	186	191	213	201	208	216	188	222	209	177	165	196	

Table 6. Number of representative tows conducted in each stratum during the period 2010 to 2020 and for the whole 1970 to 2020 period.

Stratum	NAFO Div.	Area (km2)	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	Total
440	4VN	3173.016	4	5	4	4	4	4	4	4	0	5	4	190
441	4VN	3434.000	6	7	6	6	6	6	6	6	0	7	4	238
442	4VN	4934.658	5	6	6	6	6	6	6	6	0	6	5	240
443	4VSW	4526.012	4	6	5	5	3	7	4	5	0	9	4	214
444	4VSW	13478.450	11	13	9	8	9	9	11	10	0	6	8	352
445	4VSW	3512.982	4	7	2	4	3	4	4	4	0	6	3	215
446	4VSW	1686.094	3	4	3	3	3	2	3	2	0	3	2	145
447	4VSW	5549.344	6	8	6	7	7	7	7	7	0	6	5	291
448	4VSW	4975.866	7	10	8	8	8	7	6	6	0	7	4	299
449	4VSW	494.496	2	4	2	2	2	2	2	2	0	2	2	100
450	4VSW	1315.222	3	3	3	3	3	3	3	2	0	3	2	144
451	4VSW	504.798	2	2	2	2	2	2	2	2	0	2	2	104
452	4VSW	1184.730	2	2	2	2	1	4	3	3	0	3	3	110
453	4VSW	889.406	2	1	3	2	3	2	2	1	0	2	2	116
454	4VSW	1713.566	2	4	2	2	2	2	2	2	0	3	2	121
455	4VSW	7286.948	10	10	10	11	11	9	9	8	0	9	6	429
456	4VSW	3279.470	7	9	8	8	6	5	6	6	0	6	4	331
457	4VSW	2784.974	2	4	2	2	2	3	3	3	0	3	2	113
458	4VSW	2259.572	6	9	8	6	4	5	5	5	0	6	3	269
459	4VSW	10810.232	6	7	6	6	6	7	7	6	0	9	7	262
460	4VSW	4615.296	3	4	4	3	3	5	5	5	3	6	5	151
461	4VSW	3962.836	2	3	3	2	2	3	3	3	2	3	3	113
462	4VSW	7266.344	4	6	4	4	5	5	5	5	0	5	5	212
463	4VSW	1037.068	2	3	2	2	2	3	2	2	0	2	2	107
464	4VSW	4453.898	6	7	7	7	7	6	6	4	0	6	4	288
465	4VSW	8183.222	8	10	10	10	10	10	9	7	3	10	7	397
466	4VSW	776.084	2	2	2	2	2	2	2	3	0	3	2	118
470	4X	3159.280	2	2	3	2	2	3	3	3	4	3	2	112
471	4X	3447.736	2	2	3	2	2	3	3	3	4	4	3	110

Stratum	NAFO Div.	Area (km2)	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	Total
472	4X	4289.066	4	6	4	4	4	4	4	4	4	4	4	172
473	4X	910.010	2	2	2	2	2	2	2	2	2	2	2	104
474	4X	552.874	2	2	2	2	2	2	2	2	2	2	2	100
475	4X	535.704	2	2	2	2	2	2	2	2	2	2	2	103
476	4X	5075.452	4	4	4	4	4	5	5	5	5	5	5	177
477	4X	4230.688	5	4	5	5	6	5	5	4	4	6	4	204
478	4X	800.122	2	2	2	2	2	2	2	3	2	2	2	119
480	4X	2249.270	8	7	8	8	6	7	7	7	5	7	5	306
481	4X	6438.750	8	10	9	9	9	8	10	9	6	9	6	350
482	4X	3578.228	3	4	3	3	3	3	4	4	3	4	3	141
483	4X	1826.888	2	3	2	2	2	2	3	3	2	3	2	105
484	4X	7774.576	3	5	5	5	4	6	5	7	7	7	7	186
485	4X	5432.588	5	6	5	5	5	6	6	6	4	6	5	196
490	4X	2063.834	3	4	2	4	3	4	4	4	3	4	3	173
491	4X	2359.158	4	4	4	4	4	4	4	4	3	4	3	168
492	4X	3729.324	4	6	4	4	4	3	4	4	3	4	4	171
493	4X	1830.322	3	4	4	4	3	3	4	6	3	3	3	159
494	4X	1431.978	4	4	4	4	3	4	4	3	2	4	3	128
495	4X	2005.456	3	4	4	4	2	4	4	4	3	4	3	127
		171809.888	196	243	210	208	196	212	214	208	81	227	175	9080

409 A total of 9080 representative tows were conducted for the period spanning from 1970 to 2020.

410

4 Discussion

411 This report builds on previous work and former atlases by updating a comprehensive suite of
412 indices to give a snapshot of population status and environmental preferences of 104 fish and
413 invertebrate species. The current document is not meant to replace stock assessments, species-
414 specific analyses of abundance, biomass and distribution, or any targeted attempts to integrate
415 information about species or group of species from the wide and disparate sources of data about
416 marine organisms in the area covered by the DFO Maritimes summer trawl survey. It is rather
417 meant to provide a reproducible set of tools to extract and visualize the information collected
418 in the summer groundfish research vessel survey. It is hoped that this document can provide a
419 stepping stone to conduct other ecological analyses using the trawl survey data and increase
420 reproducibility and transparency of ecological information collected annually.

421 **4.1 Diversity of approaches used for mapping fish and invertebrates in the Scotian Shelf
422 bioregion**

423 Different methods have been applied in the Northwest Atlantic, and specifically on the Scotian
424 Shelf bioregion, to map fish and invertebrate species distribution. The present report, for
425 example, builds upon the atlas of important habitat developed to map the persistence of relatively
426 high biomass for key fish species using the summer groundfish research vessel survey (Horsman
427 and Shackell 2009). Important habitat was obtained by interpolating observed weight per each
428 species using the IDW, and calculating areas with relatively persistent high biomass for periods
429 representing different fishery management eras. To compliment information from this atlas,
430 including additional representations of biomass and diversity, a similar IDW interpolation mapping
431 procedure was followed by Smith et al. (2015), Ward-Paige and Bundy (2015), and Bundy et al.
432 (2017). The summer groundfish research vessel survey is typically conducted during the month
433 of July. However, from the fall of 1978 through to the spring of 1985, DFO also conducted spring
434 and fall surveys using the same sampling design. This unique seasonal data was used to map
435 the seasonal spatial distribution of key demersal and other fish species using IDW interpolation
436 on the Scotian Shelf from the spring, summer and fall between 1978 and 1985 (Smith et al.
437 2015). Following recommendations provided by Kenchington and Kenchington (2017), the spatial
438 distribution of three indicators of biodiversity for fish and invertebrates were mapped using IDW
439 interpolation to identify areas with persistently high values across fishery management eras,
440 and compared with areas of persistently high abundance for selected species (Ward-Paige and
441 Bundy 2015). This analysis revealed a lack of consistent relationships between areas of persist
442 high diversity and persistent high biomass, suggesting that both can be used as independent
443 and important spatial indicators of the system (Ward-Paige and Bundy 2015). Groupings of
444 fishes and invertebrates based on size, habitat and feeding guild, were also mapped using
445 IDW interpolations to identify hotspots of functional group diversity (Bundy et al. 2017). This
446 analysis revealed a spatially and temporally variable distribution of functional diversity across
447 the Scotian Shelf with notable areas of high and low diversity (Bundy et al. 2017). Top quintiles
448 of each functional group using the IDW approach were used as representative layers for fish

449 and invertebrates in the MPA Network design in the Scotian Shelf Bioregion (Serdynska et al. In
450 press). IDW interpolation methods have also been used to map the distribution of individual
451 species such as sea cucumbers (*Cucumaria frondosa*) in the Scotian Shelf bioregion (Shackell et
452 al. 2013), and sea scallop (*Placopecten magellanicus*) in Georges and Browns Bank (Hubley et
453 al. 2014).

454 Species Distribution Modelling (SDM), instead of IDW, can also be used to evaluate spatio-
455 temporal dynamics by predicting and understanding past, present and future distribution
456 of species using environmental predictors (Robinson et al. 2017). A variety of modelling
457 approaches are being implemented in Maritimes Region to map and predict fish and invertebrate
458 species distribution by incorporating environmental predictors to account for seasonal and
459 temporal variability. For example, a stock assessment of snow crab (*Chionoecetes opilio*) on
460 the Scotian Shelf used data from the snow crab survey from 2005 to 2018 to map spatial data
461 products for this stock, including annual predicted interpolations of potential habitat using
462 Generalized Additive Models (GAM) and several environmental covariates including depth,
463 curvature, slope, species composition, and annual temperature (Zisserson et al. 2019). Sea
464 scallop predicted habitat using Maximum Entropy (MaxEnt) models were computed in the
465 German Bank using data compiled via benthic habitat mapping and seafloor geotechnical
466 surveys in 2006, 2009, and 2010 (Brown et al. 2012). Predictions in the Scotian Shelf bioregion
467 and the Northeast United States using datasets from DFO and the National Oceanic and
468 Atmospheric Administration from 1993 to 2012 also predicted sea scallop habitat at a wider scale
469 based on three scenarios of seasonal temperature and salinity climatologies (NOAA) (Lowen
470 et al. 2019). Offshore American lobster stock assessments (*Homarus americanus*) used data
471 from the RV, DFO Georges Bank, and National Marine Fisheries Service (NMFS) Northeast
472 Fisheries Science Center (NEFSC) bottom trawl surveys (1970 to 2015) to predict species
473 distribution using boosted regression trees and several environmental predictors (bathymetry,
474 slope, curvature, and annual temperature interpolations) (Cook et al. 2017). Information on
475 the potential for recovery of cusk (*Brosme brosme*) used data from the bottom longline Halibut
476 industry survey and Cusk absences in the Summer groundfish research vessel survey from
477 1998-2013 to predict suitable habitat using GAM, MaxEnt, and random forest models and
478 several physical environmental variables (e.g. complexity, benthic current stress and complexity,
479 temperature, salinity, primary production, chlorophyll, suspended matter) (Harris et al. 2018).
480 Atlantic halibut (*Hippoglossus hippoglossus*) assessments using Summer groundfish research
481 vessel survey and NOAA survey data from 2001 to 2013 predicted juvenile habitat using MaxEnt
482 model and environmental predictors (bathymetry, slope, bottom temperature) (French et al.
483 2018). Persistent areas of high Atlantic halibut juvenile abundance were predicted using data
484 from 27 bottom trawl surveys combined (NMFS and DFO) from 1978 to 2013 and applying
485 Bayesian hierarchical spatiotemporal models with two environmental predictors (depth and
486 temperature) (Boudreau et al. 2017).

487 These examples of mapping efforts in Maritimes Region showcase the diversity of approaches
488 relevant to a variety of important research questions and management applications. Approaches,
489 methods, datasets, and environmental predictors are selected based on individual project
490 research questions, and considerations for each species, communities or stock. This allows
491 research groups to maintain innovation and keep up with emerging methods and technologies to
492 improve assessments, predictions, and ultimately, science advice. The diversity of approaches
493 also leads to complexity when looking across studies as each data compilation and predictive
494 method carries its own independent assumptions and can lead to different spatial outputs.

495 **4.2 Interpreting spatial results for marine spatial planning purposes**

496 Fisheries and Oceans Canada is leading a marine spatial planning process that brings together
497 relevant authorities and stakeholders to better coordinate how we use and manage marine
498 spaces to achieve ecological, economic and social objectives. Operationalizing marine spatial
499 planning includes a series of steps, including the process of analyzing existing conditions
500 by collecting and mapping information about ecological, environmental and oceanographic
501 conditions (Ehler and Douvere 2009; Agardy et al. 2011). Mapping the distribution of species
502 is critical for the implementation of spatial management and as a first step in marine spatial
503 planning processes. Species distribution have supported the identification of important sites for
504 a given species or areas of high richness and diversity, which in turn can be used to inform siting
505 decisions of new activities such as Marine Protected Areas (MPA), aquaculture sites or wind
506 turbines. In the Scotian Shelf bioregion, mapping species distributions has been used to highlight
507 areas of high biological diversity to support the identification of Ecologically or Biologically
508 Significant Areas [Ricard and Shackell (2013); Ward-Paige and Bundy (2015)], to distinguish
509 important and persistent habitat of significant species and functional groups to support MPA and
510 conservation planning (Horsman and Shackell 2009; Smith et al. 2015; Ward-Paige and Bundy
511 2015; Bundy et al. 2017), to identify important habitat for Species at Risk (Harris et al. 2018) and
512 to highlight reserves for data-poor invertebrate fisheries (Shackell et al. 2013). Mapping species
513 distribution has also been used to illustrate multi-decadal scale projections of changes in species
514 distribution in the context of climate change and adaption (Stanley et al. 2018; W. et al. 2019).

515 In support of the marine spatial planning process, a public web-based atlas with relevant
516 geospatial information is being developed to support decision-making. This Atlantic Canada-
517 wide compilation of data and information will be a web-based, public platform with interactive
518 maps of ocean ecosystems, human uses and management areas. This atlas cannot host the
519 vast diversity of products and mapping approaches available in Maritimes Region. Consequently,
520 we recommend that data products presented in this report should not be used for the atlas until
521 an evaluation of the spatial information available and used in the past, is conducted.

522 This diverse portfolio of approaches and applications is not unique to the Maritimes Region. A
523 recent review of global distribution modelling efforts recommended the adoption of a consistent
524 framework that integrates multi-model approaches and a clear expression of errors and
525 uncertainties (Robinson et al. 2017). In this context, Pacific Region has developed two initiatives
526 to enable consistency and frequent publication, reproducibility, and transparency. One initiative
527 developed a fully automated reproducible report to give a synthesis of data availability, population
528 trends, fishing trends, growth and maturity patterns for 113 groundfish species in British
529 Columbia to support stock assessment (Anderson et al. 2019). The second initiative developed
530 a SDM framework that was applied to twelve species on Canada's Pacific coast as part of the
531 Regional Response Plan (Nephin et al. 2019). The Maritimes and Gulf region, through this and
532 past reports, are also using similar reproducible approaches to facilitate annual updates and
533 transparency (Ricard and Shackell 2013; Ricard and Gomez 2021).

534 Recognizing the diversity of approaches for mapping fish and invertebrates in the Scotian Shelf
535 bioregion, we recommend the development of a regional community of practice to compare
536 and evaluate approaches for mapping, interpolating and/or modelling fish and invertebrates
537 so future publications and advice related to spatial outputs can lead to more comparable work
538 and consistent science advice to support processes such as marine spatial planning. At the

539 international level, guidelines and standards related to appropriate variables and methods
540 for mapping and modeling species and communities of deep-sea habitats were proposed to
541 encourage the production of publications that will lead to more comparable work (Kenchington et
542 al. 2019). Similar general guidance for group practice approach mapping would be a worthwhile
543 product in Maritimes Region. Until then, we proposed the use of the Open Data record (DFO
544 2021) for the version 1.0 of the public web-based atlas.

545

5 Acknowledgements

546 We thank all the dedicated personnel involved in running trawl surveys in the Maritimes Region
547 and the numerous colleagues in Maritimes Region that have shared information and advice in
548 support of this report. The efforts of the Gulf Region secondary publications coordinators Alicia
549 Cassidy and Jeff Clements in getting this report published are well appreciated.

6 References

- 552 Agardy, T., Notarbartolo di Sciara, G., and Christie, P. 2011. [Addressing the shortcomings of](#)
 553 [marine protected areas through large scale marine spatial planning](#). Marine Policy 35(2):
 554 226–232.
- 555 Anderson, S.C., Grandin, C., Edwards, A.M., Grinnell, M.H., Ricard, D., and Haigh, R. (In
 556 press). Csasdown: Reproducible CSAS Reports with Bookdown.
- 557 Anderson, S.C., Keppel, E.A., and Edwards, A.M. 2019. [A reproducible data synopsis for](#)
 558 [over 100 species of British Columbia groundfish](#). DFO Can. Sci. Advis. Sec. Res. Doc.
 559 2019/041. vii + 321 p.
- 560 Appeltans, W., Bouchet, P., Boxshall, G.A., De Broyer, C., Voogd, N.J. de, Gordon, D.P.,
 561 Hoeksema, B.W., Horton, T., Kennedy, M., J., M., Poore, G.C.B., Read, G., Stöhr,
 562 S., Walter, T.C., and Costello, M.J. (*Editors*). 2012. World register of marine species.
 563 Accessed at <http://www.marinespecies.org> on 2020-12-1.
- 564 Baddeley, R., A. 2015. Spatial point patterns: Methodology and applications
 565 with R. Chapman; Hall/CRC Press, London [http://www.crcpress.com/
 566 Spatial-Point-Patterns-Methodology-and-Applications-with-R/
 567 Baddeley-Rubak-Turner/9781482210200/](http://www.crcpress.com/Spatial-Point-Patterns-Methodology-and-Applications-with-R-Baddeley-Rubak-Turner/9781482210200/).
- 568 Benoît, H.P., Abgrall, M.-J., and Swain, D.P. 2003. [An assessment of the general status of](#)
 569 [marine and diadromous fish species in the southern Gulf of St. Lawrence based on annual](#)
 570 [bottom trawl surveys \(1971-2002\)](#). Can. Tech. Rep. Fish. Aquat. Sci. 2472: iv + 183 p.
- 571 Bivand, R. 2020. classInt: Choose univariate class intervals. R package version 0.4-3 <https://CRAN.R-project.org/package=classInt>.
- 573 Bivand, R., and Lewin-Koh, N. 2020. Maptools: Tools for handling spatial objects. R package
 574 version 1.0-2 <https://CRAN.R-project.org/package=maptools>.
- 575 Bivand, R., and Rundel, C. 2020. Rgeos: Interface to geometry engine - open source
 576 ('GEOS'). R package version 0.5-5 <http://CRAN.R-project.org/package=rgeos>.
- 577 Boudreau, S.A., Shackell, N.L., Carson, S., and Heyer C. E., den. 2017. [Connectivity,](#)
 578 [persistence, and loss of high abundance areas of a recovering marine fish population in](#)
 579 [the Northwest Atlantic Ocean](#). Ecol. Evol. 7: 9739–9749.
- 580 Bourdages, H., and Ouellet, J.-F. 2012. [Geographic distribution and abundance indices of](#)
 581 [marine fish in the northern Gulf of St. Lawrence \(1990-2009\)](#). Can. Tech. Rep. Fish.
 582 Aquat. Sci. 2963: vi + 171 p.
- 583 Brown, C., Sameoto, J., and Smith, S. 2012. [Multiple methods, maps, and management](#)
 584 [applications: Purpose made seafloor maps in support of ocean management](#). Journal of
 585 Sea research 72: 113.
- 586 Bundy, A., Will, E., Serdynska, A., Cook, A., and Ward-Paige, C.A. 2017. [Defining and](#)
 587 [mapping functional groups for fishes and invertebrates in the Scotian Shelf bioregion](#). Can.
 588 Tech. Rep. Fish. Aquat. Sci. 3186: iv + 49 p.

- 589 Clark, D.W., and Emberley, J. 2011. Update of the 2010 summer scotian shelf and bay of
590 fundy research vessel survey. Can. Tech. Rep. Fish. Aquat. Sci.: 1238: ix + 98 p.
- 591 Cook, A.M., Cassista Da-Ros, M., and Denton, C. 2017. [Framework Assessment of the](#)
592 [Offshore American Lobster in Lobster Fishing Area \(LFA\) 41.](#) ICES Journal of Marine
593 Science 2017/065 viii + 186 p.
- 594 DFO. 2016. DFO maritimes research vessel trawl surveys invertebrate observations. Version
595 7 in OBIS canada digital collections. Bedford Institute of Oceanography, Dartmouth, NS,
596 Canada, Published by OBIS, Digital 2016.
- 597 DFO. 2021. Maritimes research vessel surveys [dataset]. Retrieved from
598 <https://open.canada.ca/data/en/dataset/8ddcaeeaa-b806-4958-a79f-ba9ab645f53b>.
- 599 Doubleday, W.G., and Rivard, D. 1981. Bottom trawl surveys. Can. Spec. Publ. Fish. Aquat.
600 Sci: 58: 237 pp.
- 601 Ehler, C., and Douvere, F. 2009. Marine spatial planning: A step-by-step approach.
- 602 French, K., Shackell, N., and Heyer, N. den. 2018. [Information on the Potential for Recovery](#)
603 [of Cusk in Canadian Waters.](#) Fish. Bull. 116: 107–121.
- 604 Harris, L.E., Greenlaw, M., McCurdy, D., and MacDonald, D. 2018. [Information on the](#)
605 [Potential for Recovery of Cusk in Canadian Waters.](#) DFO Can. Sci. Advis. Sec.Res. Doc.
606 2018/002. vi + 62 p.
- 607 Holstein, J. 2018. Worms: Retriving aphia information from world register of marine species.
608 R package version 0.2.2 <https://CRAN.R-project.org/package=worms>.
- 609 Horsman, T.L., and Shackell, N.L. 2009. [Atlas of important habitat for key fish species of the](#)
610 [Scotian Shelf, Canada.](#) Can. Tech. Rep. Fish. Aquat. Sci. 2835: viii + 82 p.
- 611 Hubley, P.B., Reeves, A., Smith, S.J., and Nasmith, L. 2014. [Georges Bank 'a' and Browns](#)
612 [Bank 'North' Scallop \(*Placopecten magellanicus*\) Stock Assessment.](#) DFO Can. Sci.
613 Advis. Sec. Res. Doc. 2013/079. vi + 58 p.
- 614 Kenchington, E., Callery, O., Davidson, F., Grehan, A., Morato, T., Appiott, J., Davis, A.,
615 Dunstan, P., Du Preez, C., Finney, J., González-Irusta, J.M., Howell, K., Knudby, A.,
616 Lacharité, M., Lee, J., Murillo, F.J., Beazley, L., Roberts, J.M., Roberts, M., Rooper, C.,
617 Rowden, A., Rubidge, E., Stanley, R., Stirling, D., Tanaka, K.R., Vanhatalo, J., Weigel, B.,
618 Woolley, S., and Yesson, C. 2019. [Use of Species Distribution Modeling in the Deep Sea.](#)
619 Can. Tech. Rep. Fish. Aquat. Sci. 3296: ix + 76 p.
- 620 Kenchington, T.J., and Kenchington, E.L.R. 2017. [Biodiversity metrics for use in the](#)
621 [ecosystem approach to oceans management.](#) Can. Tech. Rep. Fish. Aquat. Sci. 3186:
622 iv + 49 p.
- 623 Lohr, S. 1999. Sampling: Design and analysis. Pacific Grove, CA: Brooks/Cole Publishing
624 Company.
- 625 Losier, R.J., and Waite, L.E. 1989. Systematic listing of scientific and/or common names of
626 invertebrates and marine plants and their respective codes used by marine fish division,
627 Fisheries and Oceans, Scotia-Fundy Region. Canadian Data Report of Fisheries and

- 628 Aquatic Sciences (721).
- 629 Lowen, B., Hart, D., Stanley, R., Lehnert, S., Bradbury, I., and C., D. 2019. [Assessing effects](#)
630 [of genetic, environmental, and biotic gradients in species distribution modelling](#). ICES
631 Journal of Marine Science 76(6): 1762–1775.
- 632 Myers, R.A., and Stokes, K. 1989. Density-dependent habitat utilization of groundfish and the
633 improvement of research surveys. (D:15). International Council for the Exploration of the
634 Sea Council Meeting.
- 635 Nelson J. S., Crossman E. J., Espinosa-Perez H., Findley L. T., Gilbert C. R., Lea R. N., and
636 D., W.J. 2004. Common and scientific names of fishes from the united states canada
637 and mexico. Number 29 in Special Publication. American Fisheries Society, Bethesda,
638 Maryland, sixth edition.
- 639 Neprin, J., Gregr, E.J., St. Germain, C., Fields, C., and Finney, J.L. and. 2019. [Development](#)
640 [of a Species Distribution Modelling Framework and its Application to Twelve Species on](#)
641 [Canada's Pacific Coast](#). DFO Can. Sci. Advis. Sec. Res. Doc. 2020/004. xii + 107 p.
- 642 Neuwirth, E. 2014. RColorBrewer:ColorBrewer palettes. R package version 1.1-2 <https://CRAN.R-project.org/package=RColorBrewer>.
- 643
- 644 Page L. M., Espinosa-Pérez H., Findley L. T., Gilbert C. R., Lea R. N., Mandrak N. E., and
645 M.R.L., and S., N.J. 2013. Common and scientific names of fishes from the United States,
646 Canada, and Mexico. American Fisheries Society.
- 647 Pebesma, E. 2004. Multivariable geostatistics in S: The gstat package. *In* Computers and
648 Geosciences.
- 649 Perry, R.I., and Smith, S.J. 1994. Identifying habitat associations of marine fishes using
650 survey data: An application to the northwest atlantic. Canadian Journal of Fisheries and
651 Aquatic Sciences (51(3)): 589–602.
- 652 R Core Team. 2020. R: A language and environment for statistical computing. R Foundation
653 for Statistical Computing, Vienna, Austria.
- 654 Ricard, D., and Gomez, C. 2021. Maritimes-SUMMER-atlas. <https://github.com/dfo-gulf-science/Maritimes-SUMMER-Atlas>; GitHub.
- 655
- 656 Ricard, D., and Shackell, N.L. 2013. [Population status \(abundance/biomass, geographic](#)
657 [extent, body size and condition\), important habitat, depth, temperature and salinity of](#)
658 [marine fish and invertebrates on the Scotian Shelf and Bay of Fundy \(1970-2012\)](#). Can.
659 Tech. Rep. Fish. Aquat. Sci. 3012: viii + 180 p.
- 660 Ripley, B., and Lapsley, M. 2019. RODBC: ODBC database access. R package version 1.3-16
661 <http://CRAN.R-project.org/package=RODBC>.
- 662 Ripley, B., Venables, B., Bates, D., Hornik, K., Gebhardt, A., and Firth, D. 2020. Modern
663 applied statistics with s. R package version 7.3-53 <https://cran.r-project.org/web/packages/MASS/index.html>.
- 664
- 665 Robinson, N.M., Nelson, W.A., Costello, M.J., Sutherland, J.E., and Lundquist, C.J.
666 2017. Systematic review of marine-based species distribution models (SDMs) with

- recommendations for best practice. *Front. Mar. Sci.* (4): 421.
- Schnute, J.T., Boers, N., and Haigh, R. 2019. PBSmapping: Mapping fisheries data and spatial analysis tools. R package version 2.72.1 <https://cran.r-project.org/web/packages/PBSmapping/index.html>.
- Shackell, N., Brickman, D., and Frank, K. 2013. Reserve site selection for data-poor invertebrate fisheries using patch scale and dispersal dynamics: A case study of sea cucumber. *Aquatic Conserv: Mar. Freshw. Ecosyst.* 23: 723–731.
- Simon, J.E., and Comeau, P.A. 1994. Summer distribution and abundance trends of species caught on the Scotian Shelf from 1970-92, by the research vessel groundfish survey. *Can. Tech. Rep. Fish. Aquat. Sci.* 1953.
- Smith, C.D., Serdynska, A.R., King, M.C., and Shackell, N.L. 2015. Spring, summer and fall distribution of common demersal fishes on the Scotian Shelf between 1978 and 1985. *Can. Tech. Rep. Fish. Aquat. Sci.* 3068: vi + 38 p.
- Smith, S.J. 1996. Assessment of groundfish stocks based on bottom trawl survey results. *NAFO Scientific Council Studies* 28: 25–53.
- Stanley, R.E., DiBacco, C., Lowen, B., Beiko, R., Jeffery, N., Wyngaarden, M., Bentzen, P., Brickman, D., Benestan, L., Bernatchez, L., Johnson, C., Snelgrove, P., Wang, Z., and Wringe, I., B. Bradbury. 2018. A climate-associated multispecies cryptic cline in the northwest Atlantic. *Science Advances* 4(3): 1–7.
- Swain, D.P., and Morin, R. 1996. Relationships between geographic distribution and abundance of American plaice (*Hippoglossoides platessoides*) in the southern Gulf of St. Lawrence. *Canadian Journal of Fisheries and Aquatic Sciences* 53(1): 106–119.
- Swain, D.P., and Sinclair, A.F. 1994. Fish distribution and catchability: What is the appropriate measure of distribution? *Canadian Journal of Fisheries and Aquatic Sciences* 51(5): 1046–1054.
- Tremblay M. J., B.R., Black G. A. P. 2007. The distribution of common decapod crustaceans and other invertebrates recorded in annual ecosystem surveys of the scotian shelf 1999-2006, by the research vessel groundfish survey. *Can. Tech. Rep. Fish. Aquat. Sci.* 74.
- W., G.B.J., Shackell, N.L., Ferguson, K., P., G., Cogswell, A., Brickman, D., Wang, Z., Cook, A., Brennan, C.E., and Saba, V.S. 2019. Climate change vulnerability of American lobster fishing communities in Atlantic Canada. *Frontiers in Marine Science* 6: 579.
- Ward-Paige, C.A., and Bundy, A. 2015. Mapping Biodiversity on the Scotian Shelf and in the Bay of Fundy. *Can. Tech. Rep. Fish. Aquat. Sci.* 3068: vi + 38 p.
- Wickham, H. 2019. Tidyverse: Easily install and load the 'tidyverse'. R package version 1.3.0 <https://cran.r-project.org/web/packages/tidyverse/index.html>.
- Zisserson, B.M., Cameron, B.J., Glass, A.C., and Choi, J.S. 2019. Assessment of Scotian Shelf Snow Crab in 2017. DFO Can. Sci. Advis. Sec. Res. Doc. 2018/051. ix + 147 p.

7 Appendix

705

7.1 Atlantic cod (*Morue franche*) - species code 10 (category LF)

706

Scientific name: [Gadus morhua](#)

707




Figure 7.1A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic cod.

708




Figure 7.1B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic cod.




Figure 7.1C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic cod.






Figure 7.1D. Average fish condition in NAFO units 4X and 4VW for Atlantic cod.



711



Freq	Depth	Temp	Sal
F5	26	1.2	31.00
F25	43	3.5	32.47
F50	60	6.0	33.27
F75	82	8.1	34.40
F95	126	10.0	35.03

Figure 7.1E. Catch distribution by depth, temperature and salinity of Atlantic cod.

712




Figure 7.1F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic cod.

713

7.2 Haddock (Aiglefin) - species code 11 (category LF)

714

Scientific name: [Melanogrammus aeglefinus](#)

715




Figure 7.2A. Inverse distance weighted distribution of catch biomass (kg/tow) for Haddock.

716




Figure 7.2B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Haddock.




Figure 7.2C. Length frequency distribution in NAFO units 4X and 4VW for Haddock.






Figure 7.2D. Average fish condition in NAFO units 4X and 4VW for Haddock.



719



Freq	Depth	Temp	Sal
F5	27	1.1	31.00
F25	49	3.0	32.45
F50	66	5.5	33.14
F75	87	7.9	34.36
F95	127	10.0	35.03

Figure 7.2E. Catch distribution by depth, temperature and salinity of Haddock.




Figure 7.2F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Haddock.

721

7.3 White hake (Merluche blanche) - species code 12 (category LF)

722

Scientific name: [Urophycis tenuis](#)

723




Figure 7.3A. Inverse distance weighted distribution of catch biomass (kg/tow) for White hake.

724




Figure 7.3B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of White hake.




Figure 7.3C. Length frequency distribution in NAFO units 4X and 4VW for White hake.






Figure 7.3D. Average fish condition in NAFO units 4X and 4VW for White hake.




727



Freq	Depth	Temp	Sal
F5	40	1.1	31.00
F25	70	3.0	32.46
F50	89	5.5	33.20
F75	115	7.9	34.39
F95	163	10.0	35.04

Figure 7.3E. Catch distribution by depth, temperature and salinity of White hake.



728

Figure 7.3F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for White hake.

729

7.4 Red hake (Merluche écureuil) - species code 13 (category LF)

730

Scientific name: [Urophycis chuss](#)

731




Figure 7.4A. Inverse distance weighted distribution of catch biomass (kg/tow) for Red hake.

732




Figure 7.4B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Red hake.




Figure 7.4C. Length frequency distribution in NAFO units 4X and 4VW for Red hake.






Figure 7.4D. Average fish condition in NAFO units 4X and 4VW for Red hake.



735



Freq	Depth	Temp	Sal
F5	35	1.1	31.00
F25	62	2.9	32.43
F50	82	5.4	33.12
F75	103	7.9	34.32
F95	130	10.0	35.08

Figure 7.4E. Catch distribution by depth, temperature and salinity of Red hake.

736




Figure 7.4F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Red hake.

737

7.5 Silver hake (*Merlu argenté*) - species code 14 (category LF)

738

Scientific name: [Merluccius bilinearis](#)

739




Figure 7.5A. Inverse distance weighted distribution of catch biomass (kg/tow) for Silver hake.

740




Figure 7.5B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Silver hake.




Figure 7.5C. Length frequency distribution in NAFO units 4X and 4VW for Silver hake.






Figure 7.5D. Average fish condition in NAFO units 4X and 4VW for Silver hake.




743



Freq	Depth	Temp	Sal
F5	28	1.2	31.00
F25	55	3.1	32.46
F50	77	5.5	33.20
F75	104	8.0	34.37
F95	137	10.0	35.07

Figure 7.5E. Catch distribution by depth, temperature and salinity of Silver hake.



744

Figure 7.5F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Silver hake.

745

7.6 Pollock (Goberge) - species code 16 (category LF)

746

Scientific name: [Pollachius virens](#)

747




Figure 7.6A. Inverse distance weighted distribution of catch biomass (kg/tow) for Pollock.

748




Figure 7.6B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Pollock.




Figure 7.6C. Length frequency distribution in NAFO units 4X and 4VW for Pollock.






Figure 7.6D. Average fish condition in NAFO units 4X and 4VW for Pollock.



751



Freq	Depth	Temp	Sal
F5	37	1.1	31.00
F25	60	2.8	32.45
F50	82	5.3	33.14
F75	108	7.7	34.33
F95	137	10.0	35.03

Figure 7.6E. Catch distribution by depth, temperature and salinity of Pollock.

752




Figure 7.6F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Pollock.

753

7.7 Atlantic redfishes (Sébastes de l'Atlantique) - species code 23 (category LF)

754

Scientific name: [Sebastes](#)

755




Figure 7.7A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic redfishes.

756




Figure 7.7B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic redfishes.




Figure 7.7C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic redfishes.






Figure 7.7D. Average fish condition in NAFO units 4X and 4VW for Atlantic redfishes.



759



Freq	Depth	Temp	Sal
F5	47	1.2	31.00
F25	68	3.4	32.48
F50	86	5.9	33.29
F75	114	8.1	34.41
F95	166	10.0	35.05

Figure 7.7E. Catch distribution by depth, temperature and salinity of Atlantic redfishes.




Figure 7.7F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic redfishes.

761

7.8 Atlantic halibut (Flétan de l'Atlantique) - species code 30 (category LF)

762

Scientific name: [Hippoglossus hippoglossus](#)

763




Figure 7.8A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic halibut.

764




Figure 7.8B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic halibut.




Figure 7.8C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic halibut.






Figure 7.8D. Average fish condition in NAFO units 4X and 4VW for Atlantic halibut.



767



Freq	Depth	Temp	Sal
F5	31	1.3	31.00
F25	49	3.2	32.45
F50	60	5.8	33.16
F75	75	8.3	34.34
F95	122	10.0	35.08

Figure 7.8E. Catch distribution by depth, temperature and salinity of Atlantic halibut.

768




Figure 7.8F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic halibut.

769

7.9 American plaice (*Ple canadienne*) - species code 40 (category LF)

770

Scientific name: [Hippoglossoides platessoides](#)

771




Figure 7.9A. Inverse distance weighted distribution of catch biomass (kg/tow) for American plaice.

772




Figure 7.9B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of American plaice.




Figure 7.9C. Length frequency distribution in NAFO units 4X and 4VW for American plaice.






Figure 7.9D. Average fish condition in NAFO units 4X and 4VW for American plaice.



775



Freq	Depth	Temp	Sal
F5	29	1.3	31.00
F25	46	3.7	32.48
F50	67	6.1	33.27
F75	87	8.1	34.41
F95	133	10.0	35.05

Figure 7.9E. Catch distribution by depth, temperature and salinity of American plaice.




Figure 7.9F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for American plaice.

777

7.10 Witch flounder (*Ple grise*) - species code 41 (category LF)

778

Scientific name: [Glyptocephalus cynoglossus](#)

779




Figure 7.10A. Inverse distance weighted distribution of catch biomass (kg/tow) for Witch flounder.

780




Figure 7.10B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Witch flounder.




Figure 7.10C. Length frequency distribution in NAFO units 4X and 4VW for Witch flounder.






Figure 7.10D. Average fish condition in NAFO units 4X and 4VW for Witch flounder.




783



Freq	Depth	Temp	Sal
F5	32	1.3	31.00
F25	57	3.5	32.49
F50	77	6.1	33.30
F75	102	8.2	34.45
F95	152	10.0	35.06

Figure 7.10E. Catch distribution by depth, temperature and salinity of Witch flounder.



784

Figure 7.10F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Witch flounder.

785

7.11 Yellowtail flounder (Limande à queue jaune) - species code 42 (category LF)

786

Scientific name: [Limanda ferruginea](#)

787




Figure 7.11A. Inverse distance weighted distribution of catch biomass (kg/tow) for Yellowtail flounder.

788




Figure 7.11B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Yellowtail flounder.




Figure 7.11C. Length frequency distribution in NAFO units 4X and 4VW for Yellowtail flounder.






Figure 7.11D. Average fish condition in NAFO units 4X and 4VW for Yellowtail flounder.



791



Freq	Depth	Temp	Sal
F5	22	1.4	31.00
F25	32	3.7	32.52
F50	43	6.3	33.25
F75	56	8.2	34.41
F95	81	10.0	35.06

Figure 7.11E. Catch distribution by depth, temperature and salinity of Yellowtail flounder.

792




Figure 7.11F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Yellowtail flounder.

793

7.12 Winter flounder (Limande-plie rouge) - species code 43 (category LF)

794

Scientific name: [Pseudopleuronectes americanus](#)

795




Figure 7.12A. Inverse distance weighted distribution of catch biomass (kg/tow) for Winter flounder.

796




Figure 7.12B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Winter flounder.




Figure 7.12C. Length frequency distribution in NAFO units 4X and 4VW for Winter flounder.






Figure 7.12D. Average fish condition in NAFO units 4X and 4VW for Winter flounder.



799



Freq	Depth	Temp	Sal
F5	22	1.1	31.00
F25	31	3.0	32.48
F50	42	5.5	33.17
F75	54	8.3	34.47
F95	84	10.0	35.10

Figure 7.12E. Catch distribution by depth, temperature and salinity of Winter flounder.




Figure 7.12F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Winter flounder.

801 **7.13 Atlantic wolffish (*Loup atlantique*) - species code 50 (category LF)**

802 Scientific name: [Anarhichas lupus](#)

803




Figure 7.13A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic wolffish.

804




Figure 7.13B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic wolffish.




Figure 7.13C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic wolffish.






Figure 7.13D. Average fish condition in NAFO units 4X and 4VW for Atlantic wolffish.



Freq	Depth	Temp	Sal
F5	34	1.4	31.00
F25	49	3.6	32.48
F50	62	6.0	33.25
F75	77	8.1	34.33
F95	112	10.0	35.05

Figure 7.13E. Catch distribution by depth, temperature and salinity of Atlantic wolffish.




Figure 7.13F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic wolffish.

809

7.14 Atlantic herring (Hareng de l'Atlantique) - species code 60 (category LF)

810

Scientific name: [Clupea harengus](#)

811




Figure 7.14A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic herring.

812




Figure 7.14B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic herring.




Figure 7.14C. Length frequency distribution in NAFO units 4X and 4VW for Atlantic herring.






Figure 7.14D. Average fish condition in NAFO units 4X and 4VW for Atlantic herring.



815



Freq	Depth	Temp	Sal
F5	32	1.2	31.00
F25	54	3.2	32.45
F50	74	5.6	33.22
F75	95	8.3	34.38
F95	132	10.0	35.10

Figure 7.14E. Catch distribution by depth, temperature and salinity of Atlantic herring.

816




Figure 7.14F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Atlantic herring.

817 **7.15 Longhorn sculpin (Chaboisseau à dix-huit épines) - species code 300 (category**
 818 **LF)**

819 Scientific name: [Myoxocephalus octodecemspiniferus](#)

820




Figure 7.15A. Inverse distance weighted distribution of catch biomass (kg/tow) for Longhorn sculpin.

821




Figure 7.15B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Longhorn sculpin.




Figure 7.15C. Length frequency distribution in NAFO units 4X and 4VW for Longhorn sculpin.







Figure 7.15D. Average fish condition in NAFO units 4X and 4VW for Longhorn sculpin.




824



Freq	Depth	Temp	Sal
F5	23	1.2	31.00
F25	33	3.3	32.51
F50	48	5.8	33.29
F75	64	8.1	34.38
F95	96	10.0	35.05

Figure 7.15E. Catch distribution by depth, temperature and salinity of Longhorn sculpin.



825

Figure 7.15F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Longhorn sculpin.

826

7.16 Moustache sculpin (Faux-trigle armé) - species code 304 (category LF)

827

Scientific name: [Triglops murrayi](#)

828




Figure 7.16A. Inverse distance weighted distribution of catch biomass (kg/tow) for Moustache sculpin.

829




Figure 7.16B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Moustache sculpin.




Figure 7.16C. Length frequency distribution in NAFO units 4X and 4VW for Moustache sculpin.






Figure 7.16D. Average fish condition in NAFO units 4X and 4VW for Moustache sculpin.



832



Freq	Depth	Temp	Sal
F5	30	1.5	31.00
F25	40	4.0	32.48
F50	50	6.5	33.31
F75	63	8.2	34.39
F95	88	10.0	35.06

Figure 7.16E. Catch distribution by depth, temperature and salinity of Moustache sculpin.

833




Figure 7.16F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Moustache sculpin.

834

7.17 Sea raven (Hémithriptère atlantique) - species code 320 (category LF)

835

Scientific name: [Hemitripterus americanus](#)

836




Figure 7.17A. Inverse distance weighted distribution of catch biomass (kg/tow) for Sea raven.

837




Figure 7.17B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Sea raven.




Figure 7.17C. Length frequency distribution in NAFO units 4X and 4VW for Sea raven.





Figure 7.17D. Average fish condition in NAFO units 4X and 4VW for Sea raven.



Freq	Depth	Temp	Sal
F5	22	1.1	31.00
F25	32	3.1	32.43
F50	45	5.4	33.15
F75	61	8.0	34.35
F95	89	10.0	35.05

Figure 7.17E. Catch distribution by depth, temperature and salinity of Sea raven.




Figure 7.17F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Sea raven.

842

7.18 Alligatorfish (Poisson-alligator atlantique) - species code 340 (category LF)

843

Scientific name: [Aspidophoroides monopterygius](#)

844




Figure 7.18A. Inverse distance weighted distribution of catch biomass (kg/tow) for Alligatorfish.

845




Figure 7.18B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Alligatorfish.




Figure 7.18C. Length frequency distribution in NAFO units 4X and 4VW for Alligatorfish.






Figure 7.18D. Average fish condition in NAFO units 4X and 4VW for Alligatorfish.



848



Freq	Depth	Temp	Sal
F5	32	1.5	31.00
F25	44	3.7	32.53
F50	57	6.1	33.28
F75	72	8.2	34.45
F95	102	10.0	35.10

Figure 7.18E. Catch distribution by depth, temperature and salinity of Alligatorfish.

849




Figure 7.18F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Alligatorfish.

850 **7.19 Monkfish (Baudroie d'Amérique) - species code 400 (category LF)**

851 Scientific name: [Lophius americanus](#)




Figure 7.19A. Inverse distance weighted distribution of catch biomass (kg/tow) for Monkfish.




Figure 7.19B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Monkfish.




Figure 7.19C. Length frequency distribution in NAFO units 4X and 4VW for Monkfish.






Figure 7.19D. Average fish condition in NAFO units 4X and 4VW for Monkfish.



856



Freq	Depth	Temp	Sal
F5	33	1.1	31.00
F25	60	2.9	32.43
F50	84	5.3	33.07
F75	110	7.8	34.31
F95	148	10.0	35.03

Figure 7.19E. Catch distribution by depth, temperature and salinity of Monkfish.




Figure 7.19F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Monkfish.

858

7.20 Ocean pout (Loquette d'Amérique) - species code 640 (category LF)

859

Scientific name: [Zoarces americanus](#)

860




Figure 7.20A. Inverse distance weighted distribution of catch biomass (kg/tow) for Ocean pout.

861




Figure 7.20B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Ocean pout.




Figure 7.20C. Length frequency distribution in NAFO units 4X and 4VW for Ocean pout.







Figure 7.20D. Average fish condition in NAFO units 4X and 4VW for Ocean pout.




864



Freq	Depth	Temp	Sal
F5	31	1.0	31.00
F25	52	2.8	32.46
F50	69	5.1	33.22
F75	85	7.7	34.34
F95	116	10.0	35.03

Figure 7.20E. Catch distribution by depth, temperature and salinity of Ocean pout.



865

Figure 7.20F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Ocean pout.

866 **7.21 Thorny skate (Raie épineuse) - species code 201 (category LF)**

867 Scientific name: [Amblyraja radiata](#)

868




Figure 7.21A. Inverse distance weighted distribution of catch biomass (kg/tow) for Thorny skate.

869




Figure 7.21B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Thorny skate.




Figure 7.21C. Length frequency distribution in NAFO units 4X and 4VW for Thorny skate.






Figure 7.21D. Average fish condition in NAFO units 4X and 4VW for Thorny skate.




872



Freq	Depth	Temp	Sal
F5	29	1.3	31.00
F25	45	3.7	32.50
F50	67	6.2	33.30
F75	91	8.1	34.40
F95	148	9.9	35.03

Figure 7.21E. Catch distribution by depth, temperature and salinity of Thorny skate.



873

Figure 7.21F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Thorny skate.

874

7.22 Smooth skate (Raie lisse) - species code 202 (category LF)

875

Scientific name: [Malacoraja senta](#)

876




Figure 7.22A. Inverse distance weighted distribution of catch biomass (kg/tow) for Smooth skate.

877




Figure 7.22B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Smooth skate.




Figure 7.22C. Length frequency distribution in NAFO units 4X and 4VW for Smooth skate.




Figure 7.22D. Average fish condition in NAFO units 4X and 4VW for Smooth skate.




Figure 7.22E. Catch distribution by depth, temperature and salinity of Smooth skate.




Figure 7.22F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Smooth skate.

882

7.23 Winter skate (Raie tachetée) - species code 204 (category LF)

883

Scientific name: [Leucoraja ocellata](#)

884




Figure 7.23A. Inverse distance weighted distribution of catch biomass (kg/tow) for Winter skate.

885




Figure 7.23B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Winter skate.




Figure 7.23C. Length frequency distribution in NAFO units 4X and 4VW for Winter skate.






Figure 7.23D. Average fish condition in NAFO units 4X and 4VW for Winter skate.




888



Freq	Depth	Temp	Sal
F5	21	1.1	31.00
F25	32	3.5	32.44
F50	45	5.9	33.19
F75	71	8.3	34.42
F95	140	10.0	35.03

Figure 7.23E. Catch distribution by depth, temperature and salinity of Winter skate.



889

Figure 7.23F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Winter skate.

890 **7.24 Picked dogfish (Aiguillat commun) - species code 220 (category LF)**

891 Scientific name: [Squalus acanthias](#)




Figure 7.24A. Inverse distance weighted distribution of catch biomass (kg/tow) for Picked dogfish.




Figure 7.24B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Picked dogfish.




Figure 7.24C. Length frequency distribution in NAFO units 4X and 4VW for Picked dogfish.






Figure 7.24D. Average fish condition in NAFO units 4X and 4VW for Picked dogfish.



896



Freq	Depth	Temp	Sal
F5	35	1.1	31.00
F25	60	2.8	32.47
F50	83	5.2	33.28
F75	108	7.7	34.37
F95	139	10.0	35.02

Figure 7.24E. Catch distribution by depth, temperature and salinity of Picked dogfish.




Figure 7.24F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Picked dogfish.

898 **7.25 Northern shortfin squid (*Encornet rouge nordique*) - species code 4511 (category**
899 **LF)**

900 Scientific name: [Illex illecebrosus](#)

901




Figure 7.25A. Inverse distance weighted distribution of catch biomass (kg/tow) for Northern shortfin squid.

902




Figure 7.25B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Northern shortfin squid.




Figure 7.25C. Length frequency distribution in NAFO units 4X and 4VW for Northern shortfin squid.






Figure 7.25D. Average fish condition in NAFO units 4X and 4VW for Northern shortfin squid.



905



Freq	Depth	Temp	Sal
F5	31	1.3	31.00
F25	53	3.5	32.48
F50	71	5.9	33.28
F75	93	8.1	34.39
F95	139	10.0	35.05

Figure 7.25E. Catch distribution by depth, temperature and salinity of Northern shortfin squid.

906






Figure 7.25F. DDHS slopes versus median stratum abundance. The last two digits of each stratum number is shown in the figure for Northern shortfin squid.

907 **7.26 Atlantic hagfish (*Myxine du nord*) - species code 241 (category LI)**

908 Scientific name: [Myxine glutinosa](#)



909 Figure 7.26A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic hagfish.



910 Figure 7.26B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic hagfish.

911

7.27 Cusk (Brosme) - species code 15 (category LI)

912

Scientific name: [Brosme brosme](#)

913




Figure 7.27A. Inverse distance weighted distribution of catch biomass (kg/tow) for Cusk.

914




Figure 7.27B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Cusk.

915

7.28 Greenland halibut (Flétan noir) - species code 31 (category LI)

916

Scientific name: [Reinhardtius hippoglossoides](#)

917




Figure 7.28A. Inverse distance weighted distribution of catch biomass (kg/tow) for Greenland halibut.

918




Figure 7.28B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Greenland halibut.

919 **7.29 Gulf Stream flounder (Plie du Gulf Stream) - species code 44 (category LI)**

920 Scientific name: [Citharichthys arctifrons](#)




Figure 7.29A. Inverse distance weighted distribution of catch biomass (kg/tow) for Gulf Stream flounder.




Figure 7.29B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Gulf Stream flounder.

923

7.30 American shad (*Alose savoureuse*) - species code 61 (category LI)

924

Scientific name: [Alosa sapidissima](#)

925




Figure 7.30A. Inverse distance weighted distribution of catch biomass (kg/tow) for American shad.

926




Figure 7.30B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of American shad.

927

7.31 Alewife (Gaspareau) - species code 62 (category LI)

928

Scientific name: *Alosa pseudoharengus*

929




Figure 7.31A. Inverse distance weighted distribution of catch biomass (kg/tow) for Alewife.

930




Figure 7.31B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Alewife.

931

7.32 Capelin (Capelan) - species code 64 (category LI)

932

Scientific name: [Mallotus villosus](#)

933




Figure 7.32A. Inverse distance weighted distribution of catch biomass (kg/tow) for Capelin.

934




Figure 7.32B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Capelin.

935 **7.33 Atlantic mackerel (*Maquereau commun*) - species code 70 (category LI)**

936 Scientific name: *Scomber scombrus*




Figure 7.33A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic mackerel.




Figure 7.33B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic mackerel.

939

7.34 Longfin hake (Merluche à longues nageoires) - species code 112 (category LI)

940

Scientific name: [Phycis chesteri](#)

941




Figure 7.34A. Inverse distance weighted distribution of catch biomass (kg/tow) for Longfin hake.

942




Figure 7.34B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Longfin hake.

943

7.35 Fourbeard rockling (Motelle à quatre barbillons) - species code 114 (category LI)

944

Scientific name: [Enchelyopus cimbrius](#)

945




Figure 7.35A. Inverse distance weighted distribution of catch biomass (kg/tow) for Fourbeard rockling.

946




Figure 7.35B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Fourbeard rockling.

947

7.36 Blackbelly rosefish (Sébaste chèvre) - species code 123 (category LI)

948

Scientific name: [Helicolenus dactylopterus](#)

949




Figure 7.36A. Inverse distance weighted distribution of catch biomass (kg/tow) for Blackbelly rosefish.

950




Figure 7.36B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Blackbelly rosefish.

951

7.37 Greater argentine (Grande argentine) - species code 160 (category LI)

952

Scientific name: [Argentina silus](#)

953




Figure 7.37A. Inverse distance weighted distribution of catch biomass (kg/tow) for Greater argentine.

954




Figure 7.37B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Greater argentine.

955 **7.38 Arctic hookear sculpin (*Hameçon neigeux*) - species code 306 (category LI)**

956 Scientific name: [Artediellus uncinatus](#)

957




Figure 7.38A. Inverse distance weighted distribution of catch biomass (kg/tow) for Arctic hookear sculpin.

958




Figure 7.38B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Arctic hookear sculpin.

959 **7.39 Atlantic poacher (*Agone atlantique*) - species code 350 (category LI)**

960 Scientific name: [Leptagonus decagonus](#)




Figure 7.39A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic poacher.




Figure 7.39B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic poacher.

963 **7.40 Marlin-spike grenadier (Grenadier du Grand Banc) - species code 410 (category**
 964 **LI)**

965 Scientific name: [Nezumia bairdii](#)




Figure 7.40A. Inverse distance weighted distribution of catch biomass (kg/tow) for Marlin-spike grenadier.




Figure 7.40B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Marlin-spike grenadier.


968

7.41 Lumpfish (Lompe) - species code 501 (category LI)

969

Scientific name: [Cyclopterus lumpus](#)

970



971




Figure 7.41B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Lumpfish.

972 **7.42 Atlantic spiny lumpsucker (Petite poule de mer atlantique) - species code 502**
 973 (**category LI**)

974 Scientific name: [Eumicrotremus spinosus](#)




Figure 7.42A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic spiny lumpsucker.




Figure 7.42B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic spiny lumpsucker.

977

7.43 Sand lance (Lançon) - species code 610 (category LI)

978

Scientific name: [Ammodytes dubius](#)

979




Figure 7.43A. Inverse distance weighted distribution of catch biomass (kg/tow) for Sand lance.

980




Figure 7.43B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Sand lance.

981

7.44 Snakeblenny (Lompénie-serpent) - species code 622 (category LI)

982

Scientific name: [Lumpenus lampretaeformis](#)

983




Figure 7.44A. Inverse distance weighted distribution of catch biomass (kg/tow) for Snakeblenny.

984




Figure 7.44B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Snakeblenny.

985

7.45 Daubed shanny (Lompénie tachetée) - species code 623 (category LI)

986

Scientific name: [Leptoclinus maculatus](#)

987




Figure 7.45A. Inverse distance weighted distribution of catch biomass (kg/tow) for Daubed shanny.

988




Figure 7.45B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Daubed shanny.

989

7.46 Vahl's eelpout (*Lycodes vahlii*) - species code 647 (category LI)

990

Scientific name: [Lycodes vahlii](#)

991




Figure 7.46A. Inverse distance weighted distribution of catch biomass (kg/tow) for Vahl's eelpout.

992




Figure 7.46B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Vahl's eelpout.

993

7.47 Atlantic butterfish (*Stromaté fossette*) - species code 701 (category LI)

994

Scientific name: [Peprilus triacanthus](#)

995




Figure 7.47A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic butterfish.

996




Figure 7.47B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic butterfish.

997 **7.48 Atlantic hookear sculpin (*Hameçon atlantique*) - species code 880 (category LI)**

998 Scientific name: [Artediellus atlanticus](#)

999




Figure 7.48A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic hookear sculpin.




Figure 7.48B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic hookear sculpin.

1001

7.49 Barndoor skate (Grande raie) - species code 200 (category LI)

1002

Scientific name: [Dipturus laevis](#)

1003




Figure 7.49A. Inverse distance weighted distribution of catch biomass (kg/tow) for Barndoor skate.

1004




Figure 7.49B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Barndoor skate.

1005

7.50 Little skate (Raie hérisson) - species code 203 (category LI)

1006

Scientific name: [Leucoraja erinacea](#)

1007




Figure 7.50A. Inverse distance weighted distribution of catch biomass (kg/tow) for Little skate.

1008




Figure 7.50B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Little skate.

1009

7.51 Northern prawn (Crevette nordique) - species code 2211 (category SF)

1010

Scientific name: [Pandalus borealis](#)

1011




Figure 7.51A. Inverse distance weighted distribution of catch biomass (kg/tow) for Northern prawn.

1012




Figure 7.51B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Northern prawn.

1013

7.52 Jonah crab (*Tourteau jona*) - species code 2511 (category SF)

1014

Scientific name: [Cancer borealis](#)

1015




Figure 7.52A. Inverse distance weighted distribution of catch biomass (kg/tow) for Jonah crab.

1016




Figure 7.52B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Jonah crab.

1017

7.53 Atlantic rock crab (Tourteau poïnclos) - species code 2513 (category SF)

1018

Scientific name: [Cancer irroratus](#)

1019




Figure 7.53A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic rock crab.

1020




Figure 7.53B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic rock crab.

1021

7.54 Arctic lyre crab (*Crabe Hyas coarctatus*) - species code 2521 (category SF)

1022

Scientific name: [Hyas coarctatus](#)

1023




Figure 7.54A. Inverse distance weighted distribution of catch biomass (kg/tow) for Arctic lyre crab.

1024




Figure 7.54B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Arctic lyre crab.

1025 **7.55 Atlantic king crab (Crabe épineux du nord) - species code 2523 (category SF)**

1026 Scientific name: [Lithodes maja](#)




Figure 7.55A. Inverse distance weighted distribution of catch biomass (kg/tow) for Atlantic king crab.




Figure 7.55B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Atlantic king crab.

1029

7.56 Queen crab (Crabe des neiges) - species code 2526 (category SF)

1030

Scientific name: [Chionoecetes opilio](#)

1031




Figure 7.56A. Inverse distance weighted distribution of catch biomass (kg/tow) for Queen crab.

1032




Figure 7.56B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Queen crab.

1033

7.57 Great spider crab (Crabe lyre araignée) - species code 2527 (category SF)

1034

Scientific name: [Hyas araneus](#)

1035




Figure 7.57A. Inverse distance weighted distribution of catch biomass (kg/tow) for Great spider crab.

1036




Figure 7.57B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of Great spider crab.

1037

7.58 American lobster (Homard américain) - species code 2550 (category SF)

1038

Scientific name: [Homarus americanus](#)

1039




Figure 7.58A. Inverse distance weighted distribution of catch biomass (kg/tow) for American lobster.

1040




Figure 7.58B. Stratified random estimates of biomass (kg/tow), D75 and D95 and the correlation between D75 and biomass of American lobster.

1041

7.59 Sea lamprey (*Lamproie marine*) - species code 240 (category LR)

1042

Scientific name: [Petromyzon marinus](#)

1043




Figure 7.59A. Catch distribution for Sea lamprey.

1044

7.60 Atlantic tomcod (*Poulamon atlantique*) - species code 17 (category LR)

1045

Scientific name: [Micogadus tomcod](#)

1046




Figure 7.60A. Catch distribution for Atlantic tomcod.

1047

7.61 Offshore silver hake (Merlu argenté du large) - species code 19 (category LR)

1048

Scientific name: [Merluccius albidus](#)

1049




Figure 7.61A. Catch distribution for Offshore silver hake.

1050

7.62 Spotted wolffish (Loup tacheté) - species code 51 (category LR)

1051

Scientific name: [Anarhichas minor](#)

1052




Figure 7.62A. Catch distribution for Spotted wolffish.

1053

7.63 Northern wolffish (Loup à tête large) - species code 52 (category LR)

1054

Scientific name: [Anarhichas denticulatus](#)

1055




Figure 7.63A. Catch distribution for Northern wolffish.

1056

7.64 Rainbow smelt (Éperlan arc-en-ciel) - species code 63 (category LR)

1057

Scientific name: [Osmerus mordax](#)

1058




Figure 7.64A. Catch distribution for Rainbow smelt.

1059

7.65 Cunner (Tanche-tautogue) - species code 122 (category LR)

1060

Scientific name: [Tautogolabrus adspersus](#)

1061




Figure 7.65A. Catch distribution for Cunner.

1062

7.66 Fourspot flounder (Cardeau à quatre ocelles) - species code 142 (category LR)

1063

Scientific name: [Hippoglossina oblonga](#)

1064




Figure 7.66A. Catch distribution for Fourspot flounder.

1065

7.67 Windowpane flounder (Turbot de sable) - species code 143 (category LR)

1066

Scientific name: [Scophthalmus aquosus](#)

1067




Figure 7.67A. Catch distribution for Windowpane flounder.

1068

7.68 Longnose greeneye (Oeil-vert à long nez) - species code 149 (category LR)

1069

Scientific name: [Parasudis triculenta](#)

1070




Figure 7.68A. Catch distribution for Longnose greeneye.

1071

7.69 Lanternfishes (Poissons-lanternes) - species code 150 (category LR)

1072

Scientific name: [Myctophidae](#)

1073




Figure 7.69A. Catch distribution for Lanternfishes.

1074

7.70 Shortnose greeneye (Éperlan du large) - species code 156 (category LR)

1075

Scientific name: [Chlorophthalmus agassizi](#)

1076




Figure 7.70A. Catch distribution for Shortnose greeneye.

1077

7.71 Silvery lightfish (Brossé améthyste) - species code 158 (category LR)

1078

Scientific name: [Maurolicus muelleri](#)

1079




Figure 7.71A. Catch distribution for Silvery lightfish.

1080

7.72 Boa dragonfish (Dragon-boa) - species code 159 (category LR)

1081

Scientific name: [Stomias boa](#)

1082




Figure 7.72A. Catch distribution for Boa dragonfish.

1083 **7.73 Shorthorn sculpin (Chabosseau à épines courtes) - species code 301 (category**
1084 **LR)**

1085 Scientific name: [Myoxocephalus scorpius](#)




Figure 7.73A. Catch distribution for Shorthorn sculpin.

1087

7.74 Grubby (Chabosseau bronzé) - species code 303 (category LR)

1088

Scientific name: [Myoxocephalus aenaeus](#)

1089




Figure 7.74A. Catch distribution for Grubby.

1090

7.75 Polar sculpin (Cotte polaire) - species code 307 (category LR)

1091

Scientific name: [Cottunculus microps](#)

1092




Figure 7.75A. Catch distribution for Polar sculpin.

1093

7.76 Spatulate sculpin (Icèle spatulée) - species code 314 (category LR)

1094

Scientific name: *Icelus spatula*

1095




Figure 7.76A. Catch distribution for Spatulate sculpin.

1096

7.77 Arctic alligatorfish (Poisson-alligator arctique) - species code 341 (category LR)

1097

Scientific name: [Ulcina olrikii](#)

1098




Figure 7.77A. Catch distribution for Arctic alligatorfish.


1099

7.78 Alligatorfishes (Poissons-alligator) - species code 351 (category LR)

1100

Scientific name: [Agonidae](#)

1101



1102

7.79 Roughnose grenadier (Grenadier-scie) - species code 412 (category LR)

1103

Scientific name: [Trachyrincus murrayi](#)

1104




Figure 7.79A. Catch distribution for Roughnose grenadier.

1105 **7.80 Roundnose grenadier (Grenadier de roche) - species code 414 (category LR)**

1106 Scientific name: [Coryphaenoides rupestris](#)




Figure 7.80A. Catch distribution for Roundnose grenadier.


1108

7.81 Atlantic seasnail (*Limace atlantique*) - species code 503 (category LR)

1109

Scientific name: [Liparis atlanticus](#)

1110



1111

Figure 7.81A. Catch distribution for Atlantic seasnail.

1111 7.82 Gelatinous snailfish (Limace gélatineuse) - species code 505 (category LR)

1112 Scientific name: [Liparis fabricii](#)




Figure 7.82A. Catch distribution for Gelatinous snailfish.

1114

7.83 Variegated snailfish (*Limace marbée*) - species code 512 (category LR)

1115

Scientific name: [Liparis gibbus](#)

1116




Figure 7.83A. Catch distribution for Variegated snailfish.

1117 **7.84 Sea tadpole (Petite limace de mer) - species code 520 (category LR)**

1118 Scientific name: [Careproctus reinhardtii](#)




Figure 7.84A. Catch distribution for Sea tadpole.

1120 7.85 Wolf eelpout (*Lycodes à tête longue*) - species code 603 (category LR)

1121 Scientific name: *Lycenchelys verrillii*




Figure 7.85A. Catch distribution for Wolf eelpout.

1123

7.86 Slender snipe eel (*Avocette ruban*) - species code 604 (category LR)

1124

Scientific name: [Nemichthys scolopaceus](#)

1125




Figure 7.86A. Catch distribution for Slender snipe eel.

1126

7.87 Newfoundland eelpout (*Lycodes* du Labrador) - species code 619 (category LR)

1127

Scientific name: [Lycodes terraenovae](#)

1128




Figure 7.87A. Catch distribution for Newfoundland eelpout.

1129 7.88 Newfoundland eelpout (*Lycodes* du Labrador) - species code 620 (category LR)

1130 Scientific name: [Lycodes lavalaei](#)




Figure 7.88A. Catch distribution for Newfoundland eelpout.

1132

7.89 Rock gunnel (Sigouine de roche) - species code 621 (category LR)

1133

Scientific name: [Pholis gunnellus](#)

1134




Figure 7.89A. Catch distribution for Rock gunnel.

1135

7.90 Radiated shanny (*Ulvaria deux-lignes*) - species code 625 (category LR)

1136

Scientific name: [Ulvaria subbifurcata](#)

1137




Figure 7.90A. Catch distribution for Radiated shanny.

1138

7.91 Fourline snakeblenny (Quatre-lignes atlantique) - species code 626 (category LR)

1139

Scientific name: [Eumesogrammus praecisus](#)

1140




Figure 7.91A. Catch distribution for Fourline snakeblenny.

1141

7.92 Wrymouth (Terrassier tacheté) - species code 630 (category LR)

1142

Scientific name: [Cryptacanthodes maculatus](#)

1143




Figure 7.92A. Catch distribution for Wrymouth.

1144

7.93 Spotfin dragonet (Dragonnet tacheté) - species code 637 (category LR)

1145

Scientific name: [Foetorepus agassizii](#)

1146




Figure 7.93A. Catch distribution for Spotfin dragonet.

1147 7.94 Arctic eelpout (*Lycodes arctique*) - species code 641 (category LR)

1148 Scientific name: [Lycodes reticulatus](#)




Figure 7.94A. Catch distribution for Arctic eelpout.

1150 **7.95 Atlantic soft pout (*Molasse atlantique*) - species code 646 (category LR)**

1151 Scientific name: [Melanostigma atlanticum](#)




Figure 7.95A. Catch distribution for Atlantic soft pout.

1153

7.96 Silvery John dory (Saint Pierre argenté) - species code 704 (category LR)

1154

Scientific name: [Zenopsis conchifer](#)

1155




Figure 7.96A. Catch distribution for Silvery John dory.

1156

7.97 White barracudina (*Lussion blanc*) - species code 712 (category LR)

1157

Scientific name: [Arctozenus risso](#)

1158




Figure 7.97A. Catch distribution for White barracudina.

1159

7.98 Atlantic saury (*Balaou atlantique*) - species code 720 (category LR)

1160

Scientific name: [Scomberesox saurus](#)

1161




Figure 7.98A. Catch distribution for Atlantic saury.

1162

7.99 Hatchetfishes (Haches d'argent) - species code 741 (category LR)

1163

Scientific name: [Sternopychidae](#)

1164




Figure 7.99A. Catch distribution for Hatchetfishes.


1165

7.100 Atlantic batfish (*Malthe atlantique*) - species code 742 (category LR)

1166

Scientific name: [Dibranchus atlanticus](#)

1167



1168

7.101 Spottedfin tonguefish (Langue fil noir) - species code 816 (category LR)

1169

Scientific name: [Symphurus diomedeanus](#)

1170




Figure 7.101A. Catch distribution for Spottedfin tonguefish.

1171

7.102 Black dogfish (Aiguillat noir) - species code 221 (category LR)

1172

Scientific name: [Centroscyllium fabricii](#)

1173




Figure 7.102A. Catch distribution for Black dogfish.

1174

7.103 Longfin inshore squid (*Calmar totam*) - species code 4512 (category LR)

1175

Scientific name: [Doryteuthis pealeii](#)

1176




Figure 7.103A. Catch distribution for Longfin inshore squid.

1177

7.104 Red deepsea crab (Crabe rouge) - species code 2532 (category SR)

1178

Scientific name: [Chaceon quinquedens](#)

1179




Figure 7.104A. Catch distribution for Red deepsea crab.

INDEX

1180	Agone atlantique, 121	1226	Atlantic hookear sculpin, 130
1181	Agonidae, 160	1227	Atlantic king crab, 137
1182	Agonidae, 160	1228	Atlantic mackerel, 115
1183	Aspidophoroides monopterygius, 84	1229	Atlantic poacher, 121
1184	Leptagonus decagonus, 121	1230	Atlantic redfishes, 51
1185	Ulcina olrikii, 159	1231	Atlantic rock crab, 135
1186	Aiglefin, 36	1232	Atlantic saury, 180
1187	Aiguillat commun, 102	1233	Atlantic seasnail, 163
1188	Aiguillat noir, 184	1234	Atlantic soft pout, 177
1189	Alewife, 113	1235	Atlantic spiny lumpsucker, 124
1190	Alligatorfish, 84	1236	Atlantic tomcod, 142
1191	Alligatorfishes, 160	1237	Atlantic wolffish, 69
1192	Alosa pseudoharengus, 113	1238	Avocette ruban, 168
1193	Alosa sapidissima, 112	1239	Balaou atlantique, 180
1194	Alose savoureuse, 112	1240	Barndoor skate, 131
1195	Amblyraja radiata, 93	1241	Baudroie d'Amérique, 87
1196	American lobster, 140	1242	Black dogfish, 184
1197	American plaice, 57	1243	Blackbelly rosefish, 118
1198	American shad, 112	1244	Boa dragonfish, 154
1199	Ammodytes dubius, 125	1245	Brosme, 109
1200	Ammodytidae	1246	Brosme brosme, 109
1201	Ammodytes dubius, 125	1247	Brossé améthyste, 153
1202	Anarhichadidae	1248	Callionymidae
1203	Anarhichas denticulatus, 145	1249	Foetorepus agassizii, 175
1204	Anarhichas lupus, 69	1250	Calmar totam, 185
1205	Anarhichas minor, 144	1251	Cancer borealis, 134
1206	Anarhichas denticulatus, 145	1252	Cancer irroratus, 135
1207	Anarhichas lupus, 69	1253	Cancridae
1208	Anarhichas minor, 144	1254	Cancer borealis, 134
1209	Arctic alligatorfish, 159	1255	Cancer irroratus, 135
1210	Arctic eelpout, 176	1256	Capelan, 114
1211	Arctic hookear sculpin, 120	1257	Capelin, 114
1212	Arctic lyre crab, 136	1258	Cardeau à quatre ocelles, 148
1213	Arctozenus risso, 179	1259	Careproctus reinhardti, 166
1214	Argentina silus, 119	1260	Centroscyllium fabricii, 184
1215	Argentinidae	1261	Chabosseau bronzé, 156
1216	Argentina silus, 119	1262	Chabosseau à dix-huit épines, 75
1217	Artediellus atlanticus, 130	1263	Chabosseau à épines courtes, 155
1218	Artediellus uncinatus, 120	1264	Chaceon quinquedens, 186
1219	Aspidophoroides monopterygius, 84	1265	Chionoecetes opilio, 138
1220	Atlantic batfish, 182	1266	Chlorophthalmidae
1221	Atlantic butterfish, 129	1267	Chlorophthalmus agassizi, 152
1222	Atlantic cod, 33	1268	Parasudis truculenta, 150
1223	Atlantic hagfish, 108	1269	Chlorophthalmus agassizi, 152
1224	Atlantic halibut, 54		
1225	Atlantic herring, 72		

1270	Citharichthys arctifrons, 111	1316	Faux-trigle armé, 78
1271	Clupea harengus, 72	1317	Flétan de l'Atlantique, 54
1272	Clupeidae	1318	Flétan noir, 110
1273	Alosa pseudoharengus, 113	1319	Foetorepus agassizii, 175
1274	Alosa sapidissima, 112	1320	Fourbeard rockling, 117
1275	Clupea harengus, 72	1321	Fourline snakeblenny, 173
1276	Coryphaenoides rupestris, 162	1322	Fourspot flounder, 148
1277	Cotte polaire, 157		
1278	Cottidae	1323	Gadidae
1279	Artediellus atlanticus, 130	1324	Gadus morhua, 33
1280	Artediellus uncinatus, 120	1325	Melanogrammus aeglefinus, 36
1281	Icelus spatula, 158	1326	Microgadus tomcod, 142
1282	Myoxocephalus aenaeus, 156	1327	Pollachius virens, 48
1283	Myoxocephalus octodecemspinosis, 73	1328	Gadus morhua, 33
1284	Myoxocephalus scorpius, 155	1329	Gaspareau, 113
1285	Triglops murrayi, 78	1330	Gelatinous snailfish, 164
1286	Cottunculus microps, 157	1331	Geryonidae
1287	Crabe des neiges, 138	1332	Chaceon quinquedens, 186
1288	Crabe Hyas coarctatus, 136	1333	Glyptocephalus cynoglossus, 60
1289	Crabe lyre araignée, 139	1334	Goberge, 48
1290	Crabe rouge, 186	1335	Grande argentine, 119
1291	Crabe épineux du nord, 137	1336	Grande raie, 131
1292	Crevette nordique, 133	1337	Great spider crab, 139
1293	Cryptacanthodes maculatus, 174	1338	Greater argentine, 119
1294	Cryptacanthodidae	1339	Greenland halibut, 110
1295	Cryptacanthodes maculatus, 174	1340	Grenadier de roche, 162
1296	Cunner, 147	1341	Grenadier du Grand Banc, 122
1297	Cusk, 109	1342	Grenadier-scie, 161
1298	Cyclopteridae	1343	Grubby, 156
1299	Cyclopterus lumpus, 123	1344	Gulf Stream flounder, 111
1300	Eumicrotremus spinosus, 124		
1301	Cyclopterus lumpus, 123	1345	Haches d'argent, 181
1302	Cynoglossidae	1346	Haddock, 36
1303	Symphurus diomedeanus, 183	1347	Hameçon atlantique, 130
1304	Daubed shanny, 127	1348	Hameçon neigeux, 120
1305	Dibranchus atlanticus, 182	1349	Hareng de l'Atlantique, 72
1306	Dipturus laevis, 131	1350	Hatchetfishes, 181
1307	Doryteuthis pealeii, 185	1351	Helicolenus dactylopterus, 118
1308	Dragon-boa, 154	1352	Hemitripteridae
1309	Dragonnet tacheté, 175	1353	Hemitripterus americanus, 81
1310	Enchelyopus cimbrius, 117	1354	Hemitripterus americanus, 81
1311	Encornet rouge nordique, 105	1355	Hippoglossina oblonga, 148
1312	Etmopteridae	1356	Hippoglossoides platessoides, 57
1313	Centroscyllium fabricii, 184	1357	Hippoglossus hippoglossus, 54
1314	Eumesogrammus praecisus, 173	1358	Homard américain, 140
1315	Eumicrotremus spinosus, 124	1359	Homarus americanus, 140
		1360	Hyas araneus, 139
		1361	Hyas coarctatus, 136

1362	Hémitriptère atlantique, 81	1407	Loquette d'Amérique, 90
1363	Icelus spatula, 158	1408	Lotidae
1364	lcèle spatulée, 158	1409	Brosme brosme, 109
1365	Illex illecebrosus, 105	1410	Enchelyopus cimbrius, 117
1366	Jonah crab, 134	1411	Loup atlantique, 69
1367	Labridae	1412	Loup tacheté, 144
1368	Tautogolabrus adspersus, 147	1413	Loup à tête large, 145
1369	Lamproie marine, 141	1414	Lumpenus lampretaeformis, 126
1370	Langue fil noir, 183	1415	Lumpfish, 123
1371	Lanternfishes, 151	1416	Lussion blanc, 179
1372	Lançon, 125	1417	Lycenchelys verrillii, 167
1373	Leptagonus decagonus, 121	1418	Lycode arctique, 176
1374	Leptoclinus maculatus, 127	1419	Lycode du Labrador, 169, 170
1375	Leucoraja erinacea, 132	1420	Lycode à carreaux, 128
1376	Leucoraja ocellata, 99	1421	Lycode à tête longue, 167
1377	Limace atlantique, 163	1422	Lycodes lavalaei, 170
1378	Limace gélatineuse, 164	1423	Lycodes reticulatus, 176
1379	Limace marbée, 165	1424	Lycodes terraenovae, 169
1380	Limanda ferruginea, 63	1425	Lycodes vahlii, 128
1381	Limande à queue jaune, 63	1426	Macrouridae
1382	Limande-plie rouge, 66	1427	Coryphaenoides rupestris, 162
1383	Liparidae	1428	Nezumia bairdii, 122
1384	Careproctus reinhardtii, 166	1429	Trachyrincus murrayi, 161
1385	Liparis atlanticus, 163	1430	Malacoraja senta, 96
1386	Liparis fabricii, 164	1431	Mallotus villosus, 114
1387	Liparis gibbus, 165	1432	Malthe atlantique, 182
1388	Liparis atlanticus, 163	1433	Maquereau commun, 115
1389	Liparis fabricii, 164	1434	Marlin-spike grenadier, 122
1390	Liparis gibbus, 165	1435	Maurolicus muelleri, 153
1391	Lithodes maja, 137	1436	Melanogrammus aeglefinus, 36
1392	Lithodidae	1437	Melanostigma atlanticum, 177
1393	Lithodes maja, 137	1438	Merlu argenté, 45
1394	Little skate, 132	1439	Merlu argenté du large, 143
1395	Lolinidae	1440	Merlucciidae
1396	Doryteuthis pealeii, 185	1441	Merluccius albidus, 143
1397	Lompe, 123	1442	Merluccius bilinearis, 45
1398	Lompénie tachetée, 127	1443	Merluccius albidus, 143
1399	Lompénie-serpent, 126	1444	Merluccius bilinearis, 45
1400	Longfin hake, 116	1445	Merluche blanche, 39
1401	Longfin inshore squid, 185	1446	Merluche à longues nageoires, 116
1402	Longhorn sculpin, 75	1447	Merluche écureuil, 42
1403	Longnose greeneye, 150	1448	Microgadus tomcod, 142
1404	Lophiidae	1449	Molasse atlantique, 177
1405	Lophius americanus, 87	1450	Monkfish, 87
1406	Lophius americanus, 87	1451	Morue franche, 33
		1452	Motelle à quatre barbillons, 117
		1453	Moustache sculpin, 78

1454	Myctophidae, 151	1499	Petite poule de mer atlantique, 124
1455	Myctophidae, 151	1500	Petromyzon marinus, 141
1456	Myoxocephalus aenaeus, 156	1501	Petromyzontidae
1457	Myoxocephalus octodecemspinulosus, 75	1502	Petromyzon marinus, 141
1458	Myoxocephalus scorpius, 155	1503	Pholidae
1459	Myxine du nord, 108	1504	Pholis gunnellus, 171
1460	Myxine glutinosa, 108	1505	Pholis gunnellus, 171
1461	Myxinidae	1506	Phycidae
1462	Myxine glutinosa, 108	1507	Phycis chesteri, 116
1463	Nemichthyidae	1508	Urophycis chuss, 42
1464	Nemichthys scolopaceus, 168	1509	Urophycis tenuis, 39
1465	Nemichthys scolopaceus, 168	1510	Phycis chesteri, 116
1466	Nephropidae	1511	Picked dogfish, 102
1467	Homarus americanus, 140	1512	Pleuronectidae
1468	Newfoundland eelpout, 169, 170	1513	Glyptocephalus cynoglossus, 60
1469	Nezumia bairdii, 122	1514	Hippoglossoides platessoides, 57
1470	Northern prawn, 133	1515	Hippoglossus hippoglossus, 54
1471	Northern shortfin squid, 105	1516	Limanda ferruginea, 63
1472	Northern wolffish, 145	1517	Pseudopleuronectes americanus, 66
1473	Ocean pout, 90	1518	Reinhardtius hippoglossoides, 110
1474	Oeil-vert à long nez, 150	1519	Plie canadienne, 57
1475	Offshore silver hake, 143	1520	Plie du Gulf Stream, 111
1476	Ogcocephalidae	1521	Plie grise, 60
1477	Dibranchus atlanticus, 182	1522	Poisson-alligator arctique, 159
1478	Ommastrephidae	1523	Poisson-alligator atlantique, 84
1479	Illex illecebrosus, 105	1524	Poissons-alligator, 160
1480	Oregoniidae	1525	Poissons-lanternes, 151
1481	Chionoecetes opilio, 138	1526	Polar sculpin, 157
1482	Hyas araneus, 139	1527	Pollachius virens, 48
1483	Hyas coarctatus, 136	1528	Pollock, 48
1484	Osmeridae	1529	Poulamon atlantique, 142
1485	Mallotus villosus, 114	1530	Pseudopleuronectes americanus, 66
1486	Osmerus mordax, 146	1531	Psychrolutidae
1487	Osmerus mordax, 146	1532	Cottunculus microps, 157
1488	Pandalidae	1533	Quatre-lignes atlantique, 173
1489	Pandalus borealis, 133	1534	Queen crab, 138
1490	Pandalus borealis, 133	1535	Radiated shanny, 172
1491	Paralepididae	1536	Raie hérisson, 132
1492	Arctozenus risso, 179	1537	Raie lisse, 96
1493	Paralichthyidae	1538	Raie tachetée, 99
1494	Citharichthys arctifrons, 111	1539	Raie épineuse, 93
1495	Hippoglossina oblonga, 148	1540	Rainbow smelt, 146
1496	Parasudis triculenta, 150	1541	Rajidae
1497	Peprilus triacanthus, 129	1542	Amblyraja radiata, 93
1498	Petite limace de mer, 166	1543	Dipturus laevis, 131
		1544	Leucoraja erinacea, 132

1545	Leucoraja ocellata, 99	1592	Leptoclinus maculatus, 127
1546	Malacoraja senta, 96	1593	Lumpenus lampretaeformis, 126
1547	Red deepsea crab, 186	1594	Ulvaria subbifurcata, 172
1548	Red hake, 42	1595	Stomias boa, 154
1549	Reinhardtius hippoglossoides, 110	1596	Stomiidae
1550	Rock gunnel, 171	1597	Stomias boa, 154
1551	Roughnose grenadier, 161	1598	Stromateidae
1552	Roundnose grenadier, 162	1599	Peprilus triacanthus, 129
1553	Saint Pierre argenté, 178	1600	Stromaté fossette, 129
1554	Sand lance, 125	1601	Syphurus diomedeanus, 183
1555	Scomber scombrus, 115	1602	Sébaste chèvre, 118
1556	Scomberesocidae	1603	Sébastes de l'Atlantique, 51
1557	Scomberesox saurus, 180	1604	Tanche-tautogue, 147
1558	Scomberesox saurus, 180	1605	Tautogolabrus adspersus, 147
1559	Scombridae	1606	Terrassier tacheté, 174
1560	Scomber scombrus, 115	1607	Thorny skate, 93
1561	Scophthalmidae	1608	Tourteau jona, 134
1562	Scophthalmus aquosus, 149	1609	Tourteau poïnclos, 135
1563	Scophthalmus aquosus, 149	1610	Trachyrincus murrayi, 161
1564	Sea lamprey, 141	1611	Triglops murrayi, 78
1565	Sea raven, 81	1612	Turbot de sable, 149
1566	Sea tadpole, 166		
1567	Sebastes, 51	1613	Ulcina olrikii, 159
1568	Sebastidae	1614	Ulvaire deux-lignes, 172
1569	Helicolenus dactylopterus, 118	1615	Ulvaria subbifurcata, 172
1570	Sebastes, 51	1616	Urophycis chuss, 42
1571	Shorthorn sculpin, 155	1617	Urophycis tenuis, 39
1572	Shortnose greeneye, 152		
1573	Sigouine de roche, 171	1618	Vahl's eelpout, 128
1574	Silver hake, 45	1619	Variegated snailfish, 165
1575	Silvery John dory, 178	1620	White barracudina, 179
1576	Silvery lightfish, 153	1621	White hake, 39
1577	Slender snipe eel, 168	1622	Windowpane flounder, 149
1578	Smooth skate, 96	1623	Winter flounder, 66
1579	Snakeblenny, 126	1624	Winter skate, 99
1580	Spatulate sculpin, 158	1625	Witch flounder, 60
1581	Spotfin dragonet, 175	1626	Wolf eelpout, 167
1582	Spotted wolffish, 144	1627	Wrymouth, 174
1583	Spottedfin tonguefish, 183		
1584	Squalidae	1628	Yellowtail flounder, 63
1585	Squalus acanthias, 102		
1586	Squalus acanthias, 102	1629	Zeidae
1587	Sternopychidae, 181	1630	Zenopsis conchifer, 178
1588	Maurolicus muelleri, 153	1631	Zenopsis conchifer, 178
1589	Sternopychidae, 181	1632	Zoarces americanus, 90
1590	Stichaeidae	1633	Zoarcidae
1591	Eumesogrammus praecisus, 173	1634	Lycenchelys verrillii, 167

- 1635 Lycodes lavalaei, 170
 - 1636 Lycodes reticulatus, 176
 - 1637 Lycodes terraenovae, 169
 - 1638 Lycodes vahlii, 128
 - 1639 Melanostigma atlanticum, 177
 - 1640 Zoarces americanus, 90
-
- 1641 Éperlan arc-en-ciel, 146
 - 1642 Éperlan du large, 152