

THESIS TITLE
SECOND LINE IF NECESSARY

by

DILRAJ GHUMAN

A thesis submitted to the
Department of Physics, Engineering Physics and Astronomy
in conformity with the requirements for
the degree of Master of Science

Queen's University
Kingston, Ontario, Canada

May 2021

Copyright © Dilraj Ghuman, 2021

Abstract

This is my abstract.

Acknowledgments

Blah blah blah.

Statement of Originality

Contents

Abstract	i
Acknowledgments	ii
Statement of Originality	iii
Contents	iv
List of Tables	vi
List of Figures	vii
Chapter 1: Introduction	1
1.1 Neutrinos	1
1.1.1 Oscillations	2
1.1.2 Interactions	5
1.1.3 Production & Sources	6
Chapter 2: Background	7
2.1 Detection Techniques	7
2.2 Neutrino Telescopes	7
2.2.1 IceCube	7
2.2.2 ANTARES	7
2.2.3 KM3NET	7
Chapter 3: The Pacific Ocean Neutrino Explorer	8
3.1 Detectors	8
3.1.1 Geometry	8
3.2 Ocean Networks Canada	8
Chapter 4: Simulation	9
4.1 IceCube Framework	9

4.2	Simulating Neutrinos	9
4.3	Simulating Muons	9
4.4	Detector Response	9
Chapter 5:	Reconstruction	10
5.1	Linefit	10
5.2	Likelihood	10
Chapter 6:	Results	11
6.1	Likelihood	11
Chapter 7:	Summary and Conclusions	12
7.1	Summary	12
7.2	Future Work	12
7.3	Conclusion	12
	Bibliography	13

List of Tables

List of Figures

1.1	The Feynmann diagrams for the vertices that would be included in neutrino interactions using the charged W^\pm boson on the left and the neutral Z^0 boson on the right.	5
-----	--	---

Chapter 1

Introduction

The cosmic sky has entranced humans for as far as recorded history can trace. As technology evolved, so too did the observation of the universe around us; from the naked eye to primitive telescopes, and eventually to present day space telescopes, like the Hubble Space Telescope and the upcoming James Web Space Telescope (**NEED TO CITE THESE**). These growing technological leaps have also resulted in the exploration of the incredibly small and eventually resulted in the discovery of the neutrino [15]. It was perhaps inevitable that these two seemingly separate areas of physics would eventually meet.

1.1 Neutrinos

The neutrino is a fundamental particle first proposed by Wolfgang Pauli [3], and then later discovered in 1956 using the byproducts of β^- decay [15]. As research continued into the elusive neutrino, another flavour of neutrino was discovered in 1962 called the muon neutrino (ν_μ) [5] and eventually the final flavour of the tau neutrino (ν_τ) [12].

1.1. NEUTRINOS

1.1.1 Oscillations

Alongside the discovery of the neutrino and their flavours, another problem arose in the field of neutrino physics: the solar neutrino problem [13]. During the 1960's, an experiment was proposed by Bahcall and Davis to measure the solar neutrino flux, referred to as the Homestake experiment [6, 1]. This was a tank of ^{37}Cl , built underground to avoid cosmic backgrounds, and used the simple reaction [6, 1]



to measure the expected solar neutrino flux from the sun.

Solar neutrinos originate from nuclear processes that occur in the sun, such as the PP chain, or the CNO cycle, and can be detected in experiments on Earth [10]. Depending on the energy and process in producing the neutrino, we can expect to detect particular flavours of neutrinos in experiments like the Homestake experiment. In particular, using the predicted distribution of the internal electron density of the Sun, and a spectrum of the produced electron flavours, one could predict the expected flux of solar neutrinos [10]. In particular, one could predict the influx of electron neutrinos, as was exactly done for the Homestake experiment. It was found that the measured flux was consistently around 30% the theoretical amount [6, 1, 10], and hence was coined the solar neutrino problem.

Neutrino detectors continued to be constructed to research and understand these fundamental particles, such as Super-Kamiokande [17], the collaboration of Kamiokande [11] and the IMB [4] experiments. Another class of solar neutrino detectors were those

1.1. NEUTRINOS

that used the Gallium chain

$$\nu_e + {}^{71}\text{Ga} \rightarrow e^- + {}^{71}\text{Ge}, \quad (1.2)$$

such as GALLEX [16]. Regardless, the same issue persisted as there continued to be a distinct dissonance between the theoretical expectations of solar neutrinos and the observed experimental results. That was, until the Sudbury Neutrino Observatory (SNO) made a distinct change in their approach to solar neutrino detection compared to predecessors by using heavy water [2]. This allowed for the following interactions [2],

$$\nu_e + d \rightarrow p + p + e^- \quad (1.3)$$

$$\nu_l + d \rightarrow p + n + \nu_l \quad (1.4)$$

where we have the Charged Current (CC) interaction in equation 1.3 and the Neutral Current (NC) interaction in equation 1.4. This meant that all flavours of neutrinos could be detected, and using it to detect solar neutrinos showed the theoretical flux originally predicted [2].

This result had an astounding implication; the neutrinos were changing on their journey from the Sun [2]. In the Standard Model, all the neutrino flavours have masses that are identically zero, and this would mean that there is no possible way for the neutrinos to somehow change flavour on their journey to the detectors [10]. Clearly there was a change in flavour, and thus the Standard Model must be incorrect about the masses of the neutrinos.

The classic demonstrative method to see this is to consider the mixing of two

1.1. NEUTRINOS

neutrino flavours (like ν_μ and ν_e) [10]. In analogy to quark flavour mixing [7], we know the mixing of the flavours occurs in the transformation from the mass to the flavour basis. In particular, for two mass and flavour states one can find [10],

$$P(\nu_e \rightarrow \nu_\mu, ct) = \sin^2 2\theta \sin^2 \left(\frac{\pi ct}{L} \right) \quad (1.5)$$

where θ is the mixing between the two flavour states, $L = \frac{4\pi E}{\Delta m^2}$ is the vacuum oscillation length, and $\Delta m^2 = m_2^2 - m_1^2$. Here it is easy enough to see that the oscillation probability vanishes if the masses are identical, and this naturally extends into the three flavour case. The vacuum oscillation length, L , is an important and useful quantity as it describes the distance a neutrino must travel before an oscillation is expected [10]. Experiments like the long-baseline neutrino oscillation experiment Tokai-to-Kamioka (T2K) attempt to use this length to probe the mixing angles of the three neutrino flavours.

Similar to the CKM matrix for quark mixing [7], the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix [14] gives a relation between the mass and flavour states:

$$\begin{bmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{bmatrix} = U \begin{bmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{bmatrix} \quad (1.6)$$

1.1. NEUTRINOS




Figure 1.1: The Feynmann diagrams for the vertices that would be included in neutrino interactions using the charged W^\pm boson on the left and the neutral Z^0 boson on the right.

and we see that

$$U = \begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{\text{CP}}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{\text{CP}}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{\text{CP}}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{\text{CP}}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{\text{CP}}} & c_{23}c_{13} \end{bmatrix} \begin{bmatrix} e^{i\eta_1} & 0 & 0 \\ 0 & e^{i\eta_2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (1.7)$$

where $s_{ij} = \sin \theta_{ij}$, $c_{ij} = \cos \theta_{ij}$, δ_{CP} is the Charge-Parity violation phase [14], and η_i are the Majorana phases. If neutrinos are not their own anti-particles, or in other words are Dirac fermions, we can expect $\eta_i = 0$. If they are their own anti-particles, also known as Majorana, then the phases η_i play a more important role [9].

1.1.2 Interactions

Neutrinos are neutral and interact only through the Weak interaction. The Weak interaction is a force that is mediated by the W^\pm and Z^0 massive bosons, and is the force responsible for decays. The main vertices involved in neutrino interactions are shown in Figure 1.1, where the interacting lepton corresponds with the interacting neutrino flavour.

All interaction involving neutrino production or detection utilize these vertices in

1.1. NEUTRINOS

some shape or form. We refer to interactions that use the W^\pm boson as the Charged Current (CC) interaction [8], and those that use the Z^0 boson as being Neutral Current (NC) interactions [8].

It is natural to notice that these interactions require something to interact with, or in other words, the neutrinos must propagate through non-vacuum media and hit targets. We have up to this point only considered oscillations of neutrinos in vacuum, and another important aspect is to consider the effect interactions could have on these oscillations. In particular, it is noted that certain flavours of neutrinos can be more strongly influenced by media than others [10]. In particular, electron neutrinos (ν_e) being first generation and majority of regular everyday matter being first generation would result in this stronger coupling [10]. This difference in coupling would result in changes in the oscillation that could be complex.

1.1.3 Production & Sources

Solar: Neutrinos produced from the fusion process inside the sun. Go into detail about the process.

Atmospheric: Cosmic rays that interact with the atmosphere produce neutrinos that then shower the earth. Go into detail about process

Geo: Rare decays in the crust eject neutrinos which can be detected.

Active Galactic Nucleus: Active centres of galaxies that are hubs for producing high energy neutrinos

Supernova: Massive Explosions of dying stars that eject high energy neutrinos.

Chapter 2

Background

2.1 Detection Techniques

Literature dive into different types of neutrino detection techniques.

Cherenkov indirect, ect..?

2.2 Neutrino Telescopes

Generally use Cherenkov radiation as a method of detecting high energy neutrinos from cosmic sources.

2.2.1 IceCube

2.2.2 ANTARES

2.2.3 KM3NET

Chapter 3

The Pacific Ocean Neutrino Explorer

3.1 Detectors

3.1.1 Geometry

3.2 Ocean Networks Canada

Chapter 4

Simulation

- 4.1 IceCube Framework
- 4.2 Simulating Neutrinos
- 4.3 Simulating Muons
- 4.4 Detector Response

Chapter 5

Reconstruction

5.1 Linefit

5.2 Likelihood

Chapter 6

Results

6.1 Likelihood

Chapter 7

Summary and Conclusions

7.1 Summary

7.2 Future Work

7.3 Conclusion

BIBLIOGRAPHY

Bibliography

- [1] John N. Bahcall. Solar neutrinos. i. theoretical. *Phys. Rev. Lett.*, 12:300–302, Mar 1964.
- [2] A. Bellerive, J.R. Klein, A.B. McDonald, A.J. Noble, and A.W.P. Poon. The sudbury neutrino observatory. *Nuclear Physics B*, 908:30–51, Jul 2016.
- [3] Laurie M. Brown. The idea of the neutrino. *Physics Today*, 31(9):23–28, September 1978.
- [4] D. Casper, R. Becker-Szendy, C. B. Bratton, D. R. Cady, R. Claus, S. T. Dye, W. Gajewski, M. Goldhaber, T. J. Haines, P. G. Halverson, T. W. Jones, D. Kielczewska, W. R. Kropp, J. G. Learned, J. M. LoSecco, C. McGrew, S. Matsuno, J. Matthews, M. S. Mudan, L. Price, F. Reines, J. Schultz, D. Sinclair, H. W. Sobel, J. L. Stone, L. R. Sulak, R. Svoboda, G. Thornton, and J. C. van der Velde. Measurement of atmospheric neutrino composition with the imb-3 detector. *Phys. Rev. Lett.*, 66:2561–2564, May 1991.
- [5] G. Danby, J. Gaillard, K. Goulianos, L. Lederman, N. Mistry, M. Schwartz, and J. Steinberger. Observation of high-energy neutrino reactions and the existence of two kinds of neutrinos. *Physical Review Letters*, 9:36–44, 1962.

BIBLIOGRAPHY

- [6] Raymond Davis. Solar neutrinos. ii. experimental. *Phys. Rev. Lett.*, 12:303–305, Mar 1964.
- [7] W.J. Marciano E. Blucher. The cabibbo angle and ckm unitarity. <https://pdg.lbl.gov/2020/reviews/rpp2020-rev-vud-vus.pdf>.
- [8] Carlotta Giusti and Martin V Ivanov. Neutral current neutrino-nucleus scattering: theory. *Journal of Physics G: Nuclear and Particle Physics*, 47(2):024001, Jan 2020.
- [9] Particle Data Group and et al. Zyla. Review of Particle Physics. *Progress of Theoretical and Experimental Physics*, 2020(8), 08 2020. 083C01.
- [10] Mike Guidry and Jay Billings. A basic introduction to the physics of solar neutrinos, 2018.
- [11] K. S. Hirata et al. Observation of a small atmospheric muon-neutrino / electron-neutrino ratio in Kamiokande. *Phys. Lett. B*, 280:146–152, 1992.
- [12] K. Kodama, N. Ushida, C. Andreopoulos, N. Saoulidou, G. Tzanakos, P. Yager, B. Baller, D. Boehnlein, W. Freeman, B. Lundberg, and et al. Observation of tau neutrino interactions. *Physics Letters B*, 504(3):218–224, Apr 2001.
- [13] Andrew John Lowe. Neutrino physics & the solar neutrino problem, 2009.
- [14] Ziro Maki, Masami Nakagawa, and Shoichi Sakata. Remarks on the Unified Model of Elementary Particles. *Progress of Theoretical Physics*, 28(5):870–880, 11 1962.

BIBLIOGRAPHY

- [15] F. Reines, C. L. Cowan, F. B. Harrison, A. D. McGuire, and H. W. Kruse. Detection of the free antineutrino. *Phys. Rev.*, 117:159–173, Jan 1960.
- [16] D. Vignaud. The gallex solar neutrino experiment. *Nuclear Physics B - Proceedings Supplements*, 60(3):20–29, 1998.
- [17] Christopher W. Walter. The super-kamiokande experiment. *Neutrino Oscillations*, page 19–43, Mar 2008.