Result 1 (direct). [Callable evidence, direct assumption]
Let n be a natural number. Assume (for every $x : \mathbb{N},\ x = x$) (`h`). Let $p = h(n)$. Then

$$p = p.$$

Result 2 (named). [Callable evidence, named assumption]
Let n be a natural number. Assume reflexive() (`h`). Let $p = h(n)$. Then

$$p = p.$$

Result 3 (nested). [Callable evidence, nested assumption]
Let n and m be natural numbers. Assume (for every $x : \mathbb{N},$ (for every $y : \mathbb{N},\ x + y = y + x$)) (`h`). Let $k = h(n)$. Let $p = k(m)$. Then

$$p = p.$$

Result 4 (projected). [Callable evidence, projected universal evidence]
Let w be a family of proofs that every element of $\mathbb{N}$ equals itself. Let n be a natural number. Let $p = w.\mathrm{all}(n)$. Then

$$p = p.$$

Result 5 (proof domain). [Callable evidence, universal evidence as an argument]
Assume (for every $h :$ (for every $x : \mathbb{N},\ x = x$) , (for every $n : \mathbb{N},\ n = n$)) (`r`). Assume (for every $x : \mathbb{N},\ x = x$) (`h`). Let n be a natural number. Let $k = r(h)$. Let $p = k(n)$. Then

$$p = p.$$

Result 6 (function domain). [Callable evidence, proof family as an argument]
Assume (for every $h : \prod_{x:\mathbb{N}} x = x,$ (for every $n : \mathbb{N},\ n = n$)) (`r`). Assume (for every $x : \mathbb{N},\ x = x$) (`h`). Let n be a natural number. Let $k = r(h)$. Let $p = k(n)$. Then

$$p = p.$$

Result 7 (captured type). [Callable evidence, type scope preservation]
Assume (for every $x : \mathbb{R},\ x = x$) (`h`). Let z be a real number. Let *Real* be a type. Let $p = h(z)$. Then

$$p = p.$$

Result 8 (captured free). [Callable evidence, free variable preservation]
Let n be a natural number. Assume (for every $x : \mathbb{N}$, (for every $y : \mathbb{N}$, $x + y = n$)) (`h`). Let y be a natural number. Let $k = h(y)$. Let $p = k(n)$. Then

$$p = p.$$

Result 9 (implication). [Callable evidence, implication application]
Let n be a natural number. Assume $n = 0$ implies $n = n$ (`h`). Assume $n = 0$ (`p`). Let $q = h(p)$. Then

$$q = q.$$

Result 10 (saved assumption). [Callable evidence, local evidence alias]
Assume (for every $x : \mathbb{N}$, $x = x$) (`all`). Let $h = all$. Let n be a natural number. Let $p = h(n)$. Then

$$p = p.$$

Result 11 (local precedence). [Callable evidence, callable local precedence]
Let n be a natural number. Assume (for every $x : \mathbb{N}$, $x = x$) (`evidence`). Let $p = evidence(n)$. Then

$$p = p.$$

Result 12 (global fallback). [Callable evidence, noncallable local preserves module resolution]
Let n be a natural number. Assume $n = n$ (`evidence`). Let $p = e(n)$. Then

$$p = p.$$

Result 13 (inferred carrier). [Callable evidence, inferred condition carrier]
Let T be a type. Let f and g be functions from T to T. Let x be an element of T. Assume pointwise equal(f, g) (`h`). Let $p = h(x)$. Then

$$p = p.$$

Result 14 (existential result). [Callable evidence, existential proof as result]
Assume (for every $x : \mathbb{N}$, (there exists $y : \mathbb{N}$, $x = y$)) (`h`). Let n be a natural number. Let $p = h(n)$. Then

$$p = p.$$

Result 15 (nested function domain). [Callable evidence, nested proof-family argument]
Let n be a natural number. Assume $\Big($for every $h : \prod_{x:\mathbb{N}} \prod_{y:\mathbb{N}} x + y = y + x,\ n = n\Big)$ (`r`). Assume (for every $x : \mathbb{N}$, (for every $y : \mathbb{N}$, $x + y = y + x$)) (`h`). Let $p = r(h)$. Then

$$p = p.$$

LLM-assisted language specimens and proofs (gpt-6, 2026-11-07). See PROVENANCE.md. These statements validate scoped callable evidence.