Result 1 (basis at a restricted vertex). [Clean topology study, basis at a restricted vertex]
Let K be a type. Let *field* be a field structure on K. Let M be a module over K. Let I be a type. Let n, a and m be natural numbers. Assume $a+m \leq n$ (`bound`). Let L be a family of ordered interval labels indexed by I on vertices 0 through n. Let v be a vertex in $\{0, \ldots, m+1-1\}$. Let b be a basis of M indexed by $A(L, a+v)$. Let *vectors* be a function from I to M. Assume (for every $x : A(L, a+v)$, $b_x = \mathit{vectors}(x)$) (`realizes`). Then

$$(\text{there exists } c : \mathrm{Basis}_K(M, R(L, v)),\ (\text{for every } y : R(L, v),\ c_y = \mathit{vectors}(y)))\,.$$

Result 2 (interval basis restricts). [Clean topology study, interval basis restricts]
Let K be a type. Let *field* be a field structure on K. Let n, a and m be natural numbers. Assume $a + m \leq n$ (`bound`). Let d be a function from $\{0, \ldots, n-1\}$ to $\{\text{false}, \text{true}\}$. Assume subpath sources preserved$(n, a, m, \mathit{bound}, d)$ (`sources`). Assume subpath targets preserved$(n, a, m, \mathit{bound}, d)$ (`targets`). Let Z be a module diagram on Q_d. Let I be a type. Let L be a family of ordered interval labels indexed by I on vertices 0 through n. Let B be an interval basis of Z with labels L. Then

$$\big(\text{there exists } C : \mathrm{IBasis}(Z|_{[a,a+m]}, L|_{[a,a+m]}),\ C.\mathrm{vectors} = B.\mathrm{vectors}|_{[a,a+m]}\big)\,.$$

LLM-assisted statement and proof (gpt-6, 2026-11-07). See PROVENANCE.md. This study addresses pointwise basis transport and full interval-basis restriction. The paper's pyramid theorem remains a separate proof obligation.