Result 1 (point equality). [Carrier bundle study, equality reflexivity]
Let P be a pointed topological space. Let x be an element of P.carrier. Then

$$x = x.$$

Result 2 (constant map continuous). [Carrier bundle study, Mathlib continuous_const]
Let P be a pointed topological space. Then

$$c_P \text{ is continuous from } P \text{ to } P.$$

Result 3 (carrier alias). [Carrier bundle study, authored carrier reduction]
Let P be a pointed topological space. Let x be an element of $|P|$. Then

$$(y \mapsto y)\,(x) = x.$$

Result 4 (constructor point). [Carrier bundle study, constructor projection]
Let T be a type. Let t be a topology on T. Let x be an element of T. Then

$$\text{pointed space}(T, t, x).\text{point} = x.$$

Result 5 (selected point projection). [Carrier bundle study, dependent return substitution]
Let Q be a pointed topological space. Then

$$p_Q = Q.\text{point}.$$

Result 6 (family evaluation). [Carrier bundle study, dependent family evaluation]
Let P be a pointed topological space. Then

$$p.\text{value}(P) = P.\text{point}.$$

Isolated carrier-field prototype. LLM-assisted statements and proofs (gpt-6, 2026-11-07); see PROVENANCE.md. Main Clean does not yet accept carrier fields.