Result 1 (addition with one). [Mathlib]
Let x be an element of $\mathbb{R}$. Then

$$x + (1 \,:\, \mathbb{R}) = (1 \,:\, \mathbb{R}) + x.$$

Result 2 (named numeral assumption). [Numeral study, identity]
Assume $0 \leq (001 \,:\, \mathbb{R})$ (`nonnegative((1: Real))`). Then

$$0 \leq (1 \,:\, \mathbb{R}).$$

Result 3 (inferred real addition). [Mathlib]
Let x be an element of $\mathbb{R}$. Then

$$x + 1 = 1 + x.$$

Result 4 (inferred nested addition). [Mathlib]
Let x be an element of $\mathbb{R}$. Then

$$1 + 2 + x = x + (1 + 2).$$

Result 5 (inferred counting addition). [Mathlib]
Let n be an element of $\mathbb{N}$. Then

$$n + 1 = 1 + n.$$

Result 6 (inferred named assumption). [Numeral study, contextual assumption]
Assume $0 \leq 1$ (`nonnegative(1)`). Then

$$0 \leq 001.$$

LLM-assisted prototype examples (gpt-6, 2026-11-07). Explicit numeral annotations remain visible; inferred numerals print as ordinary digits. See PROVENANCE.md.